arXiv:1009.1648v1 [math.SG] 8 Sep 2010

LAGRANGIAN FLOER THEORY AND MIRROR SYMMETRY ON COMPACT TORIC MANIFOLDS

KENJI FUKAYA, YONG-GEUN OH, HIROSHI OHTA, KAORU ONO

Abstract. In this paper we study Lagrangian Floer theory on toric manifolds from the point of view of mirror symmetry. We construct a natural isomorphism between the Frobenius manifold structures of the (big) quantum cohomology of the toric manifold and of Saito's theory of singularities of the potential function constructed in [FOOO5] via the Floer cohomology deformed by ambient cycles. Our proof of the isomorphism involves the open-closed Gromov-Witten theory of one-loop.

Date: Sep. 8, 2010.

Key words and phrases. Floer cohomology, mirror symmetry, toric manifolds, open-closed Gromov-Witten invariant, Saito theory, Landau-Ginzburg model, weakly unobstructed Lagrangian submanifolds, potential function, Jacobian ring, Frobenius structure.

KF is supported partially by JSPS Grant-in-Aid for Scientific Research No.19104001 and Global COE program G08, YO by US NSF grant # 0904197, HO by JSPS Grant-in-Aid for Scientific Research No.19340017, and KO by JSPS Grant-in-Aid for Scientific Research, Nos. 21244002 and 18340014.

Contents

1. Introduction 4
2. Statement of the results 11
3. Projective space: an example 24

Part 1. Ring isomorphism 27

4. Norms and completions of polynomial ring over Novikov ring 27
5. Localization of Jacobian ring at moment polytope 30
6. Operator $\mathfrak{q}$: review 33
7. Well-definedness of Kodaira-Spencer map 41
8. Well-definedness of potential function 49
9. Kodaira-Spencer map is a ring homomorphism 53
10. Surjectivity of Kodaira-Spencer map 68
11. Versality of the potential function with bulk 72
12. Algebraization of Jacobian ring 73
13. Seidel homomorphism and a result by McDuff-Tolman 78
14. Injectivity of Kodaira-Spencer map 81
14.1. The case $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$ 81
14.2. The case $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ 94
15. The Chern class c_1 and critical values of $\mathfrak{PO}_{\mathfrak{b}}$ 100
16. Hirzebruch surface F_2: an example 101

Part 2. Coincidence of pairings 104

17. Operator $\mathfrak{p}$ and Poincaré duality 104
18. Cyclic symmetry in the toric case 109
19. Operator $\mathfrak{p}$ in the toric case 118
20. Moduli space of holomorphic annuli 123
21. Poincaré duality is residue pairing 141
22. Clifford algebra and Hessian matrix 142
23. Residue pairing and Hessian determinant 146
24. Cyclic homology and variation of the invariant Z 150
25. Orientation 159
25.1. Operators D_1 and D_2 159
25.2. Orientation of Index D_1 161
25.3. Orientation of Index D_2 162
25.4. Continuation of linear Kuranishi models 162
26. Sign in Theorem 20.1 and Proposition 21.2 166
26.1. Some lemmas 166
26.2. Proofs of signs in Theorem 20.1 and Proposition 21.2 170

Part 3. Appendix 176

27. Appendix 1: Coincidence of the two definitions of $\delta^{\mathfrak{b},b}$ 176
28. Appendix 2: Interpolation between component-wise and T^n equivariant Kuranishi structures 180
29. Appendix 3: T^n equivariant and cyclically symmetric Kuranishi structure 185
29.1. The case with one disc component 190
29.2. The case with more than one disk components 210

30. Appendix 4: T^n equivariant continuous family of multisections 220
31. Appendix 5: Hochschild and quantum cohomologies 223
References 229

1. Introduction

The purpose of this paper is to prove a version of mirror symmetry between compact toric A-model and Landau-Ginzburg B-model.

Let X be a compact toric manifold and take a toric Kähler structure on it. In this paper X is not necessarily assumed to be Fano. In [FOOO5] we defined a potential function with bulk, $\mathfrak{PO}_{\mathfrak b}$, which is a family of functions of n variables $y_1,\dots,y_n$ parameterized by the cohomology class $\mathfrak b \in H(X;\Lambda_0)$. (More precisely, it is parameterized by T^n-invariant cycles. We explain this point later in Sections 2 and 8.) (Λ_0 is defined in Definition 1.6.)

$\mathfrak{PO}_{\mathfrak b}$ is an element of an appropriate completion of $\Lambda_0[y_1,y_1^{-1}\dots,y_n,y_n^{-1}]$, the Laurent polynomial ring over (universal) Novikov ring Λ_0. We denote this completion by $\Lambda_0^{\overset{\circ}{P}}\langle\langle y,y^{-1}\rangle\rangle$. (See Definition 2.1 for its definition.) We put

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle y,y^{-1}\rangle\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i\frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i=1,\dots,n\right)}.$$

Differentiation of $\mathfrak{PO}_{\mathfrak b}$ with respect to the ambient cohomology class $\mathfrak b$ gives rise to a Λ_0 module homomorphism

$$\mathfrak{ks}_{\mathfrak b} : H(X;\Lambda_0) \to \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}). \tag{1.1}$$

We define a quantum ring structure $\cup^{\mathfrak b}$ on $H(X;\Lambda_0)$ by deforming the cup product by $\mathfrak b$ using Gromov-Witten theory. (See Definition 2.26.) The main result of this paper can be stated as follows:

Theorem 1.1. *Equip $H(X;\Lambda_0)$ with the ring structure $\cup^{\mathfrak b}$. Then*

1. *The homomorphism $\mathfrak{ks}_{\mathfrak b}$ in (1.1) is a ring isomorphism.*
2. *If $\mathfrak{PO}_{\mathfrak b}$ is a Morse function, i.e., all its critical points are nondegenerate, the isomorphism $\mathfrak{ks}_{\mathfrak b}$ above intertwines the Poincaré duality pairing on $H(X;\Lambda)$ and the residue pairing on $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \otimes_{\Lambda_0} \Lambda$.*

We refer readers to Section 2 for the definition of various notions appearing in Theorem 1.1. Especially the definition of residue pairing which we use in this paper is given in Definition 2.23 and Theorem 2.24.

We now explain how Theorem 1.1 can be regarded as a mirror symmetry between toric A-model and Landau-Ginzburg B-model:

(a) First the surjectivity of (1.1) implies that the family $\mathfrak{PO}_{\mathfrak b}$ is a versal family in the sense of deformation theory of singularities. (See Theorem 11.1.)

(b) Therefore the tangent space $T_{\mathfrak b}H(X;\Lambda_0)$ carries a ring structure pulled-back from the Jacobian ring by (1.1). Theorem 1.1 (1) implies that this pull-back coincides with the quantum ring structure $\cup^{\mathfrak b}$.

(c) Theorem 1.1 (2) implies that the residue paring, which is defined when $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$ is a Morse function, can be extended to arbitrary class $\mathfrak b \in H(X;\Lambda_0)$. In fact the Poincaré duality pairing of $H(X;\Lambda_0)$ is independent of $\mathfrak b$ and so obviously extended.

    This (extended) residue pairing is nondegenerate and defines a 'Riemannian metric' on $H(X;\Lambda_0)$. (Here we put parenthesis since our metric is over the field of fractions Λ of Novikov ring and is not over $\mathbb{R}$.) As in the standard Riemannian geometry, it determines the Levi-Civita connection

∇ on $H(X;\Lambda_0)$. Theorem 1.1 (2) then implies that this connection is flat: In fact, in the A-model side this connection is noting but the canonical affine connection on the affine space $H(X;\Lambda_0)$.

Let w_i be affine coordinates of $H(X;\Lambda_0)$ i.e. the coordinates satisfying $\nabla_{\partial/\partial w_i}\partial/\partial w_j = 0$. Then we have a function Φ on $H(X;\Lambda_0)$ that satisfies

$$\langle \mathbf{f}_i \cup^{\mathfrak b} \mathbf{f}_j, \mathbf{f}_k\rangle_{\mathrm{PD}_X} = \frac{\partial^3\Phi}{\partial w_i \partial w_j \partial w_k}. \tag{1.2}$$

Here $\{\mathbf{f}_i\}$ is the basis of $H(X;\Lambda_0)$ corresponding to the affine coordinates w_i and $\langle\cdot,\cdot\rangle_{\mathrm{PD}_X}$ is the Poincaré duality pairing.

In fact Φ is constructed from Gromov-Witten invariants and is called Gromov-Witten potential. (See [Ma], for example .) Using the isomorphism given in Theorem 1.1 (1) we find that the third derivative of Φ also gives the structure constants of the Jacobian ring.

(d) In Section 10 [FOOO5], we defined an Euler vector field $\mathfrak{E}$ on $H(X;\Lambda_0)$. We also remark that the Jacobian ring carries a unit that is parallel with respect to the Levi-Civita connection: This is obvious in the A-model side and hence the same holds in the B-model side.

(e) The discussion above implies that the miniversal family $\mathfrak{PO}_{\mathfrak b}$ determines the structure of Frobenius manifold on $H(X;\Lambda_0)$. K. Saito [Sa] and M. Saito [MSa] defined a Frobenius manifold structure on the parameter space of miniversal deformation of a holomorphic function germ with isolated singularities. Their proof is based on a very deep notion of primitive forms. On the other hand, our construction described above uses mirror symmetry which is different from theirs. Strictly speaking, our situation is also slightly different from Saito's in that we work over the Novikov ring but not over $\mathbb{C}$, and deals with a more global version than that of function germs. See [Sab] (especially its Section 4) for some discussion of a global version of Saito theory over $\mathbb{C}$. It is not clear to the authors how difficult it is to adapt Saito's argument to our situation and prove existence of a Frobenius manifold structure using the primitive forms.

(f) Dubrovin [Dub] showed that quantum cohomology also defines a Frobenius manifold structure on $H(X;\Lambda_0)$. Theorem 1.1 can be regarded as the coincidence of the two Frobenius manifolds, one defined by quantum cohomology (= toric A-model) and the other defined by Saito's theory (= Landau-Ginzburg B-model).

Remark 1.2. (1) Saito invented a notion, which he calls *flat structure*. It is the same structure as Frobenius manifold structure which Dubrovin found in Gromov-Witten theory.

(2) For the case of Fano X and with $\mathfrak b = 0$, a statement similar to Theorem 1.1 (1) was known to Batyrev [B1] and Givental [Gi2]. ([Gi2] contains its proof relying on the virtual localization formula which was later established in [GrPa].) For the case of $\mathbb{C}P^n$, a result closely related to Theorem 1.1 was known to Takahashi [Ta] ($n = 1$) and Baranikov [Bar]. Proofs of both Takahashi and Baranikov are rather computational. On the other hand, our proof does not rely on computations but uses a natural construction of the map $\mathfrak{ks}_{\mathfrak b}$ which we prove is an isomorphism. We prove this isomorphism property using some geometric argument without carrying out separate

calculations on both sides. In addition to this geometric argument, we use some important partial calculation of $\mathfrak{PO}_{\mathfrak b}$ carried out by Cho-Oh [CO] in our proof of Theorem 1.1.

(3) In the work [HV] of Hori-Vafa a function similar to the potential function $\mathfrak{PO}_0$ appears[1] and their discussion is somewhat close to Theorem 1.1. Also there are various conjectures closely related to Theorem 1.1 in the literature such as those stated by Saito-Takahashi [SaTa], Iritani [Iri2] and others. These conjectures are more wide-ranging than Theorem 1.1 in their scope in that they also discuss the Hodge structure, higher residues and others. Especially in [Iri2] a similar statement as Theorem 1.1 is announced as a theorem, whose proof is deferred to subsequent papers. On the other hand, in [Iri2] and other references, superpotential in the B-model side are defined 'by hand' and the role of potential function *with bulk* $\mathfrak{PO}_{\mathfrak b}$ was not discussed. In that sense, Theorem 1.1 does not seem to have appeared in the previous literature even as a conjecture. See also [KKP].

(4) We also remark that our potential function $\mathfrak{PO}_{\mathfrak b}$ in Lagrangian Floer theory is slightly different from the superpotential appearing in the literature we mentioned above: The 'superpotential' in the literature coincides with the leading order potential function $\mathfrak{PO}_0$ in our terminology. $\mathfrak{PO}_0$ coincides with potential function $\mathfrak{PO}_{\mathfrak b}$ for the case of X Fano and $\mathfrak b = \mathbf{0}$. There is an example where $\mathfrak{PO}_0$ is different from $\mathfrak{PO}_{\mathfrak b}$, $\mathfrak b = \mathbf{0}$ in non-Fano case. (Theorem 16.1.) However we can prove that $\mathfrak{PO}_0$ is the same as $\mathfrak{PO}_{\mathfrak b}$ for some $\mathfrak b$ up to an appropriate coordinate change (Theorem 11.1).

(5) We believe that using Theorem 1.1 we can prove a version of homological mirror symmetry conjecture ([Ko2]) between toric A-model and Landau-Ginzburg B-model, namely an equivalence of Fukaya category of X and the category of matrix factorization of Jac($\mathfrak{PO}_{\mathfrak b}$). We plan to explore this point elsewhere in the near future, with M. Abouzaid.

(6) It seems important to push our approach further and study the mirror symmetry along the lines of Saito, Givental and Hori-Vafa. For example, including higher residues, primitive forms, quantum D-module and hypersurfaces of toric manifolds into our story are some interesting projects. We hope to pursue this line of research in the future.

(7) After the main results of this paper was announced, an interesting result of Gross [Gro] appears which studies a similar problem in the case of $\mathbb{C}P^2$. See also [GPS].

We can apply Theorem 1.1 to the study of Lagrangian Floer theory of toric manifolds. Let $\mathfrak{M}(X,\mathfrak b)$ be the set of pairs (u,b) with

$$u \in \text{Int}P, \quad b \in H^1(L(u);\Lambda)/H^1(L(u);2\pi\sqrt{-1}\mathbb{Z})$$

such that

$$HF((L(u),\mathfrak b,b),(L(u),\mathfrak b,b);\Lambda) \neq 0.$$

Here $HF((L(u),\mathfrak b,b),(L(u),\mathfrak b,b);\Lambda)$ is the Floer cohomology with bulk deformation. (Λ is defined in Definition 1.6 (See [FOOO2], [FOOO5].)

---

[1] Hori and Vafa conjectured that the superpotential in their situation coincides with the potential in Lagrangian Floer theory. A similar conjecture was also made by Givental around the same time. This conjecture was one of the motivations of the work [CO].

Theorem 1.3. *If* $\mathfrak{PO}_{\mathfrak b}$ *is a Morse function, we have*

$$\#\mathfrak{M}(X,\mathfrak b) = \sum_k \operatorname{rank} H_k(X;\mathbb{Q}). \tag{1.3}$$

If $\mathfrak{PO}_{\mathfrak b}$ *is not a Morse function, we have*

$$0 < \#\mathfrak{M}(X,\mathfrak b) < \sum_k \operatorname{rank} H_k(X;\mathbb{Q}). \tag{1.4}$$

Theorem 1.3 settles a conjecture in the preprint version of [FOOO4]. It also removes the rationality hypothesis we put in the statement of Theorem 1.4 [FOOO4]. Theorem 1.3 is proved in Section 2 right after Theorem 2.32, using Theorem 1.1 (1).

We have another application of Theorem 1.1 which is proved in Section 15.

Theorem 1.4. *Assume* $\mathfrak b \in H^2(X;\Lambda_0)$. *The set of eigenvalues of the map* $x \mapsto c_1(X) \cup^{\mathfrak b} x : H(X;\Lambda_0) \to H(X;\Lambda_0)$ *coincides with the set of critical values of* $\mathfrak{PO}^{\mathfrak b}$, *with multiplicities counted.*

Here $\cup^{\mathfrak b}$ is the quantum cup product associated with the bulk deformation by $\mathfrak b$. (See Definition 2.26 for its definition.)

Remark 1.5. Theorem 1.4 was conjectured by M. Kontsevich. See also [Aur1]. Theorem 1.4 is proved in [FOOO4] in case X is Fano and $\mathfrak b = \mathbf{0}$.

Outlines of the proof of Theorem 1.1 and of the contents of the following sections are in order. The proof of Theorem 1.1 (1) goes along the line we indicated in Remark 6.15 [FOOO4]. However the details of the proof are much more involved partially because we have much amplified the statement.

In Section 2 we define various notions used in the statement of Theorem 1.1 and make a precise statement thereof. In Section 3, we make explicit computations for the case of $X = \mathbb{C}P^n$, $\mathfrak b = \mathbf{0}$ to illustrate Theorem 1.1. Section 4 contains several (technical) issues related to the completion of Laurent polynomial ring over Novikov ring.

As we already mentioned, the potential function we defined in [FOOO5] is parametrized by the group of T^n-invariant cycles. The set of T^n-invariant cycles, which we denote by $\mathcal{A}(\Lambda_0)$, is mapped surjectively on the cohomology group $H(X;\Lambda_0)$. However this map is not injective. In Section 8, we prove that if $\mathfrak b$ and $\mathfrak b'$ give the same cohomology class in $H(X;\Lambda_0)$, $\mathfrak{PO}_{\mathfrak b}$ and $\mathfrak{PO}_{\mathfrak b'}$ coincide up to an appropriate change of coordinates. (See Theorem 8.7.) In Section 7 we prove that the map $\mathfrak{ks}_{\mathfrak b}$, which is originally defined on $\mathcal{A}(\Lambda_0)$, descends to a map defined on $H(X;\Lambda_0)$. (See Theorem 7.1.)

In Section 9, we prove that the map $\mathfrak{ks}_{\mathfrak b}$ is a ring homomorphism. In Section 31 we explain its relation to various results, ideas, and conjectures appearing in the previous literature. We also explain there how the discussion of Section 9 can be generalized beyond the case of toric manifolds.

In Section 10, surjectivity of $\mathfrak{ks}_{\mathfrak b}$ is proved. We first show that its $\mathbb{C}$-reduction $\overline{\mathfrak{ks}}_{\mathfrak b}$ is surjective and then use it to prove surjectivity of $\mathfrak{ks}_{\mathfrak b}$ itself.

Sections 11 - 14 are devoted to the proof of injectivity of $\mathfrak{ks}_{\mathfrak b}$. Since this proof is lengthy and nontrivial, we give some outline of this proof here.

We first remark that $\mathfrak{PO}_{\mathfrak b}$ is not a Laurent polynomial in general but a formal Laurent power series of $y_1,\dots,y_n$ with elements of Λ_0 as its coefficients, which converges in an appropriate T-adic topology. We note that even in the Fano case we need to work with such formal power series as soon as we turn on bulk deformations. Therefore we first show that $\mathfrak{PO}_{\mathfrak b}$ can be transformed into a Laurent *polynomial* by an appropriate change of variables. This is proved in Section 12. This is possible because of the *versality* of our family which we prove in Section 11 using the surjectivity of $\overline{\mathfrak{ks}}_{\mathfrak b}$.

The second point to take care of is the fact that we actually need to take a closure of the ideal generated by $y_i\frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i}$. This is especially important when we work over a Laurent polynomial ring and not over its completion. For example, the equality (1.3) will not hold in the non Fano case, if we do not take the closure of the ideal. (See Example 8.2 [FOOO4].) This point is discussed in Section 14.

After these preparations, we proceed with the proof of injectivity of $\mathfrak{ks}_{\mathfrak b}$ as follows. Using the surjectivity of $\mathfrak{ks}_{\mathfrak b}$ and the fact that $H(X;\Lambda_0)$ is a free Λ_0 module, we observe that the injectivity of $\mathfrak{ks}_{\mathfrak b}$ will follow from the inequality

$$\mathrm{rank}_\Lambda(\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})\otimes_{\Lambda_0}\Lambda) \ge \mathrm{rank}_\Lambda H(X;\Lambda). \tag{1.5}$$

One possible way of proving the inequality (1.5) is to explicitly construct the inverse homomorphism from $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$ to $H(X;\Lambda)$. We have not been able to directly construct such a map on $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$ itself but will construct a map from a ring similar to $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$ instead. An outline of this construction is now in order:

In [MT], McDuff-Tolman proved that the Seidel's [Se1] representation

$$\pi_1(Ham(X)) \to H(X;\Lambda_0)$$

provides m elements $z'_i \in H(X;\Lambda_0)$ (Recall m is the number of T^n-invariant cycles (or toric divisors) of X of (real) codimension 2.) that satisfy the quantum Stanley-Reisner relations (see Definition 6.4 [FOOO4] and Section 13 of this paper) and which are congruent modulo Λ_+ to the Poincaré dual of the generators of the ring of T^n-invariant cycles. See Theorem 13.2. We use this to show that the quantum cohomology ring is expressed as the quotient of $\Lambda_0[z_1,\dots,z_m]$ by an ideal which is a closure of the ideal generated by quantum Stanley-Reisner relations and other relations congruent to the relations $y_i\frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} = 0$. In other words, quantum cohomology $(H(X;\Lambda),\cup^{\mathfrak b})$ is represented as the quotient of polynomial ring of m variables over Λ by the relations which are congruent to those used for the Jacobian ring $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$. Using this congruence, we find a family of schemes parameterized by one variable s which interpolates $Spec(\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})\otimes_{\Lambda_0}\Lambda)$ and $Spec(H(X;\Lambda),\cup^{\mathfrak b})$. This is a family of zero dimensional schemes. Therefore if it is *proper* and *flat* over $Spec(\Lambda[s])$, the orders (counted with multiplicity) of the fibers, which is nothing but the rank over Λ of the coordinate ring of the fiber, are independent of s and hence (1.5) follows by the standard fact in algebraic geometry. The proof of the properness is based on the fact that $\mathfrak{PO}_{\mathfrak b}$ has no critical point whose valuation lies in a sufficiently small neighborhood of the boundary of the moment polytope. The proof of Theorem 1.1 (1) is completed in Section 14.

In Section 15 we prove Theorem 1.4. We use it in Section 16 to calculate the potential function of second Hirzebruch surface F_2 at $\mathfrak b = \mathbf{0}$. (The same result was previously obtained by Auroux [Aur2] by a different method. See also [FOOO7].) We also calculate residue pairing in that case and verify Theorem 1.1 (2) by a direct calculation.

Sections 17 - 21 are devoted to the proof of coincidence of the Poincaré duality and the residue pairing. We summarize the main idea of the proof now.

Using the Poincaré duality applied to both X and $L(u)$, we dualize the map $\mathfrak{ks}_{\mathfrak b}$ and obtain the adjoint map $\mathfrak{ks}_{\mathfrak b}^*$. A description of this map in terms of the operator $\mathfrak p$ introduced in Section 3.8 [FOOO3] is given in Section 17. (We recall that another operator $\mathfrak q$ was used in [FOOO5] in the definition of the potential function with bulk.) We then calculate the pairing

$$\langle \mathfrak{ks}_{\mathfrak b}^*(PD[pt]), \mathfrak{ks}_{\mathfrak b}^*(PD[pt])\rangle_{\mathrm{PD}_X} \tag{1.6}$$

where $PD[pt]$ is the top dimensional cohomology class in $H(L(u);\Lambda_0)$ as a Floer cohomology class in

$$HF((L(u),\mathfrak b,b);(L(u),\mathfrak b,b);\Lambda_0) \cong H(L(u);\Lambda_0).$$

By inspecting the definition of $\mathfrak{ks}_{\mathfrak b}^*$, we find that the paring (1.6) can be calculated by counting the number of holomorphic maps w with boundary lying on $L(u)$ from certain bordered semi-stable curve Σ which is obtained by attaching two discs at their centers. We also require w to satisfy an appropriate constraint related to the class $[pt]$. Note that Σ is a degeneration of the annulus, i.e., of a compact bordered Riemann surface of genus 0 with 2 boundary components. So calculation of (1.6) is turned into a counting problem of appropriate holomorphic annuli. By deforming complex structures of the annuli, we prove that (1.6) is equal to an appropriate double trace of the operation $\mathfrak m_2$ on $HF((L(u),\mathfrak b,b);(L(u),\mathfrak b,b);\Lambda_0)$, which is our definition of residue pairing. (See Theorem 20.1.) Section 20 contains several constructions related to open-closed Gromov-Witten theory of one-loop. These results are new. For example, they did not appear in [FOOO3].

One important point in carrying out this proof is that we need to reconstruct a filtered A_∞ structure so that it becomes cyclically symmetric. In other words we need to take the perturbation of the moduli space of pseudo-holomorphic discs so that it is invariant under the cyclic permutation of the boundary marked points. This is because fiber products of the moduli spaces of pseudo-holomorphic discs appear at the boundary of the moduli spaces of pseudo-holomorphic annuli where we need cyclic symmetry for the consistency of the perturbation.

The perturbation we used to define the operator $\mathfrak q$ in [FOOO5] is not cyclically symmetric. In [Fu2], the cyclically symmetric perturbations of the moduli space of pseudo-holomorphic discs are produced. (In [Fu2] it is not assumed that X is a toric manifold.) For this purpose, *continuous family of* multisections is used therein. In Section 18 we adapt this scheme to the present situation where X is toric and L is a T^n-orbit, and construct a cyclically symmetric versions $\mathfrak q^{\mathfrak c}$, $\mathfrak m^{\mathfrak c}$ of our operators $\mathfrak q$, $\mathfrak m$. We also discuss their relationship with the operators $\mathfrak q$ and $\mathfrak m$ in [FOOO5].

The proof of Theorem 1.1 is completed in Section 21.

If a primitive form takes a simple form $dy_1\wedge\cdots\wedge dy_n/(y_1\cdots y_n)$, the residue pairing is the inverse of the determinant of the Hessian matrix of the potential function. (See Definition 2.17.) We will prove that this is the case if either (1) $n=2$ or (2) X is nef and $\mathfrak b$ is of degree 2, as follows. It was proved by Cho [Cho2] that the Floer cohomology ring $(HF((L(u),\mathfrak b,b),(L(u),\mathfrak b,b);\Lambda),\mathfrak m_2)$ is isomorphic to the Clifford algebra associated to the Hessian matrix of $\mathfrak{PO}_{\mathfrak b}$. This isomorphism between Clifford algebra and Lagrangian Floer cohomology holds in general if $\mathfrak{PO}$ is Morse. However in general we do not know whether this isomorphism respects Poincaré duality of $L(u)$. We prove this under the condition that (1) or (2) above holds. (See Theorem 22.2.) In Section 23 we use it to prove coincidence of the residue pairing (in our sense) and the inverse of the Hessian determinant modulo the higher order term in the general case.

In Section 24 we calculate cyclic (co)homology of Clifford algebra using Connes' periodicity exact sequence. We use this to prove that residue paring in our sense detects the second of the nontrivial elements of cyclic cohomology. Note the first nontrivial element of cyclic cohomology is detected by the critical value, which in turn becomes the first Chern class. (We also review the relation between cyclic cohomology and deformation of cyclic A_∞ algebra.)

In Sections 25-26 we discuss orientation and sign. Especially we study the case of moduli space of pseudo-holomorphic annuli, which was not discussed in [FOOO3].

In Sections 28-30 we discuss the construction of Kuranishi structure of the moduli space of pseudo-holomorphic discs which is invariant under T^n action and other symmetries. Those sections are rather technical but necessary and important substance for the constructions of this paper.

Hochschild and cyclic (co)homology of Lagrangian Floer cohomology and of Fukaya category is known to be related to the quantum cohomology of the ambient symplectic manifold. We prove and use this relationship in this paper. In Section 31 we summarize various known facts and previous works on this relationship.

A part of the results of this paper is announced in [FOOO5] Remark 10.3 and [FOOO4] Remark 6.15, as well as several lectures by the authors.

The authors would like to thank M. Abouzaid for helpful discussions especially on those related to Theorem 1.1 (2), Remark 1.2 (5) and Proposition 24.3. The second named-author thanks NIMS in Korea for its financial support and hospitality during his stay in the fall of 2009 when a large chunk of the current research was carried out. The first named-author thanks MSRI for its financial support and hospitality during his stay in the spring of 2010 when he was working on Section 29 of this paper. The third and fourth named authors thank KIAS for its financial support and hospitality during their stay in the summer of 2010, when the final version of this paper was prepared.

Notations and terminologies

Definition 1.6. Let R be a commutative ring with unit. We define the *universal Novikov ring* $\Lambda_0(R)$ as the set of all formal sums

$$\sum_{i=0}^{\infty} a_i T^{\lambda_i} \tag{1.7}$$

where $a_i \in R$ and $\lambda_i \in \mathbb{R}_{\geq 0}$ such that $\lim_{i\to\infty} \lambda_i = \infty$. Here T is a formal parameter.

We allow $\lambda_i \in \mathbb{R}$ in (1.7) (namely negative λ_i) to define $\Lambda(R)$ which we call *universal Novikov field*. It is a field of fraction of $\Lambda_0(R)$.

We require $\lambda_i > 0$ in (1.7) to define $\Lambda_+(R)$, which is the maximal ideal of $\Lambda_0(R)$.

In this paper we mainly use the case $R = \mathbb{C}$. In such a case we omit $\mathbb{C}$ and write Λ_0, Λ, Λ_+.

Definition 1.7. A *discrete monoid* G is a submonoid of additive monoid $\mathbb{R}_{\geq 0}$ which is discrete. An element (1.7) is said to be *G-gapped* if all the exponents λ_i are contained in G.

For $\beta \in H_2(X, L; \mathbb{Z})$, we denote by $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta)$ the compactified moduli space of the genus zero bordered holomorphic maps in class $\beta \in H_2(X, L(u); \mathbb{Z})$ with $k+1$ boundary marked points and ℓ interior marked points. Namely it is the set of all isomorphism classes of $(\Sigma, \vec{z}, \vec{z}^+, u)$, where Σ is a genus zero bordered Riemann surface with one boundary component, $u : (\Sigma, \partial\Sigma) \to (X, L)$ is a holomorphic map of homology class β, $\vec{z} = (z_0, \dots, z_k)$ is an ordered set of $k+1$ distinct boundary marked points, and $\vec{z}^+ = (z_1^+, \dots, z_\ell^+)$ is an ordered set of ℓ distinct interior marked points. (Sometimes, we denote the interior marked points by $\vec{z}^{\text{int}} = (z_1^{\text{int}}, \dots, z_\ell^{\text{int}})$, instead of $\vec{z}^+ = (z_1^+, \dots, z_\ell^+)$.) We require the order of $k+1$ boundary marked points respects the counter-clockwise cyclic order of the boundary $\partial\Sigma$ and also assume appropriate stability conditions.

This space is a compact Hausdorff space and has a Kuranishi structure with boundary and corners. (See [FOOO3] Section 7.1.)

Definition 1.8. A ($\mathbb{R}$ valued non-Archimedean) *valuation* of a commutative ring R with unit is by definition a map $\mathfrak{v} : R \to \mathbb{R} \cup \{\infty\}$ such that

1. $\mathfrak{v}(ab) = \mathfrak{v}(a) + \mathfrak{v}(b)$.
2. $\mathfrak{v}(a+b) \geq \min\{\mathfrak{v}(a), \mathfrak{v}(b)\}$.
3. $\mathfrak{v}(0) = \infty$.

It uniquely induces a valuation on its field of fractions $F(R)$. (See [BGR] p41 Proposition 4.) We denote the induced valuation on $F(R)$ by the same symbol $\mathfrak{v}$. A ring R with a valuation $\mathfrak{v}$ is called a *valuation ring* if

$$R = \{x \in F(R) \mid \mathfrak{v}(x) \geq 0\}.$$

In general, a ring R with a valuation $\mathfrak{v}$ may not coincide with $\{x \in F(R) \mid \mathfrak{v}(x) \geq 0\}$. In fact, we will see in Section 2 that the universal Novikov ring Λ_0 is a valuation ring, but we also treat several rings with valuations which are not valuation rings in the sense of above.

When a valuation $\mathfrak{v}$ is given, it defines a natural non-Archimedean norm on R given by

$$|a|_{\mathfrak{v}} := e^{-\mathfrak{v}(a)}$$

and hence defines a metric topology on R. By the property (1) in Definition 1.8, $|\cdot|_{\mathfrak{v}}$ is multiplicative, i.e., satisfies $|ab|_{\mathfrak{v}} = |a|_{\mathfrak{v}}|b|_{\mathfrak{v}}$. We say $(R, \mathfrak{v})$ is *complete* if it is complete with respect to the metric induced by the norm $|\cdot|_{\mathfrak{v}}$.

For a T^n orbit $L(u)$ of our toric manifold we denote by $\mathbf{e}_i$, $i = 1, \dots, n$ the basis of $H^1(L(u); \mathbb{Z})$ and we write an element $b \in H^1(L(u); \Lambda_0)$ as

$$b = \sum_{i=1}^{n} x_i \mathbf{e}_i.$$

We put $y_i = e^{x_i}$.

2. Statement of the results

In this section we describe the main results of this paper. We also review various results of [FOOO5] and of several other previous works and introduce various notations that we use in later sections.

We first discuss potential function and its Jacobian ring. Let (X, ω) be a compact toric manifold and $\pi : X \to P$ be the moment map. Let $n = \dim_{\mathbb{C}} X$ and $m =$ the number of irreducible components of $D = \pi^{-1}(\partial P)$ = the number of codimension one faces of P. Let B be the number of faces of P of arbitrary codimension. Let $\{P_a\}_{a=1,\dots,B}$ be the set of all faces of P such that $P_i = \partial_i P$ $(i = 1,\dots,m)$ are the faces of codimension one. Denote $D_a = \pi^{-1}(P_a)$ which are T^n-invariant cycles. We denote by $\mathcal{A}^k$ for $k \neq 0$ the free $\mathbb{Z}$ module generated by D_a's with $\operatorname{codim} P_a = k$. Then $P_0 = P$ and $D_0 = X$. We put $\mathcal{A}^0 = \mathbb{Z}[D_0]$. (We remark here this notation is slightly different from one in [FOOO5]. Namely for $k \neq 0$, the group $\mathcal{A}^k$ is the same as the one used in [FOOO5]. But in [FOOO5] we put $\mathcal{A}^0 = \{0\}$ instead.) We put $\mathcal{A} = \mathcal{A}(\mathbb{Z}) = \oplus_k \mathcal{A}^k$ and $\mathcal{A}(\Lambda_0) = \mathcal{A} \otimes \Lambda_0$.

We put $\mathbf{f}_a = [D_a]$, $a = 0, 1, \dots, B$. An element $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ is written as

$$\mathfrak{b} = \sum_{a=0}^{B} w_a(\mathfrak{b})\mathbf{f}_a$$

so w_a's are coordinates of $\mathfrak{b}$. Let $y_1,\dots,y_n$ be formal variables. For $u = (u_1,\dots,u_n) \in \operatorname{Int} P$ we define n variables $y_j(u)$ $(j = 1,\dots,n)$ by

$$y_j(u) = T^{-u_j} y_j. \tag{2.1}$$

We consider the Laurent polynomial ring:

$$\Lambda[y_1, y_1^{-1}, \dots, y_n, y_n^{-1}],$$

which is canonically isomorphic to

$$\Lambda[y_1(u), y_1(u)^{-1}, \dots, y_n(u), y_n(u)^{-1}],$$

by (2.1). We write this ring $\Lambda[y(u), y(u)^{-1}]$ or $\Lambda[y, y^{-1}]$. If $u' = (u'_1,\dots,u'_n) \in \operatorname{Int} P$ is another point we have a canonical isomorphism

$$\psi_{u',u} : \Lambda[y(u), y(u)^{-1}] \to \Lambda[y(u'), y(u')^{-1}]$$

that is

$$\psi_{u',u}(y_j(u)) = T^{u'_j - u_j} y_j(u').$$

For $i = 1,\dots,m$, we denote

$$\mathfrak{w}_i = \exp w_i = \sum_{k=0}^{\infty} \frac{1}{k!} w_i^k.$$

We consider the Laurent polynomial ring

$$\Lambda[w_0, \mathfrak{w}_1, \mathfrak{w}_1^{-1}, \dots, \mathfrak{w}_m, \mathfrak{w}_m^{-1}, w_{m+1}, \dots, w_B, y, y^{-1}],$$

and denote it by

$$\Lambda[\mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}].$$

A non-Archimedean valuation $\mathfrak{v}_T$ on Λ is given by

$$\mathfrak{v}_T\left(\sum_{i=1}^{\infty} a_i T^{\lambda_i}\right) = \inf\{\lambda_i \mid a_i \neq 0\}, \quad \mathfrak{v}_T(0) = +\infty. \tag{2.2}$$

We define a valuation $\mathfrak{v}_T^u$ on $\Lambda[\mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}]$ by assigning

$$\mathfrak{v}_T^u(\mathfrak{w}_i) = \mathfrak{v}_T^u(w_i) = \mathfrak{v}_T^u(y_i(u)) = 0, \quad \mathfrak{v}_T^u(T) = 1. \tag{2.3}$$

It implies

$$\mathfrak{v}_T^u(y_j(u')) = u_j - u'_j, \quad \mathfrak{v}_T^u(y_j) = u_j. \tag{2.4}$$

We also define

$$\mathfrak{v}_T^P = \inf_{u \in P} \mathfrak{v}_T^u.$$

Let $\Lambda^P\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$ be the completion of $\Lambda[\mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}]$ with respect to the norm $\mathfrak{v}_T^P$. We remark that $\mathfrak{v}_T^P$ is not a valuation but $e^{-\mathfrak{v}_T^P}$ still defines a norm. We also put

$$\Lambda_0^P\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle = \{x \in \Lambda^P\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle \mid \mathfrak{v}_T^P(x) \geq 0\}.$$

We define $\Lambda^P\langle\langle y, y^{-1}\rangle\rangle$, $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ in a similar way.

Let $\Lambda_0^u\langle\langle y, y^{-1}\rangle\rangle$ (resp. $\Lambda_0^u\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$) be the completion of the rings $\Lambda_0[y(u), y(u)^{-1}]$ (resp. $\Lambda_0[\mathfrak{w}, \mathfrak{w}^{-1}, w, y(u), y(u)^{-1}]$) with respect to $\mathfrak{v}_T^u$. We denote by $\Lambda_+^u\langle\langle y, y^{-1}\rangle\rangle$ the ideal of $\Lambda_0^u\langle\langle y, y^{-1}\rangle\rangle$ consisting of all the elements with $\mathfrak{v}_T^u > 0$.

To handle the case of $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ we need to use slightly different completion.

Definition 2.1. We put

$$\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle = \bigcap_{u \in \operatorname{Int} P} \Lambda_0^u\langle\langle y, y^{-1}\rangle\rangle,$$

$$\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle = \bigcap_{u \in \operatorname{Int} P} \Lambda_0^u\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle.$$

Here the definition of the intersection

$$\Lambda_0^u\langle\langle y, y^{-1}\rangle\rangle \cap \Lambda_0^{u'}\langle\langle y, y^{-1}\rangle\rangle \tag{2.5}$$

is as follows: An element $\mathfrak{a}$ of $\Lambda_0^u\langle\langle y, y^{-1}\rangle\rangle$ is a limit of a sequence of elements $\mathfrak{a}_i \in \Lambda_0[y, y^{-1}]$ in $\mathfrak{v}_T^u$ topology. We say it is in the intersection (2.5) if we may choose $\mathfrak{a}_i$ so that it converges in $\mathfrak{v}_T^{u'}$ topology also and $\lim_{i\to\infty} \mathfrak{v}_T^{u'}(\mathfrak{a}_i) \geq 0$.

In Section 4, we will define a metric on these rings in Definition 4.2 and show that indeed they are the completions of the corresponding Laurent polynomial rings with respect to the metric.

We extend the $\mathfrak{v}_T$ norm on Λ_0 to $\Lambda_0[Z_1, \dots, Z_m]$ by setting $\mathfrak{v}_T(Z_i) = 0$ $i = 1, \dots, m$. Let $\Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle$ be the completion of the polynomial ring $\Lambda_0[Z_1, \dots, Z_m]$ with respect to this $\mathfrak{v}_T$ norm thereon.

Here and hereafter we use the convention of [FOOO4] Section 2 on toric manifolds with P being its moment polytope. In [FOOO4] we use the affine functions $\ell_j : \mathbb{R}^n \to \mathbb{R}$, $(j = 1, \cdots, m)$ defined by $\ell_j(u) = \langle u, v_j\rangle - \lambda_j$ such that

$$P = \{u \mid \ell_j(u) \geq 0\} = \{u \mid \langle u, v_j\rangle \geq \lambda_j\}. \tag{2.6}$$

Let $\vec{v}_j = (v_{j,i})_{i=1}^n$ be the gradient vector. We have $v_{j,i} \in \mathbb{Z}^n$.

Definition 2.2. We define $z_j \in \Lambda_0[y(u), y(u)^{-1}]$ by

$$z_j = T^{\ell_j(u)} \prod_{i=1}^n y_i(u)^{v_{j,i}} = T^{-\lambda_j} \prod_{i=1}^n y_i^{v_{j,i}}. \tag{2.7}$$

Here $\lambda_j = \langle u, v_j\rangle - \ell_j(u)$ for $j = 1, \dots, m$. If $u \in P$, then $\ell_j(u) \geq 0$ and hence z_j defines an element of $\Lambda_0[y(u), y(u)^{-1}]$ from the first expression. Note that we use (2.1) to get the second expression, from which we can also regard $z_j \in \Lambda[y, y^{-1}]$ with $\mathfrak{v}_T^u(z_j) = -\lambda_j + \langle u, v_j\rangle = \ell_j(u) \geq 0$ by (2.3) and (2.4), if $u \in P$.

Lemma 2.3. *We have surjective continuous ring homomorphisms*

$$\Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle \to \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle, \quad \Lambda_0[[Z_1, \dots, Z_m]] \to \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle \tag{2.8}$$

which send Z_j to z_j. We also have surjective continuous ring homomorphisms

$$\begin{aligned}\Lambda_0\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, Z_1, \dots, Z_m\rangle\rangle &\to \Lambda_0^P\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle \\ \Lambda_0\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w\rangle\rangle[[Z_1, \dots, Z_m]] &\to \Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle,\end{aligned} \tag{2.9}$$

which send Z_j to z_j and preserve $\mathfrak{w}$, w.

The proof will be given in Section 4.

Definition 2.4. Let G be a discrete submonoid of $\mathbb{R}_{\ge 0}$.

1. An element of $\Lambda_0[[Z_1, \dots, Z_m]]$ is said to be *G-gapped* if the exponents of T appearing in all of its coefficients are contained in G.
2. An element of $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ or of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ is said to be *G-gapped* if it is an image of a G-gapped element by the surjective homomorphism (2.8)
3. $\Lambda_+^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ denotes the image of $\Lambda_+[[Z_1, \dots, Z_m]]$ under the map (2.8) in $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.

Remark 2.5. Any element of $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ is G-gapped for some discrete submonoid G of $\mathbb{Z}_{\ge 0}$. However this is not true for the elements of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ in general.

Let $\mathbf{e}_i$, $i = 1, \dots, n$ be the baisis of $H^1(L(u); \mathbb{Z})$ and we write an element $b \in H^1(L(u); \Lambda_0)$ as $b = \sum_{i=1}^n x_i \mathbf{e}_i$. We put $y_i = e^{x_i}$.

In [FOOO5], we defined the potential function with bulk $\mathfrak{PO}(w_1, \dots, w_B; y_1, \dots, y_n)$, that is, a 'generating function of open-closed Gromov-Witten invariant' given in Section 3.8 [FOOO3]. We enhanced this function to

$$\mathfrak{PO}(w_0, w_1, \dots, w_B; y_1, \dots, y_n) = w_0 + \mathfrak{PO}(w_1, \dots, w_B; y_1, \dots, y_n)$$

by adding additional variable w_0 corresponding to the top dimensional cycle $P^0 = P$ (and $D^0 = X$). Hereafter in this paper, $\mathfrak{PO}$ will always denote this enhanced function given in the left hand side. We also denote

$$\mathfrak{PO}_{\mathfrak{b}}(y_1, \dots, y_n) = \mathfrak{PO}(w_0, w_1, \dots, w_B; y_1, \dots, y_n)$$

where $\mathfrak{b} = \sum_{i=0}^B w_i \mathbf{f}_i$. The following lemma and the remark thereafter justify our consideration of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.

We recall

$$\begin{aligned}\mathfrak{PO}_{\mathfrak{b}}(y_1, \dots, y_n) = &\sum_{\ell,\beta} \sum_{j_1,\dots,j_\ell} \frac{w_{j_1} \dots w_{j_\ell}}{\ell!} T^{\beta\cap\omega/2\pi} y_1(u)^{\partial\beta\cap\mathbf{e}_1} \cdots y_n(u)^{\partial\beta\cap\mathbf{e}_n} \\ &\qquad \mathrm{ev}_0[\mathcal{M}_{1;\ell}(\beta; D_{j_1}, \dots, D_{j_\ell})] \cap [L] \\ = &\sum_{\ell,\beta} \frac{1}{\ell!} T^{\beta\cap\omega/2\pi} y_1(u)^{\partial\beta\cap\mathbf{e}_1} \cdots y_n(u)^{\partial\beta\cap\mathbf{e}_n} \\ &\qquad \mathrm{ev}_0[\mathcal{M}_{1;\ell}(\beta; \mathfrak{b}, \dots, \mathfrak{b})] \cap [L].\end{aligned} \tag{2.10}$$

See [FOOO5] (7.5). (The moduli space $\mathcal{M}_{1;\ell}(\beta; D_{j_1}, \dots, D_{j_\ell})$ is defined at the beginning of Section 6.) Note y_i and $y_i(u)$ are related by (2.1).

Lemma 2.6. (1) *If $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, then $\mathfrak{PO}_{\mathfrak{b}} \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$.*

(2) *If $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$, then we have $\mathfrak{PO}_{\mathfrak{b}} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.*

Proof. (1) is proved in Theorem 3.11 [FOOO5]. Since $\mathfrak{PO}_{\mathfrak{b}}$ converges in $\mathfrak{v}_T^u$ norm for any $u \in \operatorname{Int} P$, (2) follows. $\square$

Remark 2.7. We do not know whether $\mathfrak{PO}_{\mathfrak{b}} \notin \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ or not, in case of general $\mathfrak{b} \in \mathcal{A}(\Lambda_0) \setminus \mathcal{A}(\Lambda_+)$.

Definition 2.8. We define the Jacobian ring $\operatorname{Jac}(\mathfrak{PO})$ by

$$\operatorname{Jac}(\mathfrak{PO}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle}{\operatorname{Clos}_{d_{\overset{\circ}{P}}}\left(\left\{y_i \frac{\partial \mathfrak{PO}}{\partial y_i} \mid i = 1, \dots, n\right\}\right)}.$$

Here $\operatorname{Clos}_{d_{\overset{\circ}{P}}}$ stands for the closure with respect to the topology induced by the metric (4.1).

Definition 2.9. If $\mathfrak{b} = \sum w_i(\mathfrak{b})\mathbf{f}_i \in \mathcal{A}(\Lambda_0)$, we define

$$\operatorname{Jac}(\mathfrak{PO}_{\mathfrak{b}}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle}{\operatorname{Clos}_{d_{\overset{\circ}{P}}}(\{y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} \mid i = 1, \dots, n\})}.$$

In other words,

$$\operatorname{Jac}(\mathfrak{PO}_{\mathfrak{b}}) = \operatorname{Jac}(\mathfrak{PO}) \otimes_{\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle} \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle.$$

Here we regard $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ as a $\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$-module via the ring homomorphism which sends w_i to $w_i(\mathfrak{b})$.

Remark 2.10. We can prove that the ideal generated by $\{y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} \mid i = 1, \dots, n\}$ is closed if $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$. (Lemma 12.8 (2).) So, in that case, $\operatorname{Clos}_{d_{\overset{\circ}{P}}}$ in the above definition is superfluous.

Remark 2.11. If $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, we have

$$\operatorname{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \cong \frac{\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle}{\operatorname{Clos}_{\mathfrak{v}_T^P}(\{y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} \mid i = 1, \dots, n\})}, \tag{2.11}$$

where the closure $\operatorname{Clos}_{\mathfrak{v}_T^P}$ is taken with respect to the norm $e^{-\mathfrak{v}_T^P}$. (See Proposition 5.6.)

In case $\mathfrak{b} \in \mathcal{A}(\Lambda_0) \setminus \mathcal{A}(\Lambda_+)$, the equality (2.11) may not hold in general. See Example 14.39.

We now recall that

$$\pi : \mathcal{A}(\Lambda_0) \to H(X; \Lambda_0); \quad \mathfrak{b} \mapsto [\mathfrak{b}] \tag{2.12}$$

is surjective map but not injective. We can choose a subset $\{\bar{\mathbf{f}}_i \mid i = 0, \dots, B'\}$ of our basis $\{\mathbf{f}_i\}$ so that $\pi(\bar{\mathbf{f}}_i)$ forms a basis of $H(X; \mathbb{Z})$ and $\bar{\mathbf{f}}_0 = \mathbf{f}_0$, and $\deg \bar{\mathbf{f}}_i = 2$ if and only if $i \le m'$. With respect to such a basis, we identify $H(X; \Lambda_0)$ with the subspace of $\mathcal{A}(\Lambda_0)$ generated by $\{\bar{\mathbf{f}}_i\}$. Let $\overline{w}_i$ be the corresponding coordinates and $\overline{\mathfrak{w}}_i$ their exponentials. We define

$$\Lambda_0^P\langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w}, y, y^{-1}\rangle\rangle, \quad \Lambda_0^{\overset{\circ}{P}}\langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w}, y, y^{-1}\rangle\rangle$$

in the same way as we did for $\Lambda_0^P\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$, $\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$. By restriction, we may regard $\mathfrak{PO}$ as

$$\mathfrak{PO} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}, \overline{w}, y, y^{-1}\rangle\rangle.$$

We then define another quotient ring

$$\mathrm{Jac}'(\mathfrak{PO}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}, \overline{w}, y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}(\{y_i\frac{\partial \mathfrak{PO}}{\partial y_i} \mid i = 1, \dots, n\})}. \tag{2.13}$$

(The difference between Jac' and Jac is the number of variables we use.)

We next define a residue pairing on $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$. For further discussion, we need to restrict ourselves to the case where $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function. We first make this notion precise.

Define the vector of valuations

$$\vec{\mathfrak{v}}_T : (\Lambda \setminus \{0\})^n \to \mathbb{R}^n$$

by

$$\vec{\mathfrak{v}}_T(\mathfrak{y}_1, \dots, \mathfrak{y}_n) = (\mathfrak{v}_T(\mathfrak{y}_1), \dots, \mathfrak{v}_T(\mathfrak{y}_n)).$$

Definition 2.12. $\mathfrak{y} = (\mathfrak{y}_1, \dots, \mathfrak{y}_n)$ is said to be a *critical point* of $\mathfrak{PO}_{\mathfrak{b}}$ if the following holds:

$$y_i\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}(\mathfrak{y}) = 0 \quad i = 1, \dots, n \tag{2.14}$$

$$\vec{\mathfrak{v}}_T(\mathfrak{y}) \in \mathrm{Int}\, P. \tag{2.15}$$

We denote $\mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$ the set of critical points of $\mathfrak{PO}_{\mathfrak{b}}$.

The following lemma will be proved in Section 4.

Lemma 2.13. *(A) The following conditions for $\mathfrak{y} = (\mathfrak{y}_1, \dots, \mathfrak{y}_n) \in (\Lambda \setminus \{0\})^n$ are equivalent:*

1. *(2.15) holds.*
2. *The ring homomorphism $\Lambda[y, y^{-1}] \to \Lambda$, $\quad y_i \mapsto \mathfrak{y}_i$ extends to a continuous homomorphism: $\Lambda^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle \to \Lambda$, $\quad y_i \mapsto \mathfrak{y}_i$.*

(B) Similarly, the following conditions for $\mathfrak{y} = (\mathfrak{y}_1, \dots, \mathfrak{y}_n) \in (\Lambda \setminus \{0\})^n$ are equivalent:

1. *$\vec{\mathfrak{v}}_T(\mathfrak{y}) \in P$.*
2. *The ring homomorphism $\Lambda[y, y^{-1}] \to \Lambda$, $\quad y_i \mapsto \mathfrak{y}_i$ extends to a continuous homomorphism: $\Lambda^{P}\langle\langle y, y^{-1}\rangle\rangle \to \Lambda$, $\quad y_i \mapsto \mathfrak{y}_i$.*

Definition 2.14.

1. A critical point $\mathfrak{y}$ is said to be *nondegenerate* if

$$\det\left[\frac{\partial^2 \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i \partial y_j}\right]_{i,j=1}^{i,j=n}(\mathfrak{y}) \neq 0.$$

2. $\mathfrak{PO}_{\mathfrak{b}}$ is said to be a *Morse function* if all of its critical points are nondegenerate.

Proposition 2.15. (1) *There is a splitting of Jacobian ring*

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \cong \prod_{\mathfrak{y} \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})} \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}; \mathfrak{y})$$

as a direct product of rings. Each of $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$ *is a local ring.*

(2) $\mathfrak{y}$ *is nondegenerate if and only if* $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y}) \cong \Lambda$.

(3) *In particular, if* $\mathfrak{PO}_{\mathfrak{b}}$ *is a Morse function, then there exists an isomorphism*

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \cong \prod_{\mathfrak{y} \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})} \Lambda. \tag{2.16}$$

When X is Fano and $\mathfrak{b} = \mathbf{0}$, this reduces to Proposition 7.10 [FOOO4]. We will prove Proposition 2.15 in Section 5.

Definition 2.16. We denote by $1_{\mathfrak{y}} \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$ the unit of the ring $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$.

We are now ready to define the simplified residue pairing:

Definition 2.17. Assume that $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function. We then define a *simplified residue pairing*

$$\langle \cdot, \cdot \rangle^0_{\mathrm{res}} : (\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda) \otimes (\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda) \to \Lambda$$

by

$$\langle 1_{\mathfrak{y}}, 1_{\mathfrak{y}'} \rangle^0_{\mathrm{res}} = \begin{cases} 0 & \text{if } \mathfrak{y} \neq \mathfrak{y}', \\ \left(\det \left[\mathfrak{y}_i^u \mathfrak{y}_j^u \frac{\partial^2 \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i \partial y_j} \right]_{i,j=1}^{i,j=n} (\mathfrak{b}; \mathfrak{y}^u) \right)^{-1} & \text{if } \mathfrak{y} = \mathfrak{y}'. \end{cases} \tag{2.17}$$

Here $u = \vec{\mathfrak{v}}_T(\mathfrak{y})$, $T^{u_i}\mathfrak{y}_i^u = \mathfrak{y}_i$ and $\mathfrak{PO}_{\mathfrak{b}}$ is as in (2.10).

Remark 2.18.

1. We remark $e^{x_i} = y_i$. Hence $y_i \frac{\partial}{\partial y_i} = \frac{\partial}{\partial x_i}$. Thus the matrix appearing in (2.17) is the Hessian matrix of $\mathfrak{PO}_{\mathfrak{b}}$ with respect to the variable x_i.
2. In that regard, the value of $\langle 1_{\mathfrak{y}}, 1_{\mathfrak{y}} \rangle^0_{\mathrm{res}}$ is the contribution to the principal part at the critical point $x_i = \log \mathfrak{y}_i$ of the oscillatory integral
$$\int e^{\sqrt{-1}\mathfrak{PO}_{\mathfrak{b}}(x)/\hbar}\, dx$$
in the stationary phase approximation as $\hbar \to 0$.

In the case when one of the following conditions is satisfied, $\langle \cdot, \cdot \rangle^0_{\mathrm{res}}$ defined above gives the residue pairing in Theorem 1.1.

Condition 2.19. X is a compact toric manifold and $\mathfrak{b} \in H(X;\Lambda_0)$.

1. $\dim_{\mathbb{C}} X = 2$.
2. X is nef and $\deg \mathfrak{b} = 2$.

We remark that a compact toric manifold X is nef if and only if for all holomorphic map $u : S^2 \to X$ we have $\int u^* c_1(X) \geq 0$.

To define our residue pairing in the general case, we need notations.

Definition 2.20. Let C be a $\mathbb{Z}_2$ graded finitely generated free Λ module. A structure of *unital Frobenius algebra* of dimension n is $\langle \cdot, \cdot \rangle : C^k \otimes C^{n-k} \to \Lambda$, $\cup : C^k \otimes C^\ell \to C^{k+\ell}$, $1 \in C^0$, such that:

1. $\langle \cdot, \cdot \rangle$ is a graded symmetric bilinear form which induces an isomorphism $x \mapsto (y \mapsto \langle x, y \rangle)$, $C^k \to Hom_{\Lambda_0}(C^{n-k}, \Lambda)$.
2. $\cup$ is an associative product on C. 1 is its unit.
3. $\langle x \cup y, z \rangle = \langle x, y \cup z \rangle$.

We remark that we do *not* assume that $\cup$ is (graded) commutative. The cohomology group of an oriented closed manifold becomes a unital Frobenius algebra in an obvious way.

Definition 2.21. Let $(C, \langle \cdot, \cdot \rangle, \cup, 1)$ be a unital Frobenius algebra. We take a basis $\mathbf{e}_I$, $I \in \mathfrak{I}$ of C such that $\mathbf{e}_0$ is the unit. Let $g_{IJ} = \langle \mathbf{e}_I, \mathbf{e}_J \rangle$ and let g^{IJ} be its inverse matrix. We define an invariant of C by

$$\begin{aligned} Z(C) = \sum_{I_1,I_2,I_3 \in \mathfrak{I}} \sum_{J_1,J_2,J_3 \in \mathfrak{I}} & (-1)^* g^{I_1J_1} g^{I_2J_2} g^{I_30} g^{J_30} \\ & \langle \mathbf{e}_{I_1} \cup \mathbf{e}_{I_2}, \mathbf{e}_{I_3} \rangle \langle \mathbf{e}_{J_1} \cup \mathbf{e}_{J_2}, \mathbf{e}_{J_3} \rangle \end{aligned} \tag{2.18}$$

where $* = \deg \mathbf{e}_{I_1} \deg \mathbf{e}_{J_2} + \frac{n(n-1)}{2}$. We call $Z(C)$ the *trace* of unital Frobenius algebra C.

It is straightforward to check that $Z(C)$ is independent of the choices of the basis. This invariant is an example of 1-loop partition functions and can be described by the following Feynman diagram.



Figure 2.1

We will discuss the invariant Z more in Section 24.

Remark 2.22. In [Fu2], a cyclic unital filtered A_∞ algebra is assigned to an arbitrary relatively spin Lagrangian submanifold L in a symplectic manifold X. So if we fix an element b of the Maurer-Cartan scheme $\mathcal{M}_{\rm weak}(L; \Lambda_0)$, then we obtain a unital Frobenius algebra. Therefore Z in Definition 2.21 induces a map

$$Z : \mathcal{M}_{\rm weak}(L; \Lambda_0) \to \Lambda. \tag{2.19}$$

This map is an invariant of L in that generality.

For the pair (X, L) with $c_1(X) = 0$ and $\mu_L = 0$, this invariant vanishes by the degree reason. There is another invariant constructed in [Fu3] for the case of dimension 3, denoted by $\Psi : \mathcal{M}(L; \Lambda_0) \to \Lambda_0$, which is different from Z given here. For example, the unit did not enter in the definition of Ψ in [Fu3].

Let $\mathfrak{b} \in H(X; \Lambda_0)$ and $(u, b) \in \mathfrak{M}(X; \mathfrak{b})$. In [FOOO5] we constructed a (unital) filtered A_∞ algebra $(H(L(u); \Lambda_0), \{\mathfrak{m}_k^{\mathfrak{b},b}\}, {\rm PD}([L(u)]))$ such that $\mathfrak{m}_1^{\mathfrak{b},b} = 0$. In Section 18 we modify it slightly and obtain another $(H(L(u); \Lambda_0), \{\mathfrak{m}_k^{\mathfrak{c},\mathfrak{b},b}\}, {\rm PD}([L(u)]))$ which has the following cyclic property in addition:

$$\begin{aligned} &\langle \mathfrak{m}_k^{\mathfrak{c},\mathfrak{b},b}(x_1, \dots, x_k), x_0 \rangle_{\rm cyc} \\ &= (-1)^{\deg' x_0 (\deg' x_1 + \dots + \deg' x_k)} \langle \mathfrak{m}_k^{\mathfrak{c},\mathfrak{b},b}(x_0, x_1 \dots, x_{k-1}), x_k \rangle_{\rm cyc}. \end{aligned} \tag{2.20}$$

Here we put

$$\begin{aligned} x \cup^{\mathfrak{c},\mathfrak{b},b} y &= (-1)^{\deg x(\deg y+1)} \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b}(x,y), && (2.21)\\ \langle x,y\rangle_{\rm cyc} &= (-1)^{\deg x(\deg y+1)} \langle x,y\rangle_{{\rm PD}_{L(u)}}. && (2.22)\end{aligned}$$

See Remark 26.5 for the precise definition of $\langle\cdot,\cdot\rangle_{{\rm PD}_{L(u)}}$. Then we can find that $\langle\cdot,\cdot\rangle_{\rm cyc}$ satisfies the property (2.20), while the Poincaré pairing $\langle\cdot,\cdot\rangle_{{\rm PD}_{L(u)}}$ itself satisfies the property (3) in Definition 2.20. Then $(H(L(u);\Lambda), \langle\cdot,\cdot\rangle_{{\rm PD}_{L(u)}}, \cup^{\mathfrak{c},b,b}, PD[L(u)])$ becomes a unital Frobenius algebra with cyclic symmetric pairing $\langle\cdot,\cdot\rangle_{\rm cyc}$. We put

$$Z(\mathfrak{b},b) = Z((H(L(u);\Lambda), \langle\cdot,\cdot\rangle_{{\rm PD}_{L(u)}}, \cup^{\mathfrak{c},\mathfrak{b},b}, {\rm PD}([L(u)])). \tag{2.23}$$

Definition 2.23. Assume that $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function. We then define a *residue pairing*

$$\langle\cdot,\cdot\rangle_{\rm res} : ({\rm Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda) \otimes ({\rm Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda) \to \Lambda$$

by

$$\langle 1_{\mathfrak{y}}, 1_{\mathfrak{y}'}\rangle_{\rm res} = \begin{cases} 0 & \text{if } \mathfrak{y} \ne \mathfrak{y}', \\ (Z(\mathfrak{b},b))^{-1} & \text{if } \mathfrak{y} = \mathfrak{y}'. \end{cases} \tag{2.24}$$

In Section 18, we also define a unital and filtered A_∞ isomorphism $\mathfrak{F}$ between $(H(L(u);\Lambda_0), \{\mathfrak{m}_k^{\mathfrak{b}}\}, {\rm PD}([L(u)]))$ and $(H(L(u);\Lambda_0), \{\mathfrak{m}_k^{\mathfrak{c},\mathfrak{b}}\}, {\rm PD}([L(u)]))$. For $b = b_0 + b_+ \in H^1(L(u);\Lambda_0)$, $b_0 \in H^1(L(u);\mathbb{R})$, $b_+ \in H^1(L(u);\Lambda_+)$ we define

$$b^{\mathfrak{c}} = b_0 + b_+^{\mathfrak{c}}$$

by

$$b_+^{\mathfrak{c}} = \mathfrak{F}_*(b_+).$$

We have $b_+^{\mathfrak{c}} \in H^{odd}(L(u);\Lambda_+)$ in general but $b_+^{\mathfrak{c}} \in H^1(L(u);\Lambda_+)$ in case (1) or (2) of Condition 2.19 holds. We put:

$$\mathfrak{PO}_{\mathfrak{b}}^{\mathfrak{c}} = \mathfrak{PO}_{\mathfrak{b}} \circ \mathfrak{F}_*^{-1}.$$

This is related to Definition 2.17 as follows.

Theorem 2.24. (1) *Assume that* $b = \sum x_i \mathbf{e}_i$, $x_i \equiv \overline{x}_i \mod \Lambda_+$ *is a nondegenerate critical point of* $\mathfrak{PO}_{\mathfrak{b}}^u$. *Then*

$$\begin{aligned} Z(\mathfrak{b},b) &\equiv \det\left[y_i y_j \frac{\partial^2 \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i \partial y_j}\right]_{i,j=1}^{i,j=n} (\mathfrak{y}) \\ &\equiv \det\left[y_i y_j \frac{\partial^2 \mathfrak{PO}_{0,\mathfrak{b}}}{\partial y_i \partial y_j}\right]_{i,j=1}^{i,j=n} (\mathfrak{y}) \mod T^\lambda \Lambda_+. \end{aligned} \tag{2.25}$$

Here $\lambda = v_T(Z(\mathfrak{b},b))$ *and* $\mathfrak{y} = (e^{x_1},\dots,e^{x_n})$.

(2) *If Condition 2.19* (1) *holds, then we have*

$$Z(\mathfrak{b},b) = \det\left[y_i y_j \frac{\partial^2 \mathfrak{PO}_{\mathfrak{b}}^{\mathfrak{c}}}{\partial y_i \partial y_j}\right]_{i,j=1}^{i,j=n} (\mathfrak{y}). \tag{2.26}$$

(3) *If Condition 2.19* (2) *holds, then we have*

$$Z(\mathfrak{b},b) = \det\left[y_i y_j \frac{\partial^2 \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i \partial y_j}\right]_{i,j=1}^{i,j=n} (\mathfrak{y}). \tag{2.27}$$

(2.27) implies that

$$\langle\cdot,\cdot\rangle_{\rm res} = \langle\cdot,\cdot\rangle^0_{\rm res} \tag{2.28}$$

in case (2) of Condition 2.19 holds.

We prove Theorem 2.24 in Sections 22 and 23.

Remark 2.25. The authors do not know a counter example to the equality (2.27) in the non-nef case. But it is unlikely that it holds in the non-nef case.

We next turn to the Frobenius manifold structure in the big quantum cohomology. Since this story is well established (See [Dub], [Ma].), we just briefly recall it mainly to fix our notation and for readers' convenience.

To simplify the sign issue, we only consider the case where the cohomology of X consist only of even degrees. (This is certainly the case for toric manifolds, which is of our main interest in this paper.)

Let (X,ω) be a symplectic manifold. For $\alpha \in H_2(X;\mathbb{Z})$ let $\mathcal{M}_\ell(\alpha)$ be the moduli space of stable maps from genus zero semi-stable curves with ℓ marked points and of homology class α. There exists an evaluation map

$$\mathrm{ev} : \mathcal{M}_\ell(\alpha) \to X^\ell.$$

The space $\mathcal{M}_\ell(\alpha)$ has a virtual fundamental cycle and hence defines a class

$$\mathrm{ev}_*[\mathcal{M}_\ell(\alpha)] \in H_*(X^\ell;\mathbb{Q}).$$

(See [FO].) Here $* = 2n + 2c_1(X)\cap\alpha + 2\ell - 6$. Let $Q_1,\dots,Q_\ell$ be cycles such that

$$\sum \operatorname{codim} Q_i = 2n + 2c_1(X)\cap\alpha + 2\ell - 6. \tag{2.29}$$

We define Gromov-Witten invariant by

$$GW_\ell(\alpha : Q_1,\dots,Q_\ell) = \mathrm{ev}_*[\mathcal{M}_\ell(\alpha)] \cap (Q_1\times\cdots\times Q_\ell) \in \mathbb{Q}.$$

We put $GW_\ell(\alpha : Q_1,\dots,Q_\ell) = 0$ when (2.29) is not satisfied.

We now define

$$GW_\ell(Q_1,\dots,Q_\ell) = \sum_\alpha T^{\alpha\cap\omega} GW(\alpha : Q_1,\dots,Q_\ell). \tag{2.30}$$

(2.30) extends to a Λ_0 module homomorphism

$$GW_\ell : H(X;\Lambda_0)^{\otimes\ell} \to \Lambda_0.$$

Definition 2.26. Let $\mathfrak{b} \in H(X;\Lambda_0)$ be given. For each given pair $\mathfrak{c},\mathfrak{d} \in H(X;\Lambda_0)$, we define a product $\mathfrak{c} \cup^{\mathfrak{b}} \mathfrak{d} \in H(X;\Lambda_0)$ by the following formula

$$\langle \mathfrak{c} \cup^{\mathfrak{b}} \mathfrak{d}, \mathfrak{e}\rangle_{\mathrm{PD}_X} = \sum_{\ell=0}^\infty \frac{1}{\ell!} GW_{\ell+3}(\mathfrak{c},\mathfrak{d},\mathfrak{e},\mathfrak{b},\dots,\mathfrak{b}). \tag{2.31}$$

Here $\langle\cdot,\cdot\rangle_{\mathrm{PD}_X}$ denotes the Poincaré duality. The right hand side converges and defines a graded commutative and associative ring structure on $H(X;\Lambda_0)$. We call $\cup^{\mathfrak{b}}$ the *deformed quantum cup product*.

We now discuss convergence of the above deformed quantum cup product in more detail in an adic topology. Let $\overline{\mathbf{f}}_i$ $(i = 0,\dots,B')$ be a basis of $H(X;\Lambda_0)$. We assume

$\deg \overline{\mathbf{f}}_0 = 0$ and $\deg \overline{\mathbf{f}}_i = 2$ for $i = 1, \dots, m'$, $\deg \overline{\mathbf{f}}_i \geq 4$ for $i = m'+1, \dots, B'$. Then the divisor axiom of Gromov-Witten invariant (see [FO] Theorem 23.1.4) implies

$$\begin{aligned} &GW_{\ell+k}(\alpha; \underbrace{\overline{\mathbf{f}}_i, \dots, \overline{\mathbf{f}}_i}_{k\text{-times}}, Q_{k+1}, \dots, Q_{k+\ell}) \\ &= T^{\omega\cap\alpha/2\pi}(\alpha \cap \overline{\mathbf{f}}_i)^k GW_\ell(\alpha; Q_{k+1}, \dots, Q_{k+\ell}) \end{aligned} \tag{2.32}$$

for $i = 1, \dots, m'$. Now we consider $\mathfrak{b} = \sum_{i=1}^{B'} w_i \overline{\mathbf{f}}_i$ and put

$$\mathfrak{b}_+ = \sum_{i=m'+1}^{B'} \overline{w}_i \overline{\mathbf{f}}_i, \qquad \overline{\mathfrak{w}}_i = \exp \overline{w}_i \quad i = 1, \dots, m'.$$

Then (2.31) can be rewritten as

$$\begin{aligned} \langle \mathfrak{c} \cup^{\mathfrak{b}} \mathfrak{d}, \mathfrak{e} \rangle_{PD_X} = \sum_{\alpha \in H_2(X;\mathbb{Z})} \sum_{\ell=0}^{\infty} &\frac{1}{\ell!} T^{\omega\cap\alpha/2\pi} \overline{\mathfrak{w}}_1^{\alpha\cap\overline{\mathbf{f}}_1} \dots \overline{\mathfrak{w}}_{m'}^{\alpha\cap\overline{\mathbf{f}}_{m'}} \\ &GW_{\ell+3}(\alpha; \mathfrak{c}, \mathfrak{d}, \mathfrak{e}, \underbrace{\mathfrak{b}_{\text{high}}, \dots, \mathfrak{b}_{\text{high}}}_{\ell\text{-times}}). \end{aligned} \tag{2.33}$$

Remark 2.27. Since $\overline{\mathbf{f}}_0$ is a unit, the deformed quantum product $\cup^{\mathfrak{b}}$ is independent of w_0.

And as $\ell \to \infty$ in (2.33), we derive $c_1(X) \cap \alpha \to \infty$ from (2.29) and $\alpha \cap \omega \to \infty$ from Gromov compactness respectively. Therefore we obtain

Lemma 2.28. *There exist* $\lambda_i \to \infty$, $\lambda_i \geq 0$ *and*

$$P_i(\mathfrak{c}, \mathfrak{d}, \mathfrak{e}) \in \Lambda_0[\overline{\mathfrak{w}}_1, \overline{\mathfrak{w}}_1^{-1}, \dots, \overline{\mathfrak{w}}_{m'}, \overline{\mathfrak{w}}_{m'}^{-1}, \overline{w}_{m+1}, \dots, \overline{w}_{B'}]$$

such that

$$\langle \mathfrak{c} \cup^{\mathfrak{b}} \mathfrak{d}, \mathfrak{e} \rangle = \sum_i T^{\lambda_i} P_i(\mathfrak{c}, \mathfrak{d}, \mathfrak{e}).$$

We denote by

$$\Lambda_0 \langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w} \rangle\rangle$$

the completion of $\Lambda_0[\overline{\mathfrak{w}}_1, \overline{\mathfrak{w}}_1^{-1}, \dots, \overline{\mathfrak{w}}_{m'}, \overline{\mathfrak{w}}_{m'}^{-1}, \overline{w}_{m+1}, \dots, \overline{w}_{B'}]$ with respect to the non-Archimedean norm on Λ_0. Lemma 2.28 gives rise to a well-defined map

$$\cup^{\text{big}} : H(X; \Lambda_0) \otimes_{\Lambda_0} H(X; \Lambda_0) \to H(X; \Lambda_0) \otimes_{\Lambda_0} \Lambda_0 \langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w} \rangle\rangle$$

which induces a ring structure on

$$H(X; \Lambda_0 \langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w} \rangle\rangle) = H(X; \Lambda_0) \otimes_{\Lambda_0} \Lambda_0 \langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w} \rangle\rangle. \tag{2.34}$$

This finishes our description of the family of ring structures $\cup^{\mathfrak{b}}$ parameterized by the classes $\mathfrak{b} \in H(X; \Lambda_0)$.

We now specializes the above to the case of toric manifolds. The following theorem states that (2.34) is isomorphic to the Jacobian ring.

Theorem 2.29. *Let* X *be a compact toric manifold. Then, we have a ring isomorphism*

$$H(X; \Lambda_0 \langle\langle \overline{\mathfrak{w}}, \overline{\mathfrak{w}}^{-1}, \overline{w} \rangle\rangle) \cong \operatorname{Jac}'(\mathfrak{PO}).$$

See (2.13) for $\mathrm{Jac}'(\mathfrak{PO})$. The isomorphism in Theorem 2.29 can be described more explicitly. For this purpose it will be more convenient to fix $\mathfrak{b}$. We identify the tangent space

$$T_{\mathfrak{b}}H(X;\Lambda_0)$$

with $H(X;\Lambda_0)$. The coordinate basis of $T_{\mathfrak{b}}H(X;\Lambda_0)$ of the coordinates $(\overline{\mathfrak{w}},\overline{w})$ on $H(X;\Lambda_0)$ is given by the union

$$\left\{\frac{\partial}{\partial\overline{\mathfrak{w}}_i}\right\}_{i=1}^{m'}\cup\left\{\frac{\partial}{\partial\overline{w}_0}\right\}\cup\left\{\frac{\partial}{\partial\overline{w}_j}\right\}_{j=m'+1}^{B'}.$$

Definition 2.30. The *Kodaira-Spencer map*

$$\mathfrak{ks}_{\mathfrak{b}}:T_{\mathfrak{b}}H(X;\Lambda_0)\to\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$$

is defined by

$$\begin{aligned}
\mathfrak{ks}_{\mathfrak{b}}(\frac{\partial}{\partial\overline{\mathfrak{w}}_i})&\equiv\frac{\partial}{\partial\overline{\mathfrak{w}}_i}(\mathfrak{PO}_{\mathfrak{b}})\quad\mathrm{mod}\ \mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i\frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}\mid i=1,\dots,n\right),\\
\mathfrak{ks}_{\mathfrak{b}}(\frac{\partial}{\partial\overline{w}_i})&\equiv\frac{\partial}{\partial\overline{w}_i}(\mathfrak{PO}_{\mathfrak{b}})\quad\mathrm{mod}\ \mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i\frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}\mid i=1,\dots,n\right).
\end{aligned}$$

Remark 2.31. The map $\mathfrak{ks}$ is the singularity theory analogue of the Kodaira-Spencer map which is used to study deformations of complex structures.

The following is the main theorem of this paper.

Theorem 2.32. (1) *The Kodaira-Spencer map $\mathfrak{ks}_{\mathfrak{b}}$ induces a ring isomorphism*

$$\mathfrak{ks}_{\mathfrak{b}}:T_{\mathfrak{b}}H(X;\Lambda_0)\to\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}).$$

(2) *If $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function, then we have*

$$\langle\mathfrak{c},\mathfrak{d}\rangle_{\mathrm{PD}_X}=\langle\mathfrak{ks}_{\mathfrak{b}}\mathfrak{c},\mathfrak{ks}_{\mathfrak{b}}\mathfrak{d}\rangle_{\mathrm{res}}$$

for $\mathfrak{c},\mathfrak{d}\in T_{\mathfrak{b}}H(X;\Lambda_0)=H(X;\Lambda_0)$, where the left hand side is the Poincaré duality.

Proof of Theorem 1.3. Theorem 1.3 immediately follows from Theorem 2.32 (1), Proposition 2.15 in this present paper and Theorem 3.12, Proposition 8.21 in [FOOO5]. □

The following is immediate from Theorem 2.32 (1) and Proposition 2.15.

Corollary 2.33. *If $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function, then*

$$\#\mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})=\mathrm{rank}_{\Lambda}\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})=\mathrm{rank}_{\mathbb{Q}}H(X;\mathbb{Q}).$$

Sections 4-21 of this paper are devoted to the proof of Theorem 2.32.
We note that Theorem 2.32 implies the following mysterious identity:

Corollary 2.34. *If $\mathfrak{PO}^{\mathfrak{b}}$ is a Morse function, we have*

$$\sum_{(u,b)\in\mathfrak{M}(X,\mathfrak{b})}\frac{1}{Z(u,b)}=0. \tag{2.35}$$

Proof. Let $1_X \in H^0(X;\Lambda)$ be the unit. Then

$$\langle 1_X, 1_X\rangle_{\mathrm{PD}_X} = 0.$$

By Proposition 2.15 we have

$$1_X = \sum_{\mathfrak{y}\in\mathfrak{M}(X,\mathfrak{b})} 1_{\mathfrak{y}}$$

where $1_{\mathfrak{y}}$ is the unit of the Jacobian ring $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$. (2.35) now follows from (2.24) and Theorem 2.32 (2). □

Example 2.35. We take monotone toric blow up of $\mathbb{C}P^2$ at one point. Its potential function at the (unique) monotone fiber is:

$$T^{1/3}(y_1 + y_2 + (y_1y_2)^{-1} + y_1^{-1}). \tag{2.36}$$

See Example 8.1 [FOOO4]. The condition for (y_1, y_2) to be critical is

$$1 - y_1^{-2}y_2^{-1} - y_1^{-2} = 0, \quad 1 - y_1y_2^2 = 0. \tag{2.37}$$

Therefore $y_1y_2 = 1/y_2$ and

$$y_2^4 + y_2^3 - 1 = 0. \tag{2.38}$$

By Theorem 2.24 (3) we have

$$Z(0,(y_1,y_2)) = T^{2/3}\det\begin{bmatrix} y_1 + (y_1y_2)^{-1} + y_1^{-1} & (y_1y_2)^{-1} \\ (y_1y_2)^{-1} & y_2 + (y_1y_2)^{-1}\end{bmatrix} \tag{2.39}$$

Using (2.37), (2.38) we have

$$\begin{aligned} T^{-2/3}Z(0,(y_1,y_2)) &= \det\begin{bmatrix} y_2 + y_2^2 + y_2^{-2} & y_2 \\ y_2 & 2y_2\end{bmatrix} \\ &= 2y_2^3 + y_2^2 + 2y_2^{-1} = \frac{4 - y_2^3}{y_2}. \end{aligned} \tag{2.40}$$

Let z_i $(i = 1,2,3,4)$ be the 4 solutions of (2.38). Then the left hand side of (2.35) is:

$$T^{-2/3}\sum_{i=1}^4 \frac{z_i}{4 - z_i^3}. \tag{2.41}$$

We have $z^4 + z^3 - 1 = -(z+1)(4 - z^3) + 4z + 3$. Therefore (2.41) is

$$\begin{aligned} T^{-2/3}\sum_{i=1}^4 \frac{z_i(z_i+1)}{4z_i+3} &= \frac{T^{-2/3}}{16}\left(\sum_{i=1}^4 (4z_i+1) - \sum_{i=1}^4 \frac{3}{4z_i+3}\right) \\ &= -\frac{3T^{-2/3}}{16}\sum_{i=1}^4 \frac{1}{4z_i+3}. \end{aligned}$$

We use $\sum z_i = -1$ in the second equality.

We have $4(z^4+z^3-1) = (z^3 + z^2/4 - 3z/16 + 9/64)(4z+3) - 283/64$. Therefore

$$\sum_{i=1}^4 \frac{1}{4z_i+3} = \frac{64}{283}\sum_{i=1}^4\left(z_i^3 + \frac{z_i^2}{4} - \frac{3z_i}{16} + \frac{9}{64}\right).$$

By (2.38) we have $\sum z_i^3 = \sum z_i = -1$, $\sum z_i^2 = 1$. Therefore

$$\frac{64}{283}\sum_{i=1}^4\left(z_i^3 + \frac{z_i^2}{4} - \frac{3z_i}{16} + \frac{9}{64}\right) = \frac{64}{283}\left(-1 + \frac{1}{4} + \frac{3}{16} + 4\times\frac{9}{64}\right) = 0.$$

We have thus checked (2.35) directly in the case of monotone toric one point blow up of $\mathbb{C}P^2$.

3. PROJECTIVE SPACE: AN EXAMPLE

Before starting the proof of Theorem 2.32, we illustrate the theorem for the case $X = \mathbb{C}P^n$ with $\mathfrak{b} = \mathbf{0}$ by doing some explicit calculations. In this case the moment polytope P is

$$\Big\{(u_1, \ldots, u_n) \mid 0 \le u_i, \sum u_i \le 1\Big\}$$

and the potential function

$$\mathfrak{PO}_{\mathbf{0}} = \sum_{i=1}^{n} T^{u_i} y_i + T^{1-\sum u_i}(y_1 \cdots y_n)^{-1}.$$

The critical points are $\mathfrak{y}^{(k)} = T^{\frac{1}{n+1}} e^{\frac{2\pi\sqrt{-1}k}{n+1}}$ $k = 0, \ldots, n$ which are all nondegenerate. Therefore we have

$$\operatorname{Jac}(\mathfrak{PO}_{\mathbf{0}}) \otimes_{\Lambda_0} \Lambda \cong \prod_{k=0}^{n} \Lambda 1_{\mathfrak{y}^{(k)}}.$$

The isomorphism is induced by

$$\Lambda^P \langle\langle y, y^{-1} \rangle\rangle \to \prod_{k=0}^{n} \Lambda 1_{\mathfrak{y}^{(k)}}, \quad \mathcal{P} \mapsto \sum_{k=0}^{n} \mathcal{P}(\mathfrak{y}^{(k)}) 1_{\mathfrak{y}^{(k)}}.$$

If we put

$$\mathbf{f}_k = \pi^{-1}(\{(u_1, \ldots, u_n) \in P \mid u_i = 0, \, i = n-k+1, \ldots, n\}),$$

$\{\mathbf{f}_0, \ldots, \mathbf{f}_n\}$ forms a basis of $H(\mathbb{C}P^n; \Lambda)$.

We derive

$$\mathfrak{PO}_{w\mathbf{f}_1}(y) = \mathfrak{PO}_{\mathbf{0}}(y) + (e^w - 1) T^{u_n} y_n$$

from Proposition 4.9 [FOOO5] and hence

$$\mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_1) = [T^{u_n} y_n] = T^{\frac{1}{n+1}} \sum_{k=0}^{n} e^{\frac{2\pi\sqrt{-1}k}{n+1}} 1_{y^{(k)}} \tag{3.1}$$

by definition of $\mathfrak{ks}_{\mathbf{0}}$. Using the fact that $\mathfrak{ks}_{\mathbf{0}}$ is a ring homomorphism, we have

$$\mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_\ell) = T^{\frac{\ell}{n+1}} \sum_{k=0}^{n} e^{\frac{2\pi\sqrt{-1}k\ell}{n+1}} 1_{\mathfrak{y}^{(k)}}. \tag{3.2}$$

Note this holds for $\ell = 0$ also since $\mathbf{f}_0$ is a unit and $\mathfrak{ks}_{\mathbf{0}}$ is unital. We remark that

$$(\mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_1))^{n+1} = T \sum_{k=0}^{n} 1_{\mathfrak{y}^{(k)}} = T(\mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_0)).$$

Recall that

$$(\mathbf{f}_1)^{n+1} = T\mathbf{f}_0$$

is the fundamental relation of the small quantum cohomology ring of $\mathbb{C}P^n$.

We now calculate the residue pairing. We remark that both (1) and (2) of Condition 2.19 hold in this case. The Hessian of $\mathfrak{PO}_{\mathbf{0}}$ is given by

$$\mathrm{Hess}_{\mathfrak{y}^{(k)}}\mathfrak{PO}_{\mathbf{0}} = \left[T^{\frac{1}{n+1}}\frac{\partial^2}{\partial x_i \partial x_j}\left(e^{x_1}+\cdots+e^{x_n}+e^{-(x_1+\cdots+x_n)}\right)\right]_{i,j=1}^{i,j=n}(\mathfrak{x}^{(k)})$$

with $\mathfrak{x}_i^{(k)} = \frac{2\pi\sqrt{-1}k}{n+1}$. Therefore

$$\mathrm{Hess}_{\mathfrak{y}^{(k)}}\mathfrak{PO}_{\mathbf{0}} = T^{\frac{1}{n+1}}e^{\frac{2\pi\sqrt{-1}k}{n+1}}\left[\delta_{ij}+1\right]_{i,j=1}^{i,j=n}.$$

It is easy to see that the determinant of the matrix $[\delta_{ij}+1]_{i,j=1}^{i,j=n}$ is $n+1$. Therefore the residue pairing is given by

$$\langle 1_{\mathfrak{y}^{(k)}}, 1_{\mathfrak{y}^{(k')}}\rangle_{\mathrm{res}} = T^{-\frac{n}{n+1}}e^{-\frac{2\pi\sqrt{-1}kn}{n+1}}\frac{\delta_{kk'}}{1+n}. \tag{3.3}$$

Combining (3.2) and (3.3), we obtain

$$\langle \mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_\ell), \mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_{\ell'})\rangle_{\mathrm{res}} = \frac{1}{n+1}T^{-\frac{n}{n+1}}\sum_{k=0}^{n} e^{-\frac{2\pi\sqrt{-1}kn}{n+1}}T^{\frac{\ell+\ell'}{n+1}}e^{\frac{2\pi\sqrt{-1}(\ell+\ell')k}{n+1}}. \tag{3.4}$$

It follows that (3.4) is 0 unless $\ell+\ell' = n$ and

$$\langle \mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_\ell), \mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_{n-\ell})\rangle_{\mathrm{res}} = 1 = \langle \mathbf{f}_\ell, \mathbf{f}_{n-\ell}\rangle_{\mathrm{PD}_{\mathbb{C}P^n}}.$$

Thus Theorem 2.32 (2) holds in this case.

We remark that in this case a similar calculation can be found in the earlier literature [Ta], [Bar], [KKP]. The main result of the present paper generalizes this calculation to arbitrary compact toric manifolds X also with arbitrary bulk deformation parameter $\mathfrak{b} \in H(X;\Lambda_0)$.

Remark 3.1. In the above discussion of projective spaces, we derived (3.2) from (3.1) and the fact $\mathfrak{ks}_{\mathbf{0}}$ is a ring homomorphism. We can also prove (3.2) directly: Let $\beta_i \in \pi_2(X; L(u))$ as in the end of Section 1. We consider the moduli space $\mathcal{M}_{1;1}(\beta_{n-\ell+1}+\cdots+\beta_n)$ and its fiber product

$$\mathcal{M}_1(\beta_{n-\ell+1}+\cdots+\beta_n;\mathbf{f}_\ell) := \mathcal{M}_{1;1}(\beta_{n-\ell+1}+\cdots+\beta_n)_{\mathrm{ev}^{\mathrm{int}}}\times_X \mathbf{f}_\ell.$$

Since the Maslov index of $\beta_{n-\ell+1}+\cdots+\beta_n$ is 2ℓ, it follows that the (virtual) dimension of $\mathcal{M}_1(\beta_{n-\ell+1}+\cdots+\beta_n;\mathbf{f}_\ell)$ is n: Actually by inspecting the classification of holomorphic discs, Theorem 5.2 [CO] (see also Theorem 4.4 [Cho1]) we can find that $\mathcal{M}_{1;1}(\beta_{n-\ell+1}+\cdots+\beta_n)$ is Fredholm regular and the evaluation map

$$\mathrm{ev}_{\mathrm{int}} : \mathcal{M}_{1;1}(\beta_{n-\ell+1}+\cdots+\beta_n) \to X$$

is transverse to the divisor $\mathbf{f}_\ell$. Therefore $\mathcal{M}_1(\beta_{n-\ell+1}+\cdots+\beta_n;\mathbf{f}_\ell)$ is smooth, n dimensional and

$$\mathrm{ev} : \mathcal{M}_1(\beta_{n-\ell+1}+\cdots+\beta_n;\mathbf{f}_\ell) \to L(u)$$

is a diffeomorphism. We can also find that $\mathcal{M}_1(\beta;\mathbf{f}_\ell)$ is empty for other β with Maslov index 2ℓ. Therefore by definition we have

$$\mathfrak{PO}(w\mathbf{f}_\ell; y) - \mathfrak{PO}(\mathbf{0}; y) \equiv wT^{u_{n-\ell+1}+\cdots+u_n}y_{n-\ell+1}\cdots y_n \mod w^2.$$

Hence

$$\mathfrak{ks}_{\mathbf{0}}(\mathbf{f}_\ell) = \sum_{k=0}^{n}\left.\frac{\partial \mathfrak{PO}_{w\mathbf{f}_\ell}}{\partial w}\right|_{w=0}(\mathfrak{y}^{(k)})1_{\mathfrak{y}^{(k)}}$$

is given by (3.2).

Remark 3.2. In [Gro] the case of $\mathbb{C}P^2$ is studied including bulk deformation by $\mathfrak{b} \in H^4(\mathbb{C}P^2)$.

Part 1. Ring isomorphism

4. NORMS AND COMPLETIONS OF POLYNOMIAL RING OVER NOVIKOV RING

In this section we discuss various points related to the ring $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ which was defined to be

$$\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle = \bigcap_{u\in \mathrm{Int} P} \Lambda_0^{u}\langle\langle y, y^{-1}\rangle\rangle$$

in Section 2, and prove Lemma 2.3 and Lemma 2.13 whose proofs were postponed therein. This section is rather technical since we need to deal with completion of the ring of Laurent polynomials with Novikov ring coefficients. In particular, we need to make precise the definition of a metric on $\Lambda_0[y, y^{-1}]$ used taking the completion which coincides with $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.

For each sufficiently small $\varepsilon > 0$, we define the subset P_ε of P by

$$P_\varepsilon = \{u \in P \mid \ell_j(u) \geq \varepsilon,\ j = 1, \dots, m\}.$$

Here ℓ_j are affine functions appeared in (2.6).

Definition 4.1. We define $\Lambda^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle$ by the completion of $\Lambda[y, y^{-1}]$ with respect to the *bounded* metric

$$d_{P_\varepsilon}(x, x') := \min\{1, \exp(-\mathfrak{v}_T^{P_\varepsilon}(x - x'))\}$$

on $\Lambda_0[y, y^{-1}]$. And $\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle$ is the subring of elements x with $\mathfrak{v}_T^{P_\epsilon}(x) \geq 0$.

Definition 4.2. Define a metric $d_{\overset{\circ}{P}}$ on $\Lambda[y, y^{-1}]$ by

$$d_{\overset{\circ}{P}}(x, x') = \sum_{k=1}^{\infty} 2^{-k} d_{P_{1/k}}(x, x'). \tag{4.1}$$

The case of $\Lambda_0^{\overset{\circ}{P}}\langle\langle \mathfrak{w}, \mathfrak{w}^{-1}, w, y, y^{-1}\rangle\rangle$ is similar.

Proposition 4.3. *$\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ is the completion of a subring*

$$\Lambda[y, y^{-1}] \cap \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$$

of the Laurent polynomial ring with respect to the metric $d_{\overset{\circ}{P}}$.

Proof. We note that if $\varepsilon > 0$ is sufficiently small P_ε is still convex and so

$$\mathfrak{v}_T^{P_\varepsilon} = \min\{\mathfrak{v}_T^{(u(\varepsilon,j))} \mid j = 1, \dots, N\} \tag{4.2}$$

where $u(\varepsilon, 1), \dots, u(\varepsilon, N)$ are the vertices of P_ε. Therefore we have

$$\bigcap_{u\in \mathrm{Int} P} \Lambda_0^{u}\langle\langle y, y^{-1}\rangle\rangle = \bigcap_{k=1}^{\infty} \Lambda_0^{P_{1/k}}\langle\langle y, y^{-1}\rangle\rangle$$

and hence

$$\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle = \bigcap_{k=1}^{\infty} \Lambda_0^{P_{1/k}}\langle\langle y, y^{-1}\rangle\rangle.$$

But it easily follows from the definition of the metric $d_{\overset{\circ}{P}}$ that the completion of $\Lambda[y, y^{-1}] \cap \Lambda_0^{\overset{\circ}{P}} \langle\langle y, y^{-1} \rangle\rangle$ with respect to $d_{\overset{\circ}{P}}$ coincides with

$$\bigcap_{k=1}^{\infty} \Lambda_0^{P_{1/k}} \langle\langle y, y^{-1} \rangle\rangle$$

whose proof is standard and so omitted. ☐

Note that there exists a continuous homomorphism

$$\Lambda_0^{P_{\varepsilon_1}} \langle\langle y, y^{-1} \rangle\rangle \to \Lambda_0^{P_{\varepsilon_2}} \langle\langle y, y^{-1} \rangle\rangle$$

for $\varepsilon_1 < \varepsilon_2$. The following lemma is an immediate consequence of Proposition 4.3 and (4.2). Note by

Lemma 4.4. *$\Lambda_0^{\overset{\circ}{P}} \langle\langle y, y^{-1} \rangle\rangle$ is isomorphic to the projective limit $\varprojlim \Lambda_0^{P_\varepsilon} \langle\langle y, y^{-1} \rangle\rangle$.*

The following lemma is a consequence of the proof of Lemma 3.9 [FOOO5] whose proof we omit but refer thereto.

Lemma 4.5. *Suppose $T^\mu y^v$ lies in $\Lambda_0^P[y, y^{-1}]$. Then there exist $w(v) \in \mathbb{Z}_{\geq 0}^m$ such that*

$$T^\mu y^v = T^{\lambda(\mu,v)} z^{w(v)} \tag{4.3}$$

for $z^{w(v)} = z_1^{w(v)^{(1)}} \dots z_m^{w(v)^{(m)}}$ and

$$\mathfrak{v}_T^P(T^\mu y^v) = \lambda(\mu, v). \tag{4.4}$$

Now we are ready to give the proofs of Lemmas 2.3 and 2.13.

Proof of Lemma 2.3. The case of $\Lambda_0^P \langle\langle y, y^{-1} \rangle\rangle$ immediately follows from Lemma 3.9 [FOOO5] and so we will focus on the case of $\Lambda_0^{\overset{\circ}{P}} \langle\langle y, y^{-1} \rangle\rangle$.

By definition of z_j's in (2.7), we have $\mathfrak{v}_T^u(z_j) = \ell_j(u)$. In particular, if $u \in \operatorname{Int} P$, we have $\mathfrak{v}_T^u(z_j) > 0$. Therefore any formal power series of z_j with coefficient in Λ_0 converges in $\mathfrak{v}_T^u$ for any $u \in \operatorname{Int} P$. Hence the map

$$\Lambda_0[[Z_1, \dots, Z_m]] \to \Lambda_0^{\overset{\circ}{P}} \langle\langle y, y^{-1} \rangle\rangle \tag{4.5}$$

is well defined. It remains to prove its surjectivity.

Now let x be an arbitrary element $\Lambda_0^{\overset{\circ}{P}} \langle\langle y, y^{-1} \rangle\rangle$. Then for any $u \in \operatorname{Int} P$ we can write x as

$$x = \sum_{i=1}^{\infty} a_i T^{\lambda'_i} y(u)^{v_i}$$

of with $0 \leq \lambda'_i \leq \lambda'_{i+1}$, $\lim \lambda'_i \to \infty$. (Recall $\mathfrak{v}_T^u(y_j(u)) = 0$.) We assume that $(\lambda'_i, v_i) \neq (\lambda'_j, v_j)$, and $a_i \in \mathbb{C} \setminus \{0\}$. We define

$$\lambda_i = \lambda(\lambda'_i, u, v_i) = \mathfrak{v}_T^P(T^{\lambda'_i} y(u)^{v_i}), \quad w_i = w(v_i) = (w_i^{(1)}, \dots, w_i^{(m)}).$$

For each $N \in \mathbb{N}$, we define

$$I_N = \{i \mid w_i^{(1)} + \dots + w_i^{(m)} \leq N\},$$

and

$$x_N = \sum_{i \in I_N} a_i T^{\lambda'_i} y(u)^{v_i}. \tag{4.6}$$

A priori, (4.6) may be an infinite sum but the sum converges in $\mathfrak{v}_T^u$ norm for any $u \in \operatorname{Int} P$: It is a sub-series of a convergent series. Therefore the sum converges in d_{P_ε} for any $\varepsilon > 0$ by $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$, since v_T^u is an *non-Archimedean* valuation for any u. Therefore it converges in $d_{\overset{\circ}{P}}$ by the definition (4.1).

Using (4.3), we can rewrite

$$\sum_{i\in I_N} a_i T^{\lambda'_i} y(u)^{v_i} = \sum_{i \in I_N} a_i T^{\lambda_i} z^{w_i}$$

and so the right hand side also converges in d_{P_ϵ}. On the other hand, if $|w_i^{(1)} + \cdots + w_i^{(m)}| \le N$, we derive

$$|\mathfrak{v}_T^{P_\epsilon}(a_i T^{\lambda_i} z^{w_i}) - \mathfrak{v}_T^{P}(a_i T^{\lambda_i} z^{w_i})| \le \epsilon N$$

from (4.2) and hence the sum also converges in d_P and hence $x_N \in \Lambda_T^P\langle\!\langle y, y^{-1}\rangle\!\rangle$. In particular,

$$\{i \mid \lambda_i < C, w_i^{(1)} + \cdots + w_i^{(m)} < N\}$$

is a finite set for any C and N. Therefore we have proved that

$$\sum_{i\in I_N} a_i T^{\lambda_i} z^{w_i}$$

is a polynomial with $\lambda_i \ge 0$: Recall $\lambda_i = \mathfrak{v}_T^P(T^{\lambda'_i} y(u)^{v_i}) = \mathfrak{v}_T^P(T^{\lambda_i} z^{w_i}) \ge 0$. By construction, we have

$$x = \lim_{N\to\infty} x_N$$

with respect to the distance $d_{\overset{\circ}{P}}$ and so x is the image of the formal power series

$$\sum_{i \in \cup_{N=0}^\infty I_N} a_i T^{\lambda_i} Z^{w_i}.$$

This implies the surjectivity of (4.5).

The proof of the other two cases are similar and is left to the reader. □

Proof of Lemma 2.13. (A) (1) ⇒ (2) is immediate from Lemma 2.3.

We next prove (2) ⇒ (1). Recall from Definition 2.2 that $z_j = T^{-\lambda_j}\prod_{i=1}^n y_i^{v_{j,i}}$ $(j = 1, \dots, m)$. By the homomorphism $y_i \to \mathfrak{y}_i$, z_j is mapped to $\mathfrak{z}_j = T^{-\lambda_j}\prod_{i=1}^n \mathfrak{y}_i^{v_{j,i}}$. For each $j = 1, \dots, m$, $\sum_{k=1}^\infty z_j^k$ converges in $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$. This is because $\mathfrak{v}_T^u(z_j) = \ell_i(u) > 0$ for any $u \in \text{Int } P$. By the continuity hypothesis of the evaluation homomorphism substituting $y \mapsto \mathfrak{y}$, it follows that $\sum_{k=0}^\infty \mathfrak{z}_j^k$ converges in Λ_0. Therefore

$$\mathfrak{v}_T(\mathfrak{z}_j) > 0.$$

Since we have $\mathfrak{z}_j = T^{-\lambda_j}\prod_{i=1}^n \mathfrak{y}_i^{v_{j,i}}$ from (2.7), we have $0 < \mathfrak{v}_T(\mathfrak{z}_j) = -\lambda_j + \langle v_j, \vec{u}\rangle = \ell_j(\vec{u})$ for all $j = 1, \dots, m$ with $\vec{u} = \vec{\mathfrak{v}}_T(\mathfrak{y})$. By the definition of the moment polytope P, $\vec{u} \in \text{Int } P$ which implies (2.15).

(B) (1) ⇒ (2) also follows from Lemma 2.3. To prove (2) ⇒ (1), we consider a series $\sum_{k=1}^\infty T^{\epsilon k} z_j^k$ for $\epsilon > 0$. Then it converges in $\Lambda_0^P\langle\!\langle y, y^{-1}\rangle\!\rangle$. Then a similar argument above yields that $\epsilon + \mathfrak{v}_T(\mathfrak{z}_j) > 0$. Since $\epsilon > 0$ can be taken sufficiently small, we have $v_T(\mathfrak{z}_j) \ge 0$. Then the similar argument shows (1). □

Definition 4.6. We define

$$\Lambda^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle = \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle \otimes_{\Lambda_0} \Lambda.$$

Remark 4.7. We recall that in Section 2 we define $\Lambda^P\langle\langle y, y^{-1}\rangle\rangle$ by the completion of $\Lambda[y, y^{-1}]$ with respect to $\mathfrak{v}_T^P$. By considering the homomorphism (2.8) over Λ coefficients, we find that $\Lambda^P\langle\langle y, y^{-1}\rangle\rangle$ coincides with the image of

$$\Lambda\langle\langle Z_1, \dots, Z_m\rangle\rangle \to \Lambda^P\langle\langle y, y^{-1}\rangle\rangle. \tag{4.7}$$

We find that

$$\Lambda^P\langle\langle y, y^{-1}\rangle\rangle = \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle \otimes_{\Lambda_0} \Lambda.$$

Similarly, we also find that

$$\Lambda^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle = \Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle \otimes_{\Lambda_0} \Lambda.$$

However, we note that (4.7) does not extend to $\Lambda[[Z_1, \dots, Z_m]] \to \Lambda^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.

5. LOCALIZATION OF JACOBIAN RING AT MOMENT POLYTOPE

In this section, we give the proof of Proposition 2.15. We will use the following lemma for the proof.

Lemma 5.1. $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda$ *is a finite dimensional* Λ *vector space.*

Lemma 5.1 is an immediate consequence of Theorem 2.32 (1). (We will use Proposition 2.15 to define residue pairing on Jacobian ring. The residue pairing will be used to state Theorem 2.32 (2). However the proof of Theorem 2.32 (1) will be completed, without using Proposition 2.15, in Section 10 when the surjectivity of $\mathfrak{ks}_{\mathfrak{b}}$ is established. Thus the logic of arguments used in the proofs will not be circular.)

Proof of Proposition 2.15. Let $\mathcal{P} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$. If $\mathfrak{y}$ is a critical point of $\mathfrak{PO}_{\mathfrak{b}}$ then by Lemma 2.13, we can plug in $\mathfrak{y}_i = y_i$ to $\mathcal{P}$ and obtain $\mathcal{P}(\mathfrak{y})$. Since $\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}(\mathfrak{y}) = 0$ if $\mathfrak{y}$ is a critical point, we obtain a homomorphism $\Phi_{\mathfrak{y}} : \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \to \Lambda$ for each critical point $\mathfrak{y}$. We will prove that $\prod_{\mathfrak{y}\in \mathrm{Crit}(\mathfrak{PO}_\beta)} \Phi_{\mathfrak{y}}$ gives the isomorphism (2.16) when $\mathfrak{PO}_\beta$ is a Morse function.

For $y_i \in \Lambda^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ we define a Λ linear map

$$\widehat{y_i} : \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \to \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda$$

as the multiplication by y_i in the Jacobian ring. Since $y_i^{-1} \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ and the map $y_i \mapsto \widehat{y_i}$ is a ring homomorphism from $\Lambda^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ to $\mathrm{End}_\Lambda(\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda)$, $\widehat{y_i}$ is invertible.

For $\mathfrak{y} = (\mathfrak{y}_1, \dots, \mathfrak{y}_n) \in \Lambda^n$ we put

$$\begin{aligned}\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}; \mathfrak{y}) \quad = \quad & \{\mathcal{P} \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \mid \\ & (\widehat{y_i} - \mathfrak{y}_i)^N \mathcal{P} = 0 \ i = 1, \dots, n \text{ for sufficiently large } N\}. \end{aligned} \tag{5.1}$$

Since $\widehat{y_i}$ is invertible, if $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}; \mathfrak{y}) \ne 0$, $\mathfrak{y}_i \ne 0$. We denote

$$\mathfrak{Y} = \{\mathfrak{y} \mid \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}; \mathfrak{y}) \ne 0\}.$$

We have a decomposition

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \cong \prod_{\mathfrak{y}\in\mathfrak{Y}} \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}; \mathfrak{y}) \tag{5.2}$$

as a Λ vector space. Using the fact $\widehat{y}_i\widehat{y}_j = \widehat{y}_j\widehat{y}_i$ we can prove that (5.2) is a direct product decomposition as a ring. (See the proof of Proposition 6.7 [FOOO4].)

Lemma 5.2. *If* $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y) \neq 0$*, then* $\vec{\mathfrak v}_T(\mathfrak y) \in \mathrm{Int}\, P$.

Proof. Let $u = \vec{\mathfrak v}_T(\mathfrak y)$. If $u \notin \mathrm{Int}\, P$ then there exists j such that $\ell_j(u) \le 0$. There exists nonzero $\mathcal P$ such that $y_i\mathcal P = \mathfrak y_i\mathcal P$ for $i = 1,\dots,n$. We have

$$(z_j)^k\mathcal P = T^{-k\lambda_j} y_1^{kv_{j,1}}\cdots y_n^{kv_{j,n}}\mathcal P = T^{-k\lambda_j}\mathfrak y_1^{kv_{j,1}}\cdots \mathfrak y_n^{kv_{j,n}}\mathcal P \tag{5.3}$$

Since $\lim_{k\to\infty} z_j^k = 0$ in $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ and

$$\lim_{k\to\infty}\mathfrak v_T((T^{-k\lambda_j}\mathfrak y_1^{kv_{j,1}}\cdots \mathfrak y_n^{kv_{j,n}})^{-1}) = -\lim_{k\to\infty} k\ell_j(u) \ge 0,$$

it follows that

$$(z_j)^k(T^{-k\lambda_j}\mathfrak y_1^{kv_{j,1}}\cdots \mathfrak y_n^{kv_{j,n}})^{-1}$$

converges to 0 in $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$. Therefore (5.3) implies $\mathcal P = 0$. This is a contradiction. □

Lemma 5.3. $\mathfrak Y = \mathrm{Crit}(\mathfrak{PO}_{\mathfrak b})$.

Proof. Let $\mathfrak y \in \mathfrak Y$. Then there exists nonzero $\mathcal P \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y)$ such that $y_i\mathcal P = \mathfrak y_i\mathcal P$ for $i = 1,\dots,n$. It follows that $y_i \mapsto \mathfrak y_i$ induces a non-zero homomorphism $\Phi_{\mathfrak y} : \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \to \Lambda$. Therefore (2.14) is satisfied. (2.15) follows from Lemma 5.2.

We next assume $\mathfrak y \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak b})$. Then, by Lemma 2.13 and the definition, $y_i \mapsto \mathfrak y_i$ induces a homomorphism $\Phi_{\mathfrak y} : \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \to \Lambda$. For given $\mathfrak y' = (\mathfrak y'_1,\dots,\mathfrak y'_n) \in \mathfrak Y$, let $1_{\mathfrak y'}$ be the unit of $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y')$. By definition, we have $(y_i - \mathfrak y'_i)^N 1_{\mathfrak y'} = 0$ for sufficiently large N and so we have $0 = \Phi_{\mathfrak y}((y_i - \mathfrak y'_i)^N 1_{\mathfrak y'}) = (\mathfrak y_i - \mathfrak y'_i)^N\Phi_{\mathfrak y}(1_{\mathfrak y'})$. Therefore if $\mathfrak y \neq \mathfrak y'$, we obtain $\Phi_{\mathfrak y}(1_{\mathfrak y'}) = 0$. Therefore the factorization (5.2) implies $\mathrm{Crit}(\mathfrak{PO}_{\mathfrak b}) \subset \mathfrak Y$ for otherwise we would have $\Phi_{\mathfrak y} \equiv 0$, a contradiction. The proof of Lemma 5.3 is complete. □

Lemma 5.4. $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y)$ *is a local ring.*

Proof. For any homomorphism $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y) \to \Lambda$, the element $y_i 1_{\mathfrak y}$ must be sent to $\mathfrak y_i$. The lemma then follows form the fact that $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y)$ is generated by $y_i 1_{\mathfrak y}$ as a Λ algebra. □

We remark that so far we have not used the assumption that $\mathfrak{PO}_{\mathfrak b}$ is a Morse function.

Lemma 5.3 and (5.2) imply that $\prod_{\mathfrak y \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak b})}\Phi_{\mathfrak y}$ is surjective. To prove its injectivity it suffices to show the following.

Lemma 5.5. $\mathfrak y$ *is a nondegenerate critical point of* $\mathfrak{PO}_{\mathfrak b}$ *if and only if*

$$\dim_\Lambda \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y) = 1.$$

Proof. First assume $\dim_\Lambda \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y) > 1$. By an elementary linear algebra we can find $\mathcal P \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y)$ and i_0 with the following properties:

1. $(y_{i_0} - \mathfrak y_{i_0})\mathcal P \neq 0$.
2. $(y_j - \mathfrak y_j)\mathcal P = 0$ for $j \neq i_0$.
3. $(y_{i_0} - \mathfrak y_{i_0})^2\mathcal P = 0$.

We will show by contradiction that $\mathbf{e}_{i_0} = (0,\dots,0,\overset{i_0\text{-th}}{1},0,\dots,0)$ cannot be contained in the image of the Hessian matrix $\mathrm{Hess}_{\mathfrak{y}}\mathfrak{PO}_{\mathfrak{b}}$.

Suppose to the contrary that there exists a vector $\vec{c} = (c_1,\dots,c_n)$ with $c_i \in \Lambda$ such that

$$\mathbf{e}_{i_0} = \left(\frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial y_j \partial y_k}(\mathfrak{y})\right)\vec{c}. \tag{5.4}$$

In other words, we have

$$dy_{i_0}(\mathfrak{y}) = d\left(\sum_{k=1}^{n} c_k \frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_k}\right)(\mathfrak{y}).$$

Integrating this, we obtain

$$y_{i_0} - \mathfrak{y}_{i_0} = \mathcal{Q} + \sum_{k=1}^{n} c_k \frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_k}(y) \tag{5.5}$$

for $\mathcal{Q} \in \Lambda\langle\langle y, y^{-1}\rangle\rangle$ such that

$$\mathcal{Q}(\mathfrak{y}) = 0, \quad d\mathcal{Q}(\mathfrak{y}) = 0. \tag{5.6}$$

This implies $\mathcal{Q} = \sum_{j_1,j_2=1}^{n}(y_{j_1} - \mathfrak{y}_{j_1})(y_{j_2} - \mathfrak{y}_{j_2})\mathcal{R}_{j_1,j_2}$ for some $\mathcal{R}_{j_1,j_2} \in \Lambda\langle\langle y, y^{-1}\rangle\rangle$. Therefore Properties (2), (3) of $\mathcal{P}$ above imply $\mathcal{P}\mathcal{Q} = 0$ in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$. Thanks to (5.5), this in turn implies $\mathcal{P}(y_{i_0} - \mathfrak{y}_{i_0}) = 0$. This contradicts to Property (1) and so the Hessian matrix $\mathrm{Hess}_{\mathfrak{y}}\mathfrak{PO}_{\mathfrak{b}}$ cannot be nondegenerate.

We next prove the converse. Suppose $\dim_\Lambda \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y}) = 1$ and $0 \neq \mathcal{P} \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ such that $(y_i - \mathfrak{y}_i)\mathcal{P} = 0$ for all $i = 1,\dots,n$. Using the finite dimensionality of $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ over Λ and Lemma 5.4, there exists a sufficiently large N such that whenever $\mathcal{P}(\mathfrak{y}') = 0$, $\mathcal{P}^N = 0$ on $\mathrm{Jac}_{\mathfrak{b}}(\mathfrak{PO};\mathfrak{y}')$. Using this fact, we find a $\mathcal{P} \in \overset{\circ}{\Lambda}{}^{P}\langle\langle y, y^{-1}\rangle\rangle$ such that $\mathcal{P} = 0$ in $\mathrm{Jac}_{\mathfrak{b}}(\mathfrak{PO};\mathfrak{y}')$ for all $\mathfrak{y}' \neq \mathfrak{y}$ and $\mathcal{P}(\mathfrak{y}) \neq 0$. By the hypothesis $\dim_\Lambda \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y}) = 1$, we conclude $(y_i - \mathfrak{y}_i)\mathcal{P}(\mathfrak{y}) = 0$ for $i = 1,\dots,n$ in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$.

Therefore it follows from (5.2) that for each i there exists some vector $\vec{\mathcal{G}}_i = (\mathcal{G}_{i,1},\dots,\mathcal{G}_{i,n})$ with $\mathcal{G}_{i,k} \in \overset{\circ}{\Lambda}{}^{P}\langle\langle y, y^{-1}\rangle\rangle$ such that

$$(y_i - \mathfrak{y}_i)\mathcal{P} = \sum_{k=1}^{n} \mathcal{G}_{i,k} \frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_k}. \tag{5.7}$$

By substituting $y = \mathfrak{y}$ into the y_i-derivative of (5.7), we obtain

$$\begin{aligned} 0 \neq \mathcal{P}(\mathfrak{y}) &= \sum_{k=1}^{n} c_k \frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial y_k \partial y_i}(\mathfrak{y}), \\ 0 &= \sum_{k=1}^{n} c_k \frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial y_k \partial y_j}(\mathfrak{y}) \qquad j \neq i \end{aligned}$$

for $c_k = \mathcal{G}_k(\mathfrak{y})$. This implies that $\mathbf{e}_i$ is in the image of the Hessian matrix at $\mathfrak{y}$. Since i is arbitrary for $1,\dots,n$, the Hessian at $\mathfrak{y}$ is nondegenerate, which finishes the proof. □

Now the proof of Proposition 2.15 is complete. □

Proposition 5.6. *We have*

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \cong \frac{\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle}{\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1, \dots, n\right)}, \tag{5.8}$$

for sufficiently small $\epsilon > 0$. If $\mathfrak b \in \mathcal A(\Lambda_+)$ then we have

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \cong \frac{\Lambda_0^{P}\langle\langle y, y^{-1}\rangle\rangle}{\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1, \dots, n\right)}. \tag{5.9}$$

Proof. There is a continuous homomorphism

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b}) \to \frac{\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{d^{P_\epsilon}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1, \dots, n\right)}, \tag{5.10}$$

where the closure $\mathrm{Clos}_{d^{P_\epsilon}}$ is taken with respect to the metric d^{P_ϵ}. Surjectivity of (5.10) follows from Lemma 12.8 (1). (We may apply this lemma to $P = P_\epsilon$.) To prove the injectivity of (5.10) it suffices to prove that it is injective after taking $\otimes_{\Lambda_0}\Lambda$. This is because $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$ is torsion-free by Theorem 2.32 (1). We use Proposition 2.15 (1). Then the factor $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b};\mathfrak y)$ injects to the right hand side of (5.10) since $\vec{\mathfrak v}_T(\mathfrak y) \in P_\epsilon$. Therefore (5.10) is injective.

We can prove

$$\frac{\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle}{\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1, \dots, n\right)} \cong \frac{\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{d_{P^\epsilon}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1, \dots, n\right)}$$

in the same way as Lemma 12.8 (3). This implies (5.8). The proof of (5.9) is similar. □

Remark 5.7. We used Lemma 12.8 in the proof of Proposition 5.6. We do not use Proposition 5.6 during the proof of Lemma 12.8.

6. OPERATOR $\mathfrak q$: REVIEW

We first review the construction of operator $\mathfrak q$. It is described in [FOOO3] Section 7.4 in the general case and [FOOO5] Section 6 in toric case.

Let $\mathcal M^{\mathrm{main}}_{k+1;\ell}(\beta)$ be the compactified moduli space of the genus zero bordered holomorphic maps $u : (\Sigma, \partial\Sigma) \to (X, L)$, in class $\beta \in H_2(X, L(u); \mathbb Z)$ with $k+1$ boundary marked points and ℓ interior marked points. (We require the order of $k+1$ boundary marked points respects the counter-clockwise cyclic order of the boundary of the Riemann surface.) We have the evaluation maps

$$\mathrm{ev} : \mathcal M^{\mathrm{main}}_{k+1;\ell}(\beta) \to X^\ell \times L(u)^{k+1}$$

where we put

$$\mathrm{ev} = (\mathrm{ev}^{\mathrm{int}}, \mathrm{ev}) = (\mathrm{ev}^{\mathrm{int}}_1, \dots, \mathrm{ev}^{\mathrm{int}}_\ell; \mathrm{ev}_0, \dots, \mathrm{ev}_k).$$

We denote the ℓ-tuple of T^n-invariant cycles by

$$\mathbf p = (\mathbf p(1), \dots, \mathbf p(\ell)).$$

Here $\mathbf p(i)$ is a T^n-invariant cycle in $\bigoplus_{k>0} \mathcal A^k(\mathbb Z)$.

We remark that the elements of $\mathcal A^0(\mathbb Z)$ change the potential function only by constant and does not affect the Jacobian ring.

We define the fiber product

$$\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p}) = \mathcal{M}^{\rm main}_{k+1;\ell}(\beta)_{{\rm ev}^{\rm int}} \times_{X^\ell} (\mathbf{p}(1) \times \cdots \times \mathbf{p}(\ell)). \tag{6.1}$$

It has a T^n-invariant Kuranishi structure and the evaluation map

$$\mathrm{ev}_0 : \mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p}) \to L(u) \tag{6.2}$$

is T^n-invariant. Since T^n action on $L(u)$ is free, it follows from T^n-equivariance of (6.2) that the T^n action on each of the Kuranishi neighborhood of $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})$ is free. We use this fact to take an appropriate multisection of $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})$.

To describe the properties of this multisection, we need to review some notations which we introduced in Section 6 [FOOO5]. We denote the set of shuffles of ℓ elements by

$$\mathrm{Shuff}(\ell) = \{(\mathbb{L}_1,\mathbb{L}_2) \mid \mathbb{L}_1 \cup \mathbb{L}_2 = \{1,\dots,\ell\},\ \mathbb{L}_1 \cap \mathbb{L}_2 = \emptyset\}. \tag{6.3}$$

Let $\underline{B} = \{1,\dots,B\}$ and $Map(\ell,\underline{B})$ the set of all maps $\{1,\dots,\ell\} \to \underline{B}$. We will define a map

$$\mathrm{Split} : \mathrm{Shuff}(\ell) \times Map(\ell,\underline{B}) \longrightarrow \bigcup_{\ell_1+\ell_2=\ell} Map(\ell_1,\underline{B}) \times Map(\ell_2,\underline{B}), \tag{6.4}$$

as follows: Let $\mathbf{p} \in Map(\ell,\underline{B})$ and $(\mathbb{L}_1,\mathbb{L}_2) \in \mathrm{Shuff}(\ell)$. We put $\ell_j = \#(\mathbb{L}_j)$ and let $\mathrm{i}_j : \{1,\dots,\ell_j\} \cong \mathbb{L}_j$ be the order-preserving bijection. We consider the map $\mathbf{p}_j : \{1,\dots,\ell_j\} \to \underline{B}$ defined by $\mathbf{p}_j(i) = \mathbf{p}(\mathrm{i}_j(i))$, and set

$$\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p}) = (\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_1, \mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_2) := (\mathbf{p}_1,\mathbf{p}_2).$$

We now define a corresponding gluing map

$$\begin{aligned}\mathrm{Glue}&^{(\mathbb{L}_1,\mathbb{L}_2),\mathbf{p}}_{\ell_1,\ell_2;k_1,k_2;i;\beta_1,\beta_2} \\ &: \mathcal{M}^{\rm main}_{k_1+1;\ell_1}(L(u),\beta_1;\mathbf{p}_1)_{\mathrm{ev}_0} \times_{\mathrm{ev}_i} \mathcal{M}^{\rm main}_{k_2+1;\ell_2}(L(u),\beta_2;\mathbf{p}_2) \\ &\qquad\qquad \to \mathcal{M}^{\rm main}_{k+1;\ell}(L(u),\beta;\mathbf{p})\end{aligned} \tag{6.5}$$

below. Here $k = k_1 + k_2 - 1$, $\ell = \ell_1 + \ell_2$, $\beta = \beta_1 + \beta_2$, and $i = 1,\dots,k_2$. Put

$$\mathbb{S}_j = ((\Sigma_{(j)},\varphi_{(j)},\{z^+_{i,(j)}\},\{z_{i,(j)}\}) \in \mathcal{M}^{\rm main}_{k_j+1;\ell_j}(L(u),\beta_j;\mathbf{p}_j)$$

$j = 1,2$.

Remark 6.1. Here and hereafter, boundary marked points are denoted by z_i and interior marked points are denoted either by z^+_i or $z^{\rm int}_i$.

We glue $z_{0,(1)} \in \partial\Sigma_1$ with $z_{i,(2)} \in \partial\Sigma_2$ to obtain

$$\Sigma = \Sigma_1 \#_i \Sigma_2.$$

Suppose $(\mathbb{S}_1,\mathbb{S}_2)$ is an element of the fiber product in the left hand side of (6.5). Namely we assume

$$\varphi_{(1)}(z_{0,(1)}) = \varphi_{(2)}(z_{i,(2)}).$$

This defines a holomorphic map

$$\varphi = \varphi_{(1)} \#_i \varphi_{(2)} : \Sigma \to X$$

by putting $\varphi = \varphi_{(j)}$ on Σ_j.

Let $m \in \mathbb{L}_j$. Then $\mathfrak{i}_j(c) = m$, $\mathfrak{i}_j : \{1, \dots, \ell_j\} \cong \mathbb{L}_j$ be the order-preserving bijection. We define the m-th interior marked point $z_m^{\rm int}$ of φ as $z_{c;(j)}^{\rm int} \in \Sigma_j \subset \Sigma$. We define the boundary marked points $(z_0, z_1, \dots, z_k)$ on Σ by

$$(z_0, z_1, \dots, z_k) = (z_{0,(2)}, \dots, z_{i-1,(2)}, z_{1,(1)}, \dots, z_{k_1,(1)}, z_{i+1,(2)}, \dots, z_{k_2,(2)}).$$

Now we put

$$\mathbb{S} = ((\Sigma, \varphi, \{z_i^+\}, \{z_i\})$$

and

$$\mathrm{Glue}_{\ell_1,\ell_2;k_1,k_2;i;\beta_1,\beta_2}^{(\mathbb{L}_1,\mathbb{L}_2),\mathbf{p}}(\mathbb{S}_1, \mathbb{S}_2) = \mathbb{S}.$$

The following lemma is proved in [FOOO3] Section 7.1.

Lemma 6.2. *We have the following identity as the spaces with Kuranishi structure:*

$$\partial\mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p}) = \bigcup_{\substack{k_1+k_2=k+1\\ \beta_1+\beta_2=\beta}} \bigcup_{i=1}^{k_2} \bigcup_{(\mathbb{L}_1,\mathbb{L}_2)\in \mathrm{Shuff}(\ell)} \mathcal{M}_{k_1+1;|\mathbb{L}_1|}^{\rm main}(\beta_1; (\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_1)$$
$$ {}_{\mathrm{ev}_0}\times_{\mathrm{ev}_i} \mathcal{M}_{k_2+1;|\mathbb{L}_2|}^{\rm main}(\beta_2; (\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_2).$$

Let $\sigma : \{1, \dots, \ell\} \to \{1, \dots, \ell\}$ be an element of $\mathfrak{S}_\ell$, the symmetric group of order $\ell!$. We have the induced map

$$\sigma_* : \mathcal{M}_{k+1;\ell}^{\rm main}(\beta) \to \mathcal{M}_{k+1;\ell}^{\rm main}(\beta)$$

which permutates the interior marked points. We define a new ℓ-tuple $\mathbf{p}_\sigma$ by setting $\mathbf{p}_\sigma(i) = \mathbf{p}(\sigma(i))$. Then σ_* induces a map

$$\sigma_* : \mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p}) \to \mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p}_\sigma). \tag{6.6}$$

We now consider the following condition for a system of multisections $\mathfrak{q}$. (Here we use the same letter $\mathfrak{q}$ as the operator $\mathfrak{q}$ to indicate that this multisection is used to define the operator $\mathfrak{q}$.) We write $\mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}}$ for the zero set of this multisection.

Condition 6.3.
(1) The multisection $\mathfrak{q}$ is T^n-invariant and transversal to 0.
(2) The evaluation map $\mathrm{ev}_0 : \mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}} \to L(u)$ is a submersion. Here the left hand side is the zero set of our multisection $\mathfrak{q}$.)
(3) It is compatible to other multisections given at the boundary

$$\mathcal{M}_{k_1+1;\ell_1}^{\rm main}(\beta_1;\mathbf{p}_1)^{\mathfrak{q}}{}_{\mathrm{ev}_0}\times_{\mathrm{ev}_i} \mathcal{M}_{k_2+1;\ell_2}^{\rm main}(\beta_2;\mathbf{p}_2)^{\mathfrak{q}}$$

where $k_1 + k_2 = k + 1$, $\beta_1 + \beta_2 = \beta$, $i \in \{1, \dots, k_2\}$ and

$$\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p}) = (\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_1, \mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})_2) := (\mathbf{p}_1,\mathbf{p}_2).$$

(4) It is compatible with the forgetful map

$$\mathfrak{forget} : \mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}} \to \mathcal{M}_{1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}}$$

which forgets the 1-st ... k-th boundary marked points.
(5) It is invariant under the action of the symmetric group which exchanges the interior marked points and the factors of $\mathbf{p}$.

Remark 6.4. Condition 6.3 (1) implies that $\mathrm{ev}_i : \mathcal{M}_{k+1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}} \to L(u)$ is a submersion for each $i = 0, \dots, k+1$. (So (2) is a consequence of (1) actually.) It is in general impossible to take our perturbation $\mathfrak{q}$ so that

$$(\mathrm{ev}_i, \mathrm{ev}_j) : M_{k+1;\ell}^{\rm main}(\beta;\mathbf{p})^{\mathfrak{q}} \to L(u)^2$$

is a submersion as far as we consider a *single* multisection.

The following is proved in [FOOO5].

Lemma 6.5 ([FOOO5] Lemma 6.5. See also Sections 29 - 30 of this paper.)**.** *There exists a system of multisections, which satisfies Condition 6.3 above.*

Definition 6.6. Hereafter we call the multisection on $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta;\mathbf{p})$ chosen in Lemma 6.5 [FOOO5] the *'$\mathfrak{q}$'-multisection.* This is the multisection we use to define the operator $\mathfrak{q}$.

Let C be a graded module. We define its degree shift $C[1]$ by $C[1]^k = C^{k+1}$. We put

$$B_kC = \underbrace{C\otimes\cdots\otimes C}_{k \text{ times}}. \tag{6.7}$$

Let

$$\widehat{B}C = \widehat{\bigoplus_{k=0}^{\infty}} B_kC$$

be the completed direct sum of them. We also consider a map $\Delta^{k-1} : BC \to (BC)^{\otimes k}$

$$\Delta^{k-1} = (\Delta\otimes\underbrace{id\otimes\cdots\otimes id}_{k-2})\circ(\Delta\otimes\underbrace{id\otimes\cdots\otimes id}_{k-3})\circ\cdots\circ\Delta.$$

For an indecomposable element $\mathbf{x}\in BC$, it can be expressed as

$$\Delta^{k-1}(\mathbf{x}) = \sum_c \mathbf{x}_c^{k;1}\otimes\cdots\otimes\mathbf{x}_c^{k;k}$$

where c runs over some index set.

Let $\mathfrak{S}_k$ be the symmetric group of order $k!$. It acts on B_kC by

$$\sigma\cdot(x_1\otimes\cdots\otimes x_k) = (-1)^* x_{\sigma(1)}\otimes\cdots\otimes x_{\sigma(k)} \tag{6.8}$$

where $* = \sum_{i<j:\sigma(i)>\sigma(j)} \deg x_i \deg x_j$.

Definition 6.7. E_kC is the quotient of B_kC by the submodule generated by $\sigma\cdot\mathbf{x}-\mathbf{x}$, where $\sigma\in\mathfrak{S}_k$, $\mathbf{x}\in B_kC$.

$\widehat{E}C$ is a completion of the direct product $\bigoplus_{k=0}^{\infty} E_kC$.

For an element $\mathbf{x}\in B_kC$ we denote its equivalence class in E_kC by $[\mathbf{x}]$.

Remark 6.8. In [FOOO3, FOOO5] the authors defined E_kC as the *subset* of elements of B_kC which are *invariant* under the $\mathfrak{S}_k$ action. Here we take the *quotient* instead. This difference is not essential since we are working on a ring containing $\mathbb{Q}$. It is slightly more convenient to use the quotient for the description of this paper.

There are a few changes which occur accordingly. Especially for the element $\mathfrak{b}$ of C we define

$$e^{\mathfrak{b}} = \sum \frac{1}{k!}\mathfrak{b}^k \in \widehat{E}C.$$

Note in case of $\widehat{B}C$, $e^{\mathfrak{b}} = \sum \mathfrak{b}^k \in \widehat{B}C[1]$. In [FOOO3, FOOO5] we put $e^{\mathfrak{b}} = \sum \mathfrak{b}^k$ also in case of $\widehat{E}C[1]$. In fact the coalgebra structure on $\widehat{E}C[1]$ is given by

$$\begin{aligned}&\Delta([x_1\otimes\cdots\otimes x_k])\\ &= \sum_{(\mathbb{L}_1,\mathbb{L}_2)\in\text{Shuffle}(k)} (-1)^*[x_{\ell_1(1)}\otimes\cdots\otimes x_{\ell_1(\ell_1(k_1))}]\otimes[x_{\ell_2(1)}\otimes\cdots\otimes x_{\ell_2(\ell_2(k_2))}]\end{aligned}$$

where $\mathbb{L}_1 = \{\ell_1(1), \dots, \ell_1(k_1)\}$, $\mathbb{L}_2 = \{\ell_2(1), \dots, \ell_2(k_2)\}$ and

$$* = \sum_{\ell_2(i)<\ell_1(j)} \deg' x_{\ell_2(i)} \deg' x_{\ell_1(j)}.$$

Therefore

$$\Delta\left(\sum_k \frac{1}{k!}\mathfrak{b}^k\right) = \left(\sum_k \frac{1}{k!}\mathfrak{b}^k\right) \otimes \left(\sum_k \frac{1}{k!}\mathfrak{b}^k\right). \tag{6.9}$$

We fix a T^n-invariant Riemannian metric on $L(u)$. We identify $H(L;\mathbb{R})$ with the vector space of harmonic forms. The following lemma can be easily checked and so its proof is omitted. This characterization of harmonic forms for the torus turns out to be useful for later discussions.

Lemma 6.9. *Equip T^n with an invariant Riemannian metric. Then a differential form is harmonic with respect to this metric if and only if it is T^n-invariant.*

We also identify $H(L(u);\Lambda_0(\mathbb{R}))$ with $H(L(u);\mathbb{R}) \otimes_{\mathbb{R}} \Lambda_0(\mathbb{R})$. Hereafter, we write Λ_0 etc. in place of $\Lambda_0(\mathbb{R})$ etc..

We take $h_1, \dots, h_k \in H(L(u);\Lambda_0)$ and identify them with harmonic forms, that is T^n-invariant forms in our case.

Definition 6.10. We define

$$\mathfrak{q}_{\ell;k;\beta}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k) = \mathrm{ev}_{0*}\mathrm{ev}^*(h_1 \times \cdots \times h_k), \tag{6.10}$$

where

$$(\mathrm{ev}_0, \mathrm{ev}) : \mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}} \to L(u) \times L(u)^k.$$

Condition 6.3 (5) implies that the left hand side of (6.10) is independent of the representative $\mathbf{p}$ but depends only on $[\mathbf{p}] \in \widehat{E}(\mathcal{A}[2])$. We thus obtain:

$$\mathfrak{q}_{\ell;k;\beta} : E_\ell(\mathcal{A}[2]) \otimes B_{k+1}(H(L(u);\mathbb{Q})[1]) \to H(L(u);\mathbb{Q})[1].$$

Remark 6.11.
1. In our situation, all the forms h_i and the (perturbed) moduli spaces are T^n-invariant. Therefore $\mathfrak{q}_{\ell;k;\beta}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k)$ is also T^n-invariant. So we need not take the procedure to go to the canonical model by using the Green kernel to define $\mathfrak{q}$.
2. In Section 3.8 [FOOO3] (Definition 3.8.67 (3.8.68)) there is a factor $1/\ell!$ is in the right hand side. There is the same factor in [FOOO5] (6.10). We do not put it here, since we take quotient to define EC here. (See Remark 6.8.)

Convention 6.12. Hereafter we write

$$\mathrm{ev}_*(\mathrm{ev}^*(h_1 \times \cdots \times h_k); \mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}})$$

for the right hand side of (6.10). We will apply the similar notation in the general context. (See Section 28 for the usage of the same notation in the general context.)

We take $u \in \mathrm{Int}\, P$. Let $b_0 = \sum x_i \mathbf{e}_i$ $x_i \in \mathbb{C}$ and associate a representation $\rho : H_1(L(u);\mathbb{Z}) \to \mathbb{C}^*$ by $\rho(e_i) = e^{x_i}$.

Definition 6.13. Let $h_1, \dots, h_k \in H(L(u);\Lambda_0)$ be T^n-invariant forms. We put:

$$\begin{aligned}&\mathfrak{q}^\rho_{\ell;k}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k) \\ &= \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} T^{\beta\cap\omega/2\pi}\rho(\partial\beta)\mathfrak{q}_{\ell;k;\beta}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k).\end{aligned}$$

The main property of the operator $\mathfrak{q}$ above is Theorem 3.8.32 [FOOO3] or Theorem 2.1 [FOOO5]. Namely we have:

(1) For each β and $\mathbf{x} \in B_k(H(L;\mathbb{R})[1])$, $\mathbf{y} \in E_k(H[2])$, we have the following:
$$0 = \sum_{\beta_1+\beta_2=\beta} \sum_{c_1,c_2} (-1)^* \mathfrak{q}_{\beta_1}(\mathbf{y}^{2;1}_{c_1}; \mathbf{x}^{3;1}_{c_2} \otimes \mathfrak{q}_{\beta_2}(\mathbf{y}^{2;2}_{c_1}; \mathbf{x}^{3;2}_{c_2}) \otimes \mathbf{x}^{3;3}_{c_2}) \tag{6.11a}$$
where
$$* = \deg' \mathbf{x}^{3;1}_{c_2} + \deg' \mathbf{x}^{3;1}_{c_2} \deg \mathbf{y}^{2;2}_{c_1} + \deg \mathbf{y}^{2;1}_{c_1}.$$
In (6.11a) and hereafter, we write $\mathfrak{q}_\beta(\mathbf{y};\mathbf{x})$ in place of $\mathfrak{q}_{\beta;\ell,k}(\mathbf{y};\mathbf{x})$ if $\mathbf{y} \in E_\ell(H[2])$, $\mathbf{x} \in B_k(H(L;\mathbb{R})[1])$.

(2) Let $\mathbf{e} = PD([L])$ be the Poincaré dual to the fundamental class of L. Let $\mathbf{x}_i \in B(H(L;\mathbb{R})[1])$ and we put $\mathbf{x} = \mathbf{x}_1 \otimes \mathbf{e} \otimes \mathbf{x}_2 \in B(H(L;R)[1])$. Then
$$\mathfrak{q}_\beta(\mathbf{y};\mathbf{x}) = 0 \tag{6.11b}$$
except the following case:
$$\mathfrak{q}_{\beta_0}(1; \mathbf{e} \otimes x) = (-1)^{\deg x} \mathfrak{q}_{\beta_0}(1; x \otimes \mathbf{e}) = x, \tag{6.11c}$$
where $\beta_0 = 0 \in H_2(X,L;\mathbb{Z})$ and $x \in H(L;\mathbb{R})[1] = B_1(H(L;\mathbb{R})[1])$.

We can prove them by using (1) $-$ (5) Condition 6.3.

Definition 6.14. Let $\mathfrak{b} \in H^{even}(X;\Lambda_0)$ and $\rho : H_1(L(u);\mathbb{Z}) \to \mathbb{C}^*$. We write $\mathfrak{b} = \mathfrak{b}_0 + \mathfrak{b}_2 + \mathfrak{b}_{\text{high}}$ where $\mathfrak{b}_0 \in H^0(X;\Lambda_0)$, $\mathfrak{b}_2 \in H^2(X;\Lambda_0)$, $\mathfrak{b}_{\text{high}} \in H^{2m}(X;\Lambda_0)$ $(m > 1)$. We define $\mathfrak{m}^\rho_{k;\beta} : B_k(H(L;\mathbb{R})[1]) \to H(L;\mathbb{R})[1]$ by
$$\begin{aligned} &\mathfrak{m}^{\mathfrak{b},\rho}_k(h_1,\dots,h_k) \\ &= \sum_\beta \exp(\mathfrak{b}_2 \cap \beta) \sum_{\ell=0}^\infty T^{\beta\cap\omega/2\pi} \rho(\partial\beta) \mathfrak{q}_{\ell;k;\beta}(\mathfrak{b}^\ell_{\text{high}}; h_1,\dots,h_k). \end{aligned} \tag{6.12}$$

Let $b \in H^1(L(u);\Lambda_0)$. We define $b = b_0 + b_+$ where $b_0 \in H^1(L(u);\mathbb{C})$ and $b_+ \in H^1(L(u);\Lambda_+)$. We define $\rho : H_1(L(u);\mathbb{Z}) \to \mathbb{C}^*$ by $\gamma \mapsto e^{\gamma\cap b_0}$. We define
$$\mathfrak{m}^{\mathfrak{b},b}_k(h_1,\dots,h_k) = \sum_{l_0,\dots,l_k} \mathfrak{m}^\rho_{k+\sum l_i;\beta}(b_+^{l_0}, h_1, b_+^1, \dots, b_+^{l_{k-1}}, h_k, b_+^{l_k}). \tag{6.13}$$

We can use (6.11) to prove that $\mathfrak{m}^{\mathfrak{b},b}_k$ defines a unital filtered A_∞ structure.

In [FOOO5] Definition 8.2, we defined a chain complex $(\Omega(L(u)) \widehat{\otimes} \Lambda_0; \delta_1^{(\mathfrak{b},b)})$ on the de Rham complex of Lagrangian fiber $L(u)$ (tensored with the universal Novikov ring). We can descend the (strict and gapped) filtered A_∞ algebra on $\Omega(L(u)) \widehat{\otimes} \Lambda_0$ to the de Rham cohomology $H(L(u);\Lambda_0)$. Especially we can define the boundary operator $\mathfrak{m}^{\mathfrak{b},b}_1 = \pm\delta^{\mathfrak{b},b}_{\text{can}}$ on $H(L(u);\Lambda_0)$ as above[2]. This is a general fact proven in Section 5.4 [FOOO3]. In the rest of this section we will review its construction and define the chain homotopy equivalence:
$$\Pi : (\Omega(L(u)) \widehat{\otimes} \Lambda_0 \,, \delta^{\mathfrak{b},b}) \to (H(L(u);\Lambda_0), \delta^{\mathfrak{b},b}_{\text{can}}). \tag{6.14}$$

We first recall the definition of $\delta^{\mathfrak{b},b}$ in Section 8 [FOOO5].

---

[2] Actually it is zero in case Floer cohomology is isomorphic to the classical cohomology, that is the case b is a critical point of $\mathfrak{PO}^u_{\mathfrak{b}}$.

We use the correspondence by this perturbed moduli space to define operators on the de Rham complex $\Omega(L(u))$ of $L(u)$. Let $h \in \Omega^d(L(u))$ be a degree d smooth differential form on $L(u)$. Let

$$\mathrm{ev}_1^*(h) \in \Omega^{\deg h}(\mathcal{M}_{2,\ell}(\beta;\mathfrak{b}^{\otimes \ell})) \tag{6.15}$$

be the pull back of h. We define

$$\delta^{\mathfrak{b}}_{\beta;\ell}(h) = (-1)^{\deg h}(\mathrm{ev}_0)_!\mathrm{ev}_1^*(h) = \mathfrak{m}^{\mathfrak{b}}_{1,\beta;\ell}(h) \in \Omega^*(L(u)) \tag{6.16}$$

where $(\mathrm{ev}_0)_!$ is the integration along fiber. We remark that T^n equivariance of ev_0 and transitivity of the T^n action on $L(u)$ imply that ev_0 is a submersion. Therefore integration along the fiber is well-defined and defines a smooth form on L. The degree $*$ is given by

$$* = \deg h - \mu(\beta) + 1$$

where μ is the Maslov index.

Remark 6.15. (1) We remark that $\mathfrak{b}$ is not in $\mathcal{A}(\mathbb{Z})$ but in $\mathcal{A}(\Lambda_0)$. So, more precisely we need to define $\delta^{\mathfrak{b}}_{\beta;\ell}$ as follows. We put

$$\mathfrak{b} = \sum_{i=1}^{B} w_i(\mathfrak{b})\mathbf{f}_i.$$

For each multi-index $I = (i_1,\dots,i_\ell)$, we define $\mathbf{f}_I = (\mathbf{f}_{i_1},\dots,\mathbf{f}_{i_\ell})$ and consider

$$\mathrm{ev}^I = (\mathrm{ev}^I_0, \mathrm{ev}^I_1) : \mathcal{M}^{\mathrm{main}}_{2;\ell}(\beta;\mathbf{f}_I) \to L(u)\times L(u).$$

Now we put

$$\delta^{\mathfrak{b}}_{\beta;\ell}(h) = (-1)^{\deg h}\sum_{I\in\{1,\dots,B\}^\ell} w_{i_1}(\mathfrak{b})\cdots w_{i_\ell}(\mathfrak{b})(\mathrm{ev}^I_0)_!(\mathrm{ev}^I_1)^*(h). \tag{6.17}$$

Hereafter we write (6.16) in place of (6.17) whenever no confusion can occur.

(2) We remark that when we need to use a multisection rather than a single valued section, the perturbed moduli space $\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{f}_I)^{\mathfrak{s}}$ is neither a manifold nor an orbifold. However integration along the fiber and pull-back of differential forms on such space can be defined in the way similar to the case of manifold. (See Section 12 [FOOO5] for the details.)

We next use $\delta^{\mathfrak{b}}_{\beta;\ell}$ to define $\delta^{\mathfrak{b},b}$. Let

$$b = \sum_{i=1}^{n}\mathfrak{x}_i\mathbf{e}_i \in H^1(L(u);\Lambda_0).$$

We put $\mathfrak{y}_i = e^{\mathfrak{x}_i} \in \Lambda_0 \setminus \Lambda_+$, and define $\rho^b : H_1(L(u);\mathbb{Z}) \to \Lambda_0\setminus\Lambda_+$ by

$$\rho^b\left(\sum k_i\mathbf{e}_i^*\right) = \mathfrak{y}_1^{k_1}\cdots\mathfrak{y}_n^{k_n}. \tag{6.18}$$

Definition 6.16. We define $\delta^{\mathfrak{b},b} : \Omega(L(u))\,\widehat{\otimes}\,\Lambda_0 \to \Omega(L(u))\,\widehat{\otimes}\,\Lambda_0$ by

$$\delta^{\mathfrak{b},b} = \mathfrak{m}_{1,0} + \sum_{\beta,\ell}\frac{1}{\ell!}T^{\beta\cap\omega/2\pi}\rho^b(\partial\beta)\delta^{\mathfrak{b}}_{\beta,\ell},$$

Here $\mathfrak{m}_{1,0}$ is defined from the de Rham differential d by $\mathfrak{m}_{1,0}(h) = (-1)^{n+\deg h+1}dh$. See [FOOO3] Remark 3.5.8.

By (6.16) we easily find that if $h \in \Omega(L(u))$ is T^n-invariant then $\delta^{\mathfrak{b},b}_{\beta,\ell}(h) = \rho^b(\partial\beta)\delta^{\mathfrak{b}}_{\beta,\ell}(h)$ is also T^n-invariant . Therefore

$$\delta^{\mathfrak{b},b}(H(L(u);\Lambda_0)) \subset H(L(u);\Lambda_0).$$

We define

$$\delta^{\mathfrak{b},b}_{\rm can} : H(L(u);\Lambda_0) \to H(L(u);\Lambda_0)$$

as the restriction of $\delta^{\mathfrak{b},b}$.

Remark 6.17. In fact, we have $\delta^{\mathfrak{b},b}_{\rm can} = \mathfrak{m}^{\mathfrak{b},b}_1$ where the right hand side is in Definition 6.14. We will prove it in Section 27.

Let

$$\Pi : \Omega(L(u)) \to H(L(u);\mathbb{R}) \hookrightarrow \Omega(L(u))$$

be the harmonic projection.

Now we have the following:

Theorem 6.18. *We have*

$$\delta^{\mathfrak{b},b} \circ \delta^{\mathfrak{b},b} = 0. \tag{6.19}$$

Moreover Π *is a chain homotopy equivalence.*

Proof. The proof is based on the study of boundary of $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})$. Using Lemma 6.2, Condition 6.3 and the definition, we can easily show the following formula.

$$\mathfrak{m}_{1,0} \circ \delta^{\mathfrak{b}}_{\beta;\ell} + \delta^{\mathfrak{b}}_{\beta;\ell} \circ \mathfrak{m}_{1,0} = -\sum_{\beta_1+\beta_2=\beta}\sum_{\ell_1,\ell_2=0}^{\infty} \frac{\ell!}{\ell_1!\ell_2!}\delta^{\mathfrak{b}}_{\beta_1;\ell_1} \circ \delta^{\mathfrak{b}}_{\beta_2;\ell_2}. \tag{6.20}$$

Therefore

$$\begin{aligned}(\delta^{\mathfrak{b},b})^2 = \mathfrak{m}_{1,0}^2 &+ \sum_{\beta}\sum_{\ell=0}^{\infty}\frac{\rho^b(\partial\beta)}{\ell!}(\delta^{\mathfrak{b}}_{\beta;\ell} \circ \mathfrak{m}_{1,0} + \mathfrak{m}_{1,0} \circ \delta^{\mathfrak{b}}_{\beta;\ell}) \\ &+ \sum_{\beta_1+\beta_2=\beta}\sum_{\ell_1,\ell_2=0}^{\infty}\frac{\rho^b(\partial\beta_1)\rho^b(\partial\beta_2)}{\ell_1!\ell_2!}\delta^{\mathfrak{b}}_{\beta_1;\ell_1} \circ \delta^{\mathfrak{b}}_{\beta_2;\ell_2} = 0.\end{aligned}$$

We next show that Π is a chain map. We remark that

$$\Pi(h) = \int_{g\in T^n} g^* h \; dg, \tag{6.21}$$

where $g : L \to L$ for $g \in T^n$ is obtained by T^n action and dg is the normalized Haar measure. The formula (6.21) and T^n invariance of $\mathcal{M}^{\rm main}_{2;\ell}(\beta;\mathbf{f}_2)$ imply that

$$\Pi \circ \delta^{\mathfrak{b},b}_{\rm can} = \delta^{\mathfrak{b},b}_{\rm can} \circ \Pi.$$

Therefore Π is a chain map. It is then easy to show that it is a chain homotopy equivalence. $\square$

7. WELL-DEFINEDNESS OF KODAIRA-SPENCER MAP

In Section 2 we used a choice of a section $H(X;\Lambda_0) \to \mathcal{A}(\Lambda_0)$ of π to give the definition of the Kodaira-Spencer map

$$\mathfrak{ks}_{\mathfrak{b}} : H(X;\Lambda_0) \to \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}).$$

In this section, we prove that the definition of $\mathfrak{ks}_{\mathfrak{b}}$ does not depend on the choice of the section.

To make this statement precise, we need to set-up some notations. Let $w_0, \dots, w_B$ be a coordinates of $\mathfrak{b}$ with respect to the basis $\{\mathfrak{f}_i\}_{i=1}^{B}$, i.e., $\mathfrak{b} = \sum_i w_i \mathfrak{f}_i$. We consider the potential function $\mathfrak{PO}(w_0, \dots, w_B; y_1, \dots, y_n)$ for $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ and define a map $\widetilde{\mathfrak{ks}}_{\mathfrak{b}} : T_{\mathfrak{b}}\mathcal{A}(\Lambda_0) \to \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ by

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}\left(\frac{\partial}{\partial w_i}\right) = \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial w_i}.$$

Let $\pi : \mathcal{A}(\Lambda_0) \to H(X;\Lambda_0)$ be the canonical map given in (2.12).

Theorem 7.1. *The following diagram commutes:*

$$\begin{tikzcd}
\mathcal{A}(\Lambda_0) \arrow[r, "\widetilde{\mathfrak{ks}}_{\mathfrak{b}}"] \arrow[d, "\pi"'] & \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle \arrow[d] \\
H(X;\Lambda_0) \arrow[r, "\mathfrak{ks}_{\mathfrak{b}}"] & \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}).
\end{tikzcd} \tag{7.1}$$

In particular, the definition of $\mathfrak{ks}_{\mathfrak{b}}$ is independent of the section $H(X;\Lambda_0) \to \mathcal{A}(\Lambda_0)$ used in Section 2.

The rest of this section is occupied with the proof of this theorem.

Let $Q = \sum_i w_i(Q)\mathbf{f}_i$ with $[Q] = 0$ in $H(X;\Lambda_0)$. We will prove

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q) \in \mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1, \dots, n\right). \tag{7.2}$$

We may assume $w_0(Q) = 0$. (In fact, $\mathcal{A}^0 \to H^0(L(u);\mathbb{Z})$ is an isomorphism.) We consider the case of β, $\mathfrak{b}$ and $\mathfrak{f}$ for which the moduli spaces

$$\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u), \beta; \mathfrak{b}^{\otimes \ell} \otimes \mathbf{f}_i) \tag{7.3}$$

have dimension n. Then the evaluation map

$$\mathrm{ev}_0 : \mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u), \beta; \mathfrak{b}^{\otimes \ell} \otimes \mathbf{f}_i) \to L(u)$$

defines a homology class

$$\mathrm{ev}_{0*}(\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u), \beta; \mathfrak{b}^{\otimes \ell} \otimes \mathbf{f}_i)) \in H_n(L(u);\Lambda_0) \cong H^0(L(u);\Lambda_0) = \Lambda_0. \tag{7.4}$$

Remark 7.2. Actually $\mathfrak{b} = \sum \mathfrak{b}_i T^{\lambda_i}$ is not a space but contains T. So (7.3) is not a space but a kind of formal power series with space as coefficients. Namely

$$\begin{aligned}
&\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u), \beta; \mathfrak{b}^{\otimes \ell} \otimes \mathbf{f}_i) \\
&= \sum_{i_1=1}^{\infty} \cdots \sum_{i_\ell=1}^{\infty} \mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u), \beta; \mathfrak{b}_{i_1} \otimes \dots \mathfrak{b}_{i_\ell} \otimes \mathbf{f}_i) T^{\lambda_{i_1} + \dots + \lambda_{i_\ell}}.
\end{aligned}$$

Then the left hand side of (7.4) has an obvious meaning. By an abuse of notation we use the expression such as (7.3) frequently for the rest of this paper.

Lemma 7.3. *Let* $b = \sum_{i=1}^n x_i \mathbf{e}_i$ *and* $y_i = e^{x_i}$. *Then we have*

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)(y) = \sum_{i=1}^{B} \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} \frac{w_i(Q)}{\ell!} T^{\omega\cap\beta/2\pi} \rho^b(\partial\beta) \operatorname{ev}_{0*}(\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u),\beta;\mathfrak{b}^{\otimes\ell}\otimes\mathbf{f}_i)),$$

where $Q = \sum w_i(Q)\mathbf{f}_i$. *Note the right hand side is an element of* Λ_0 *by the isomorphism (7.4).*

Proof. This follows by differentiating

$$\mathfrak{PO}(\mathfrak{b};y) = \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} \frac{1}{\ell!} T^{\omega\cap\beta/2\pi} \rho^b(\partial\beta) \operatorname{ev}_{0*}(\mathcal{M}^{\mathrm{main}}_{1;\ell}(L(u),\beta;\mathfrak{b}^{\otimes\ell})) \tag{7.5}$$

with respect to w_i's. (7.5) follows from (2.10). □

For the convergence of our series, we slightly rewrite Lemma 7.3. (See the proof of Proposition 11.4 [FOOO5].) We put $\mathfrak{b} = \sum_{a=1}^B \mathfrak{b}_a \mathbf{f}_a$. (We need not consider $\mathbf{f}_0$ because it does not affect the Jacobian ring.) We put

$$\mathfrak{b}_{\mathrm{high}} = \sum_{a=m+1}^{B} \mathfrak{b}_a \mathbf{f}_a, \quad \mathfrak{b}_2 = \sum_{a=1}^{m} \mathfrak{b}_a \mathbf{f}_a. \tag{7.6}$$

Namely they correspond to the degree ≥ 4 part and the degree 2 part of $\mathfrak{b}$, respectively.

Lemma 7.4. *We have*

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)(y) = \sum_{i=1}^{B} \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(U);\mathbb{Z})} \frac{w_i(Q)}{\ell!} T^{\omega\cap\beta/2\pi} \exp(\mathfrak{b}_2 \cap \beta) \rho^b(\partial\beta) \operatorname{ev}_{0*}(\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u),\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell}\otimes\mathbf{f}_i)).$$

We are now ready to give the proof of Theorem 7.1. We split the proof into two parts. One is consideration of the case $Q \in \bigoplus_{k\geq 4} \mathcal{A}^k(\Lambda_0)$ and the other is of the case $Q \in \mathcal{A}^2(\Lambda_0)$.

We start with the first case. We use our assumption and take a singular chain R of X such that $\partial R = Q$. (Here we fix a triangulation of Q and regard it as a smooth singular chain.) We emphasize it is *impossible* to choose such an R that is T^n-invariant.

We now consider

$$\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(L(u),\beta;\mathbf{p}\otimes R) = \mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(L(u),\beta)_{\mathrm{ev}^{\mathrm{int}}} \times (\mathbf{p}(1)\times\cdots\times\mathbf{p}(\ell)\times R).$$

Here $\mathbf{p}(1),\ldots,\mathbf{p}(\ell)$ are T^n-invariant cycles. We remark that in [FOOO5] we never considered chains on X which are not T^n-invariant. Here we however need to use the chains that are not T^n-invariant. Because of this, there are several points we need to be careful in the subsequent discussions henceforth.

Lemma 7.5. *There exists a Kuranishi structure with boundary on* $\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(L(u),\beta;\mathbf{p}\otimes R)$. *Its boundary is a disjoint union of the following three types of fiber product. Here* $\beta_1+\beta_2=\beta$, $k_1+k_2=k+1$, $i=1,\ldots,k_2$ *and* $(\mathbf{p}_1,\mathbf{p}_2) = \mathrm{Split}((L_1,L_2),\mathbf{p})$ *for some* $(L_1,L_2)\in\mathrm{Shuff}(\ell)$.

$$\mathcal{M}^{\mathrm{main}}_{k_1+1;|\mathbf{p}_1|+1}(L(u),\beta_1;\mathbf{p}_1\otimes R)_{\mathrm{ev}_0} \times_{\mathrm{ev}_i} \mathcal{M}^{\mathrm{main}}_{k_2+1;|\mathbf{p}_2|}(L(u),\beta_2;\mathbf{p}_2), \tag{7.7}$$

$$\mathcal{M}^{\text{main}}_{k_1+1;|\mathbf{p}_1|}(L(u),\beta_1;\mathbf{p}_1)_{\text{ev}_0}\times_{\text{ev}_i}\mathcal{M}^{\text{main}}_{k_2+1;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R), \tag{7.8}$$

$$\mathcal{M}^{\text{main}}_{k+1}(L(u),\beta;\mathbf{p}\otimes Q). \tag{7.9}$$

Here and hereafter we write $|\mathbf{p}|=\ell$ *if* $\mathbf{p}\in E_\ell\mathcal{A}(\Lambda_0)$.

This follows from Lemmata 6.4 and 6.5 [FOOO5].

We next define a multisection (perturbation). We remark that the multisection $\mathfrak{q}$ on $\mathcal{M}^{\text{main}}_{k+1;\ell}(L(u),\beta;\mathbf{p})$ was already fixed in Lemma 6.5 (= Lemma 6.5 [FOOO5]), which we use here. The multisection we have chosen there is T^n-invariant which we use to prove that $\text{ev}_0 : \mathcal{M}^{\text{main}}_{k+1;\ell}(L(u),\beta;\mathbf{p})^{\mathfrak{q}}\to L(u)$ is a submersion. (Note $\mathcal{M}^{\text{main}}_{k+1;\ell}(L(u),\beta;\mathbf{p})^{\mathfrak{q}}$ is a zero set of *multi-valued* perturbation. Although the zero set does not define a smooth manifold, we can define the notion of submersion from such a space to a manifold in an obvious way. In fact the notion of the tangent space at each point is well defined for such a space. (See [FO], [FOOO2] for the details.)

For the case of our moduli space $\mathcal{M}^{\text{main}}_{k+1;\ell}(L(u),\beta;\mathbf{p}\otimes R)$, we do not have T^n-action on it. So we can no longer use this technique. Therefore the evaluation map $\text{ev}_0 : \mathcal{M}^{\text{main}}_{k+1;\ell}(L(u),\beta;\mathbf{p}\otimes R)\to L(u)$ is not necessarily submersive. (It may occur for example the case where its dimension is strictly smaller than n.) On the other hand, we are using de Rham cohomology here. So we need to define the integration along the fiber of our evaluation map and we need our evaluation map to be submersive. We can do this by using the notion of continuous family of multisections. This notion was explained in detail in Section 12 [FOOO5].

Lemma 7.6. *There exists a continuous family of multisections on* $\mathcal{M}^{\text{main}}_{k+1}(L(u),\beta;\mathbf{p}\otimes R)$ *with the following properties:*

1. *It is compatible with the decomposition into* (7.7), (7.8), (7.9) *of its boundary.*
2. *The restriction of the evaluation map* ev_0 *to the zero set of the family of multisections is a submersion to* $L(u)$.
3. *It is compatible with the forgetful map*

   $$\mathfrak{forget} : \mathcal{M}^{\text{main}}_{k+1;\ell+1}(L(u),\beta;\mathbf{p}\otimes R)\to\mathcal{M}^{\text{main}}_{1;\ell+1}(L(u),\beta;\mathbf{p}\otimes R)$$

   of boundary marked points.

Proof. The proof is by an induction over the symplectic area $\beta\cap\omega$ and the number of boundary marked points, using Lemma 12.19 [FOOO5]. □

Lemma 7.7. *We have an equality*

$$\text{ev}_{0*}\left(\mathcal{M}^{\text{main}}_{1;|\mathbf{p}_1|}(L(u),\beta_1;\mathbf{p}_1)_{\text{ev}_0}\times_{\text{ev}_1}\mathcal{M}^{\text{main}}_{2;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R)\right)=0 \tag{7.10}$$

as differential forms on $L(u)$.

Proof. By definition, it is easy to see that the differential form is the integration along the fiber of the differential form

$$\text{ev}_{0*}\left(\mathcal{M}^{\text{main}}_{1;|\mathbf{p}_1|}(L(u),\beta_1;\mathbf{p}_1)\right)$$

by the correspondence

$$(\text{ev}_0,\text{ev}_1):\mathcal{M}^{\text{main}}_{2;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R)\to L\times L.$$

On the other hand, it follows from Corollary 6.6 [FOOO5] that the dimension of $\mathcal{M}^{\text{main}}_{1;|\mathbf{p}_1|}(L(u),\beta_1;\mathbf{p}_1)$ is n or higher. Therefore integration along the fiber gives rise to a function, that is a degree 0 form on $L(u)$. (See (7.4). Actually it is a constant because of T^n invariance.)

By Lemma 7.6 (3), the evaluation map ev_1 factors through

$$\mathcal{M}^{\text{main}}_{2;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R) \overset{\mathfrak{forget}}{\longrightarrow} \mathcal{M}^{\text{main}}_{1;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R) \xrightarrow{\text{ev}_0} L(u)$$

and hence the fiber of $\text{ev}_1 : \mathcal{M}^{\text{main}}_{2;|\mathbf{p}_2|+1}(L(u),\beta_2;\mathbf{p}_2\otimes R) \to L$ has positive dimension and hence (7.10) follows. $\square$

Remark 7.8. We may rephrase the argument used in the above proof as follows: Since b is a weak bounding cochain, $\mathfrak{m}^{\mathfrak{b},b}_{0,\beta}(1) = cPD[L]$ $(c \in \Lambda_0)$ by definition. On the other hand $PD[L]$ is a strict unit of our filtered A_∞ algebra in the de Rham model. Therefore $\mathfrak{m}^{\mathfrak{b},b}_{1,\beta}(PD[L]) = 0$.

We remark that if we had used singular cohomology instead of de Rham cohomology, then $PD[L]$ would become only a homotopy unit rather than a strict unit which would require more technical and complicated arguments. This is one of reasons why we use the de Rham cohomology model here.

Definition 7.9. We denote

$$\begin{aligned}\tilde{\mathfrak{X}}^u(\mathfrak{b},b) = \sum_{\ell=0}^{\infty}\sum_{\beta\in H_2(X;L(u);\mathbb{Z})} & \frac{1}{\ell!}T^{\omega\cap\beta/2\pi}\rho^b(\partial\beta)\exp(\mathfrak{b}_2\cap\beta) \\ & \text{ev}_{0*}(\mathcal{M}^{\text{main}}_{1;\ell+1}(L(u),\beta;\mathfrak{b}^{\otimes\ell}_{\text{high}}\otimes R)).\end{aligned} \tag{7.11}$$

This is an element of $\Omega(L(u))\widehat{\otimes}\Lambda_0$ that is formally obtained by replacing Q by R in the formula in Lemma 7.4. We also define

$$\mathfrak{X}^u(\mathfrak{b},b) = \Pi(\tilde{\mathfrak{X}}^u(\mathfrak{b},b)) \in H(L(u);\Lambda_0).$$

We note that here we fix u. We fix $\mathfrak{b}$ and u and regard $\tilde{\mathfrak{X}}^u(\mathfrak{b},b)$, $\mathfrak{X}^u(\mathfrak{b},b)$ as functions of b. Let $2^{\{1,\dots,n\}}$ be the set of all subsets of $\{1,\dots,n\}$. For $I = \{i_1,\dots,i_k\} \in 2^{\{1,\dots,n\}}$, $i_j < i_{j+1}$, we put $\mathbf{e}_I = \mathbf{e}_{i_1}\wedge\cdots\wedge\mathbf{e}_{i_k}$. It forms a basis of $H(L(u);\mathbb{Z})$. We put $b = \sum_{I\in 2^{\{1,\dots,n\}}} x_I(u)\mathbf{e}_I$, $y_i(u) = e^{x_i(u)}$ and define

$$\tilde{\mathfrak{X}}^u_{\mathfrak{b}}(y_1(u),\dots,y_n(u)) = \tilde{\mathfrak{X}}^u(\mathfrak{b},b), \quad \mathfrak{X}^u_{\mathfrak{b}}(y_1(u),\dots,y_n(u)) = \mathfrak{X}^u(\mathfrak{b},b).$$

We fix a diffeomorphism $\psi_u : T^n \cong L(u)$ which induces an identification $H^*(T^n) \cong H(L(u))$ and $\Omega(T^n) \cong \Omega(L(u))$. We also recall that y_i, $y(u)_i$ are the coordinates of $H^1(T^n;\mathbb{C})$, $H^1(L(u);\mathbb{C})$ respectively and satisfies

$$y_i(u) = T^{-u_i}y_i.$$

(See (3.10) [FOOO5] for the precise meaning of this relation.)

We regard each of $\tilde{\mathfrak{X}}^u_{\mathfrak{b}}$, $\mathfrak{X}^u_{\mathfrak{b}}$ as a formal sum of the form

$$\sum a_i T^{\lambda_i} y^{v_i}$$

where $a_i \in \Omega(T^n)$ or $a_i \in H(T^n;\mathbb{C})$ respectively. The next proposition claims that they converge in $d_{\overset{\circ}{P}}$ topology.

Proposition 7.10. (1) *We have*

$$\tilde{\mathfrak{X}}^u_{\mathfrak{b}} \in \Omega(T^n)\widehat{\otimes}\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$$

and

$$\mathfrak{X}^u_{\mathfrak{b}} \in H(T^n; \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle).$$

(2) *In case* $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$ *we can take* $\Lambda_0^P\langle\!\langle y, y^{-1}\rangle\!\rangle$ *in place of* $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$.

Proof. We remark that Gromov compactness immediately implies that the series (7.11) converges in $\mathfrak{v}_T^u$-norm. It implies

$$\tilde{\mathfrak{X}}^u_{\mathfrak{b}} \in \Lambda_0\langle\!\langle y(u), y(u)^{-1}\rangle\!\rangle.$$

To prove Proposition 7.10 it suffices to show that (7.11) converges in $\mathfrak{v}_T^{u'}$-norm for *any* $u' \in \operatorname{Int} P$. We remark that $\tilde{\mathfrak{X}}^u_{\mathfrak{b}}$ *depends* on u. (In a similar situation in Section 7 [FOOO5] we proved that the potential function (with bulk) is independent of u. In that situation, our moduli space is T^n-invariant. However in the current situation, the moduli space $\mathcal{M}^{\rm main}_{1;\ell+1}(L(u), \beta; \mathfrak{b}^{\otimes \ell} \otimes R)$ is *not* T^n-invariant. This is because the chain R is not T^n-invariant. Nevertheless, we will still be able to prove Proposition 7.10.)

Suppose $\mathrm{ev}_{0*}(\mathcal{M}^{\rm main}_{1;\ell+1}(L(u), \beta; \mathfrak{b}_{\rm high}^{\otimes \ell} \otimes R)) \neq 0$. Then by Theorem 10.1 [FOOO4] we can write

$$\beta = \sum_{i=1}^m k_i(\beta)\beta_i + \alpha(\beta) \tag{7.12}$$

so that $k_i(\beta) \geq 0$, $\sum k_i(\beta) > 0$, and that $\alpha(\beta)$ is realized by a sum of holomorphic spheres.

Since $\partial\beta = \sum_i k_i(\beta)\partial\beta_i$, it follows from (2.7) that

$$y(u)_1^{\partial\beta\cap\mathbf{e}_1}\cdots y(u)_n^{\partial\beta\cap\mathbf{e}_n} = T^{c(\beta)}z_1^{k_1(\beta)}\cdots z_m^{k_m(\beta)}$$

for

$$c(\beta) = -\sum_{i=1}^m k_i(\beta)\ell_i(u). \tag{7.13}$$

Combining (7.12) and (7.13), we obtain

$$\beta \cap \omega/2\pi = \alpha(\beta)\cap\omega/2\pi - c(\beta). \tag{7.14}$$

We write the contribution from (β, ℓ) in (7.11) as:

$$\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell} = \frac{1}{\ell!}T^{\alpha(\beta)\cap\omega/2\pi}z_1^{k_1(\beta)}\cdots z_m^{k_m(\beta)}a(\beta,\ell) \tag{7.15}$$

where

$$a(\beta,\ell) = \exp(\mathfrak{b}_0\cap\beta)\mathrm{ev}_{0*}(\mathcal{M}^{\rm main}_{1;\ell+1}(L(u), \beta; \mathfrak{b}_{\rm high}^{\otimes \ell} \otimes R)) \in \Lambda_0 \,\widehat{\otimes}\, \Omega(T^n). \tag{7.16}$$

Lemma 7.11. *Suppose that there are infinitely many different* (β_c, ℓ_c)*'s* ($c = 1, 2, \dots$) *with* $a(\beta_c, \ell_c) \neq 0$. *If* $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, *then* $\mathfrak{v}_T^P(\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell})$ *are unbounded.*

Proof. Suppose to the contrary that $\mathfrak{v}_T^P(\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell})$ are bounded. Then since we have $\mathfrak{v}_T^P(\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell}) \geq \ell_c\mathfrak{v}_T(\mathfrak{b}_{\rm high})$, it follows that ℓ_c are bounded. By the hypothesis that the numbers of pairs (β_c, ℓ_c) are assumed to be infinite, there must be infinitely many different β_c's. We have the (virtual) dimension

$$n + \mu(\beta_c) - (\deg \mathfrak{b}_{\rm high} - 2)\ell_c - (\deg R - 2) - 2 \tag{7.17}$$

for $\mathcal{M}^{\text{main}}_{1;\ell_c+1}(L(u),\beta_c;\mathfrak{b}_{\text{high}}^{\otimes \ell}\otimes R)$. Therefore $a(\beta_c,\ell_c)\neq 0$ only when

$$0 \le n+\mu(\beta_c)-(\deg \mathfrak{b}_{\text{high}}-2)\ell_c-(\deg R-2)-2\le n. \tag{7.18}$$

Therefore since ℓ_c is bounded, the Maslov indices of β_c's must be bounded.

From (7.12) and $\mu(\beta_i)=2$, we have

$$\mu(\beta_c)=\sum_{i=1}^{m}2k_i(\beta_c)+2c_1(\alpha(\beta_c)).$$

We split our consideration into two cases:

1. If $\sum k_i(\beta_c)$ are unbounded, then the Chern numbers of $\alpha(\beta_c)$ are unbounded.
2. If $\sum k_i(\beta_c)$ are bounded, then since there are infinitely many β_c's there must be infinitely many different $\alpha(\beta_c)$'s.

We thus find that there are always infinitely many different $\alpha(\beta_c)$'s. Therefore we can conclude from Gromov compactness that the values $\omega\cap\alpha(\beta_c)$ are unbounded. But by taking the $\mathfrak{v}_T^P$-valuation of (7.15), we have

$$\mathfrak{v}_T^P(\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell})\ge \omega\cap\alpha(\beta_c)/2\pi,$$

which gives rise to a contradiction. This finishes the proof. □

For a general element $\mathfrak{b}$ in $\mathcal{A}(\Lambda_0)\setminus\mathcal{A}(\Lambda_+)$, we have the following alternatives.

Lemma 7.12. *Assume that there are infinitely many* (β_c,ℓ_c)*'s* $(c=1,2,\dots)$ *with* $a(\beta_c,\ell_c)\neq 0$*, then one of the following holds:*

1. $\mathfrak{v}_T^P(\widetilde{\mathfrak{X}}^u_{\mathfrak{b},\beta,\ell})$ *are unbounded.*
2. $\sum_{i=1}^n k_i(\beta_c)$ *are unbounded.*

Proof. If ℓ_c are bounded, then we can prove (2) in the same way as the proof of Lemma 7.11. We next assume ℓ_c are unbounded. Then $(\deg \mathfrak{b}_{\text{high}}-2)\ell_c$ are unbounded. This, combined with (7.18), implies that $\mu(\beta_c)$ are unbounded. Then it follows from (7.13), (7.14) that either (2) happens or there are infinitely many different $\alpha(\beta_c)$'s. On the other hand for the latter case, it follows from Gromov compactness that $\alpha(\beta_c)\cap\omega$ are unbounded. This implies (1) as in the end of the proof of Lemma 7.11. The proof of Lemma 7.12 is complete. □

It is easy to see that Lemmata 7.11 and 7.12 imply Proposition 7.10. □

Proposition 7.13. *We have*

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)\cdot PD[L(u)]=\delta^{\mathfrak{b},b}_{\text{can}}(\mathfrak{X}^u_{\mathfrak{b}}). \tag{7.19}$$

Proof. By Theorem 6.18 we have:

$$\delta^{\mathfrak{b},b}_{\text{can}}(\mathfrak{X}^u_{\mathfrak{b}})=\delta^{\mathfrak{b},b}_{\text{can}}(\Pi(\widetilde{\mathfrak{X}}^u_{\mathfrak{b}}))=\Pi(\delta^{\mathfrak{b},b}(\widetilde{\mathfrak{X}}^u_{\mathfrak{b}})). \tag{7.20}$$

It follows from Lemmas 7.5, 7.6 and the definition that the chain $\mathfrak{m}_{1,\beta_0}(\widetilde{\mathfrak{X}}^u_{\mathfrak{b}}))$ is a sum of the terms corresponding to (7.7)-(7.9). (Here $\mathbf{p}=\mathfrak{b}\otimes\cdots\otimes\mathfrak{b}$.)

By Lemma 7.7 the term corresponding to (7.8) vanishes.

By definition, the term corresponding to (7.7) is $\sum_{\beta\neq\beta_0}\mathfrak{m}_{1,\beta}(\widetilde{\mathfrak{X}}^u_{\mathfrak{b}})$. The term corresponding to (7.9) is $\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)$ by Lemma 7.3. Therefore we have

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)\cdot PD(L(u))=\delta^{\mathfrak{b},b}(\widetilde{\mathfrak{X}}^u_{\mathfrak{b}}).$$

The proposition now follows from (7.20). □

Proposition 7.14.

$$\mathrm{Im}\ \delta^{\mathfrak b,b}_{\mathrm{can}} \subset \mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak b}}{\partial y_i} : i = 1,\dots,n \right) H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle).$$

If $\mathfrak b \in \mathcal A(\Lambda_+)$, *we may take* Λ_0^P *in the right hand side.*

Proof. Let $G(\mathfrak b)$ be a discrete submonoid such that $\mathfrak b$ is $G(\mathfrak b)$-gapped and $G(\mathfrak b) \supset G_0$. We put $G(\mathfrak b) = \{\lambda_0, \lambda_1, \dots\}$ such that $0 = \lambda_0 < \lambda_1 < \dots$.

Let $\mathfrak I_k$ be the ideal of $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$ generated by elements

$$T^{\lambda_l} z_1^{j_1} \cdots z_m^{j_m}$$

such that $\sum_{i=1}^m j_i + l \ge k$. We recall from Definition 6.14 (see Section 27) that:

$$\mathfrak m_2^{\mathfrak b,b}(h_1,h_2) = \sum_{\ell,\beta} \frac{T^{\beta\cap\omega/2\pi}}{\ell!} \exp(\beta \cap \mathfrak b_2)\rho^b(\partial\beta)\mathfrak q_{\beta;2,\ell}(\mathfrak b_{\mathrm{high}}^{\otimes\ell}; h_1 \otimes h_2).$$

It defines a ring structure

$$h_1 \cup^{\mathfrak b,b} h_2 = (-1)^{\deg h_1(\deg h_2+1)}\mathfrak m_2^{\mathfrak b,b}(h_1,h_2) \tag{7.21}$$

on $H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle)$. Then $\cup^{\mathfrak b,b}$ is associative and $G(\mathfrak b)$-gapped. Consider the generators $\mathbf e_i$ $(i = 1,\dots,n)$ of $H(T^n;\mathbb Z)$.

Lemma 7.15. *The subring of* $(H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle), \cup^{\mathfrak b,b})$ *generated by* $\{\mathbf e_i \mid i = 1,\dots,n\}$ *is dense.*

Proof. Let $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle\langle n\rangle$ be the free associative algebra which is generated by n elements over the ring $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$. For each $\mathcal P \in \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle\langle n\rangle$ we can define $\mathcal P(\mathbf e_1,\dots,\mathbf e_n) \in H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle)$ using the product $\cup^{\mathfrak b,b}$ in an obvious way.

The lemma will be an immediate consequence of the following:

Sublemma 7.16. *For each given* k *and* $\mathfrak P \in H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle)$ *there exists* $\mathcal P \in \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle\langle n\rangle$ *such that*

$$\mathfrak P - \mathcal P(\mathbf e_1,\dots,\mathbf e_n) \in \mathfrak I_k H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle).$$

Proof. We prove this sublemma by an induction over k. Since

$$h_1 \cup^{\mathfrak b} h_2 - h_1 \wedge h_2 \in \mathfrak I_1 H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle),$$

the case $k = 1$ is a consequence that the usual cohomology ring $H(T^n;\mathbb Z)$ is generated by $\mathbf e_i$. Suppose that Sublemma 7.16 holds for $k-1$ and let

$$\mathfrak P - \mathcal P(\mathbf e_1,\dots,\mathbf e_n) \equiv \sum_{l,j_1,\dots,j_m} C(l;j_1,\dots,j_m) T^{\lambda_l} z_1^{j_1}\cdots z_m^{j_m}$$
$$\mod \mathfrak I_k H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle),$$

where the summation is taken over $l, j_1,\dots,j_m$, such that $\sum_{i=1}^m j_i + l = k-1$ and $C(l;j_1,\dots,j_m) \in H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle)$. Applying the case $k = 1$, we can find $\mathcal P(l;j_1,\dots,j_m) \in \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle\langle n\rangle$ such that

$$C(j_1,\dots,j_m) - \mathcal P(l;j_1,\dots,j_m)(\mathbf e_1,\dots,\mathbf e_n) \in \mathfrak I_1 H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle).$$

If we put

$$\mathcal{P}' = \mathcal{P} + \sum_{l,j_1,\dots,j_m} T^{\lambda_l} z_1^{j_1} \cdots z_m^{j_m} \mathcal{P}(l; j_1, \dots, j_m),$$

we have

$$\mathfrak{P} - \mathcal{P}'(\mathbf{e}_1, \dots, \mathbf{e}_n) \in \mathfrak{I}_k H(T^n; \Lambda_0^{\overset{\circ}{P}} \langle\!\langle y, y^{-1} \rangle\!\rangle)$$

by definition. Sublemma 7.16 is proved. □

Hence the proof of Lemma 7.15 is completed. □

Now, using the identity

$$\delta_{\rm can}^{\mathfrak{b},b}(h_1 \cup^{\mathfrak{b},b} h_2) = \pm \delta_{\rm can}^{\mathfrak{b},b}(h_1) \cup^{\mathfrak{b},b} h_2 \pm h_1 \cup^{\mathfrak{b},b} \delta_{\rm can}^{\mathfrak{b},b}(h_2), \tag{7.22}$$

we will prove that for each $\mathcal{P} \in \Lambda_0^{\overset{\circ}{P}} \langle\!\langle y, y^{-1} \rangle\!\rangle \langle n \rangle$

$$\delta_{\rm can}^{\mathfrak{b},b}(\mathcal{P}(\mathbf{e}_1, \dots, \mathbf{e}_n)) \in {\rm Clos}_{d_{\overset{\circ}{P}}} \left(y_i \frac{\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1, \dots, n \right) H(T^n; \Lambda_0^{\overset{\circ}{P}} \langle\!\langle y, y^{-1} \rangle\!\rangle) \tag{7.23}$$

by induction over the 'degree' of 'noncommutative polynomial' $\mathcal{P}$. We note that (7.22) is a consequence of the filtered A_∞ relation of $\mathfrak{m}^{\mathfrak{b},b}$ and the fact $\mathfrak{m}_0(1)$ and so $\mathfrak{m}_0^{\mathfrak{b},b}(1)$ is proportional to the (exact) unit. The proof of Proposition 7.14 is now complete. □

Therefore Propositions 7.14 and 7.13 imply Theorem 7.1 in the case when $\deg Q > 2$. We finally consider the case when degree of Q is 2. We put $Q = \sum_{i=1}^m Q_i \mathbf{f}_i$ and $Q^* = \sum_{i=1}^m Q_i D_i \in H_{2n-2}(D; R)$.

Lemma 7.17. *We have*

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q)(y) = \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X; L(u); \mathbb{Z})} \frac{1}{\ell!} (Q^* \cap \beta) \rho^b(\partial \beta) \exp(\mathfrak{b}_2 \cap \beta) T^{\omega \cap \beta / 2\pi}$$
$${\rm ev}_{0*}(\mathcal{M}_{1;\ell}^{\rm main}(L(u), \beta; \mathfrak{b}_{\rm high}^{\otimes \ell})).$$

Here $Q^* \cap \beta$ is defined by the intersection pairing

$$\cap : H_{2n-2}(D; \mathbb{C}) \otimes H_2(X; L(u); \mathbb{C}) \to \mathbb{C}.$$

This is well defined since $D \cap L(u) = \emptyset$.

Proof. This is a consequence of Lemma 7.4. □

Since $[Q^*] = 0$ in $H_{2n-2}(X; \mathbb{C})$ by the hypothesis, there exists R such that $\partial R = Q^*$. We have $[R] \in H_{2n-1}(X; L(u); \mathbb{C})$, using the exactness of

$$H_2(X; \mathbb{C}) \to H_2(X, L(u); \mathbb{C}) \to H_1(L(u); \mathbb{C}), \tag{7.24}$$

we can find d_i such that

$$Q^* \cap \beta = \sum_{i=1}^n d_i (\partial \beta \cap \mathbf{e}_i). \tag{7.25}$$

By differentiating (7.5), we have

$$y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} = \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X; L(u); \mathbb{Z})} \frac{T^{\omega \cap \beta / 2\pi}}{\ell!} (\partial \beta \cap \mathbf{e}_i) \rho^b(\partial \beta) \exp(\mathfrak{b}_2 \cap \beta)$$
$${\rm ev}_{0*}(\mathcal{M}_{1;\ell}^{\rm main}(L(u), \beta; \mathfrak{b}_{\rm high}^{\otimes \ell})). \tag{7.26}$$

Therefore

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(Q) = \sum_{i=1}^{n} d_i y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}.$$

The proof of Theorem 7.1 is now complete. □

8. Well-definedness of potential function

Let $\mathfrak{b}, \mathfrak{b}' \in \mathcal{A}(\Lambda_0)$, which represent the same cohomology class in $H(X;\Lambda_0)$. In this section we prove that $\mathfrak{PO}_{\mathfrak{b}}$ coincides with $\mathfrak{PO}_{\mathfrak{b}'}$ after an appropriate coordinate change of variables. Some kinds of such statement follow from the well-definedness of potential function up to filtered A_∞ homotopy equivalence, which we established in [FOOO2], for arbitrary symplectic manifold and its relatively spin Lagrangian submanifold. (See also [Aur1] for some discussion related to it.) On the other hand, for the purpose of this paper, we need to specify the type of coordinate change we use. For example, we want them to converge everywhere in the moment polytope. For this purpose we use the special feature of our situation.

We start with the definition of the class of coordinate changes.

Definition 8.1. We consider n elements $y'_i \in \Lambda^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$ $(i = 1,\dots,n)$. See Definition 4.6 for this notation.

1. We say that $y' = (y'_1,\dots,y'_n)$ is a *coordinate change converging on* $\operatorname{Int} P$ (or a *coordinate change on* $\operatorname{Int} P$) if
$$y'_i \equiv c_i y_i \mod y_i \Lambda_+^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle \tag{8.1}$$
$c_i \in \mathbb{C} \setminus \{0\}$.
2. We say that the coordinate change is *strict* if $c_i = 1$ for all i.
3. We say that the *coordinate change converges on* P if $y'_i \in \Lambda^{P}\langle\!\langle y, y^{-1}\rangle\!\rangle$ $(i = 1,\dots,n)$ in addition. Its strictness is defined in the same way. We also say that y' is a *coordinate change on* P.
4. Coordinate change y' on P is said to be *G-gapped* if $(y'_i - c_i y_i)/y_i$ is G-gapped. (See Definition 2.4.)
5. *Coordinate change converging at* u (or *coordinate change at* u) is defined as an element
$$y(u)'_i \in \Lambda_0\langle\!\langle y(u), y(u)^{-1}\rangle\!\rangle$$
such that
$$y'_i(u) \equiv c_i y_i(u) \mod y_i(u)\Lambda_+^{u}\langle\!\langle y(u), y(u)^{-1}\rangle\!\rangle.$$
Its strictness is defined in the same way.

Let $y' = (y'_1,\dots,y'_n)$ be a coordinate change converging on $\operatorname{Int} P$. If a series

$$\mathcal{P} = \sum_i a_i T^{\lambda_i} y^{v_i}$$

converges in $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$, then

$$\sum_i a_i T^{\lambda_i} (y')^{v_i} \tag{8.2}$$

also converges. Here $v_i = (v_i^{(1)}, \dots, v_i^{(n)}) \in \mathbb{Z}^n$ and

$$(y')^{v_i} = (y'_1)^{v_i^{(1)}} \cdots (y'_n)^{v_i^{(n)}}.$$

We remark that

$$(y'_i)^{-1} = c_i^{-1} y_i^{-1} \left(1 + \frac{y'_i - c_i y_i}{c_i y_i}\right)^{-1} = c_i^{-1} y_i^{-1} \sum_{k=0}^{\infty} (-1)^k \left(\frac{y'_i - c_i y_i}{c_i y_i}\right)^k$$

converges in $\mathfrak{v}_T^{\overset{\circ}{P}}$ norm by (8.1). Therefore (8.2) makes sense.

Definition 8.2. We denote (8.2) by $\mathcal{P}(y')$ and call it the *coordinate change* of $\mathcal{P}$ by y'. There are the versions corresponding to (3) or (4) of Definition 8.1, which are defined in the same way. We can define a composition of coordinate change by applying (8.2) in each factor. We write the composition of y' and y'' by $y' \circ y''$.

The proof of the following lemma is easy and so omitted.

Lemma 8.3. *Coordinate change induces a continuous endomorphism on* $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$, $\Lambda_0^{P}\langle\langle y, y^{-1}\rangle\rangle$*, and* $\Lambda_0^{u}\langle\langle y, y^{-1}\rangle\rangle$ *respectively. Therefore composition thereof induces a coordinate change again. Composition of coordinate change is associative. Composition of* G*-gapped coordinate changes are* G*-gapped.*

Convention 8.4. Let $Z_1, \dots, Z_m$ be formal variables. Let $z_1, \dots, z_m$ be the elements in $\Lambda_0\langle\langle y, y^{-1}\rangle\rangle$ defined by (2.7). For each $R \in \Lambda_0[[Z_1, \dots, Z_m]]$, we put

$$R(z) = R(z_1, \dots, z_m).$$

This is well-defined as an element of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ by Lemma 2.3.

Lemma 8.5. *Coordinate change has an inverse. Namely for* y' *there exists* y'' *such that* $y' \circ y''(y) = y'' \circ y'(y) = y$.

Proof. It suffices to prove the lemma for the case where y' is strict. We first consider the case y' is a coordinate change on P. We assume that y' is G-gapped and put $G = \{\lambda_0, \lambda_1, \dots\}$. ($\lambda_0 = 0 < \lambda_1 < \cdots$.) For each $k = 1, 2, \dots$, we are going to find a G-gapped coordinate change on P, $y''_{(k)} = (y''_{(k),1}, \dots, y''_{(k),n})$ such that

1. $y''_{(k)}$ is G-gapped and strict.
2. $((y''_{(k)} \circ y')_i - y_i)/y_i \in T^{\lambda_k}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$.

We prove this by induction over k. We assume

$$((y''_{(k)} \circ y')_i - y_i)/y_i \equiv a_i(z) T^{\lambda_k} \mod T^{\lambda_{k+1}}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$$

where $a_i \in R[Z_1, \dots, Z_m]$. We put

$$y''_{(k+1),i} = (1 - T^{\lambda_k} a_i) y''_{(k),i}.$$

Since $y'_i \equiv y''_i \equiv y_i \mod y_i\Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle$, it is easy to see that $y''_{(k+1),i}$ satisfies (1),(2).

Now $\lim_{k\to\infty} y''_{(k+1)}$ converges and gives a left inverse to y'. Existence of the right inverse can be proved in the same way. Then the right and the left inverse coincide by a standard argument of group theory.

The case of coordinate change at u is the same.

For the case of coordinate change on $\operatorname{Int} P$ we proceed as follows. We remark that if y' is a coordinate change on P, it induces an isomorphism on $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$. Now let y' be a coordinate change on $\operatorname{Int} P$. Then for each sufficiently small $\varepsilon > 0$, the coordinate change y' induces an isomorphism on $\Lambda_0^{P_\varepsilon}\langle\langle y, y^{-1}\rangle\rangle$. Therefore by taking the projective limit of these isomorphisms as $\varepsilon \to 0$, which obviously exists, the coordinate change y' induces an isomorphism on $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ by Lemma 4.4. Now it is easy to find its inverse. $\square$

Remark 8.6. The set of all coordinate changes forms a solvable group. It is related to the group of self homotopy equivalences of filtered A_∞ algebra and also relevant to the nilpotent group discussed in [KS2] and [GPS]. We remark however in [KS2] and [GPS] the coordinate changes that preserve the volume element $dy_1 \cdots dy_n/(y_1 \cdots y_n)$ are studied. We do not prove here that the coordinate change we produce (for example in Theorem 8.7 below) has this property. This point is related to the Poincaré duality and cyclic symmetry. So if we use the cyclically symmetric version of the filtered A_∞ structure that we produce in Section 18, we may obtain a coordinate change which preserves the volume element.

For $\mathcal{P} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$, we define its Jacobian ring by

$$\mathrm{Jac}(\mathcal{P}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathcal{P}}{\partial y_i} : i = 1, \ldots, n\right)}.$$

It follows from Lemmata 8.3, 8.5 that a coordinate change induces an isomorphism of Jacobian rings.

The following is the main result of this section.

Theorem 8.7. *Let $\mathfrak{b}, \mathfrak{b}' \in \mathcal{A}(\Lambda_0)$. We assume that $[\mathfrak{b}] = [\mathfrak{b}'] \in H(X;\Lambda_0)$.*

Then there exists a coordinate change y' on $\mathrm{Int}\,P$, such that

$$\mathfrak{PO}_{\mathfrak{b}}(y') = \mathfrak{PO}_{\mathfrak{b}'}(y). \tag{8.3}$$

If $\mathfrak{b} - \mathfrak{b}' \in \mathcal{A}(\Lambda_+)$, then y' can be taken to be strict.

If both $\mathfrak{b}, \mathfrak{b}' \in \mathcal{A}(\Lambda_+)$, then y' can be taken to be a strict coordinate change on P.

Example 8.8. Let $X = \mathbb{C}P^2$ with P as in Section 3. We put

$$P_i = \{(u_1, u_2) \in P \mid u_i = 0\}, \quad \mathbf{f}_i = \pi^{-1}(P_i).$$

Let $c \in \mathbb{R} \setminus \{0\}$. We consider $\mathfrak{b} = c\mathbf{f}_1$ and $\mathfrak{b}' = c\mathbf{f}_2$. Then we have

$$\mathfrak{PO}_{\mathfrak{b}} = e^c z_1 + z_2 + z_3, \qquad \mathfrak{PO}_{\mathfrak{b}'} = z_1 + e^c z_2 + z_3.$$

We put $y_1' = e^{-c} y_1$, $y_2' = e^c y_2$. It is easy to see (8.3) holds.

Proof. We first consider the case $\mathfrak{b} - \mathfrak{b}' \in \bigoplus_{k\ge 4}\mathcal{A}^k(\Lambda_0)$. Moreover we also consider $\mathfrak{b}, \mathfrak{b}' \in \bigoplus_{k\ge 4}\mathcal{A}^k(\Lambda_+)$ since the same argument applies to the case with $\mathfrak{b}$, $\mathfrak{b}' \in \mathcal{A}(\Lambda_0)$ with Λ_0^P replaced by $\Lambda_0^{\overset{\circ}{P}}$.

We take a new formal variable s and put

$$\mathfrak{b}(s) = s\mathfrak{b} + (1-s)\mathfrak{b}'.$$

Let R be a smooth singular chain $\partial R = \mathfrak{b} - \mathfrak{b}'$. For given $u \in \mathrm{Int}\,P$, we define

$$\begin{aligned}\tilde{\mathfrak{X}}^u_S(s,b) = \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} &\frac{T^{\omega\cap\beta/2\pi}}{\ell!}\rho^b(\partial\beta)\exp(\mathfrak{b}_2 \cap \beta) \\ &\mathrm{ev}_{0*}(\mathcal{M}^{\mathrm{main}}_{1;\ell+1}(L(u),\beta;\mathfrak{b}(s)^{\otimes \ell}_{\mathrm{high}} \otimes R)),\end{aligned} \tag{8.4}$$

$$\mathfrak{X}^u_S(s,b) = \Pi(\tilde{\mathfrak{X}}^u_S(s,b)), \tag{8.5}$$

where Π is as in (6.14). Here $\mathfrak{b}_{\rm high}$ is degree ≥ 4 part and $\mathfrak{b}_2$ is the degree 2 part. We remark that $\mathfrak{b}_2 = \mathfrak{b}'_2 = \mathfrak{b}(s)_2$ by the hypothesis.

We define $\Lambda_0^P\langle\langle y, y^{-1}, s\rangle\rangle$ in the same way as $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$. Then, by the same way as the proof of Proposition 7.10, we can prove that (8.4) converges in $\mathfrak{v}_T^P$ norm. Hence $\mathfrak{X}^u_S(s,b)$ can be regarded as an element of $H(L(u);\Lambda_+^P\langle\langle s,y,y^{-1}\rangle\rangle)$. In the same way as the proof of (7.2) we have

$$\mathfrak{ks}_{\mathfrak{b}(s)}(\mathfrak{b}-\mathfrak{b}') = \delta^{\mathfrak{b},b}(\mathfrak{X}^u_S(s,b)). \tag{8.6}$$

Therefore, in the same way as the proof of Proposition 7.14, we can find

$$\mathfrak{X}^u_{S,i}(s,b) \in \Lambda_0^P\langle\langle s,y,y^{-1}\rangle\rangle$$

such that

$$\delta^{\mathfrak{b},b}(\mathfrak{X}^u_S(s,b)) = \sum_i \mathfrak{X}^u_{S,i}(s,b) y_i \frac{\partial \mathfrak{PO}_{b(s)}}{\partial y_i}.$$

We remark that

$$\mathfrak{ks}_{\mathfrak{b}(s)}(\mathfrak{b}-\mathfrak{b}') = \frac{\partial \mathfrak{PO}_{\mathfrak{b}(s)}}{\partial s}. \tag{8.7}$$

Let

$$y_i(s) \in \Lambda^P\langle\langle s,y,y^{-1}\rangle\rangle$$

satisfy the same properties as the coordinate change on P. Using (8.7), we calculate

$$\frac{\partial}{\partial s}\left(\mathfrak{PO}_{\mathfrak{b}(s)}(y(s))\right) = \mathfrak{ks}_{\mathfrak{b}(s)}(\mathfrak{b}-\mathfrak{b}')\circ y(s) + \sum_i \left(\frac{\partial \mathfrak{PO}_{\mathfrak{b}(s)}}{\partial y_i}\frac{\partial y_i(s)}{\partial s}\right)\circ y(s).$$

The right hand side is zero if

$$y_i(s)\ \mathfrak{X}^u_{S,i} + \frac{\partial y_i(s)}{\partial s} = 0. \tag{8.8}$$

We will assume $y_i(s) \equiv y_i \mod y_i\Lambda_+^P\langle\langle s,y,y^{-1}\rangle\rangle$.

Motivated by (8.1) and (8.8), we set

$$Y_i(s) = \log(y_i(s)/y_i) \in \Lambda_+^P\langle\langle y,y^{-1}\rangle\rangle.$$

Then (8.8) is equivalent to

$$\frac{d}{ds}Y_i(s) + \mathfrak{X}^u_{S,i} = 0. \tag{8.9}$$

This is an ordinary differential equation and $\mathfrak{X}^u_{S,i} \in \Lambda_+^P\langle\langle s,y,y^{-1}\rangle\rangle$. Therefore it has a unique solution with $Y_i(0)=0$, which we again denote by Y_i. We now define

$$y_i(s) = \exp(Y_i(s))y_i.$$

Then we have

$$\frac{\partial}{\partial s}\left(\mathfrak{PO}_{\mathfrak{b}(s)}(y(s))\right) = 0.$$

Therefore

$$\mathfrak{PO}_{\mathfrak{b}'}(y(1)) = \mathfrak{PO}_{\mathfrak{b}(1)}(y(1)) = \mathfrak{PO}_{\mathfrak{b}(0)}(y(0)) = \mathfrak{PO}_{\mathfrak{b}}(y).$$

Hence $y' = y(1)$ is the coordinate change we are looking for.

We finally consider the case $\mathfrak{b}' - \mathfrak{b} \in \mathcal{A}^2(\Lambda_0)$. We use the notation of the proof of the corresponding case of Theorem 7.1. We put $Q = \mathfrak{b}' - \mathfrak{b}$ and put $\partial R = Q^*$. In the same way as (7.25) we can find d_i such that

$$Q^* \cap \beta = \sum_{i=1}^{n} d_i(\partial\beta \cap \mathbf{e}_i). \tag{8.10}$$

We then put

$$y_i' = e^{d_i} y_i. \tag{8.11}$$

Now we have (see (2.10))

$$\begin{aligned}\mathfrak{PO}_{\mathfrak{b}}(y') &= \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} \frac{T^{\omega\cap\beta/2\pi}}{\ell!} \exp(\mathfrak{b}_2 \cap \beta) y'(u)_1^{\partial\beta\cap\mathbf{e}_1} \cdots y'(u)_n^{\partial\beta\cap\mathbf{e}_n} \\ &\qquad \mathrm{ev}_{0*}(\mathcal{M}_{1;\ell}^{\mathrm{main}}(L(u),\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})) \\ &= \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} \frac{T^{\omega\cap\beta/2\pi}}{\ell!} \exp(\mathfrak{b}_2' \cap \beta) y(u)_1^{\partial\beta\cap\mathbf{e}_1} \cdots y(u)_n^{\partial\beta\cap\mathbf{e}_n} \\ &\qquad \mathrm{ev}_{0*}(\mathcal{M}_{1;\ell}^{\mathrm{main}}(L(u),\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})) \\ &= \mathfrak{PO}_{\mathfrak{b}'}(y).\end{aligned}$$

The proof of Theorem 8.7 is complete. □

9. Kodaira-Spencer map is a ring homomorphism

In this section we prove the following theorem.

Theorem 9.1. *The map* $\mathfrak{ks}_{\mathfrak{b}} : (H(X;\Lambda_0), \cup^{\mathfrak{b}}) \to \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ *is a ring homomorphism.*

Remark 9.2. Various statements related to this theorem has been discussed in the previous literature sometimes in a greater generality. We explain some of them in Section 31.

We fix a basis $\bar{\mathbf{f}}_i$ $i = 0, \dots, B'$ of $H(X;\mathbb{C})$ as in Section 2. We define $c_{ab}^c \in \Lambda_0$ by

$$\bar{\mathbf{f}}_a \cup^{\mathfrak{b}} \bar{\mathbf{f}}_b = \sum_{c=0}^{B'} c_{ab}^c \bar{\mathbf{f}}_c. \tag{9.1}$$

We will prove

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\bar{\mathbf{f}}_a)\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\bar{\mathbf{f}}_b) - \sum_{c=0}^{B'} c_{ab}^c \widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\bar{\mathbf{f}}_c) \equiv 0 \mod \mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1, \dots, n\right). \tag{9.2}$$

(Here $\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\bar{\mathbf{f}}_a)\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\bar{\mathbf{f}}_b)$ denotes multiplication as the elements of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.)

Let $\mathcal{M}_{1;\ell+2}^{\mathrm{main}}(\beta)$ be the moduli space of stable maps from genus zero semi-stable curve with 1 boundary and $\ell + 2$ interior marked points and of homology class β. We denote the fibre product

$$\mathcal{M}_{1;\ell+2}^{\mathrm{main}}(\beta) \times_{X^{\ell+2}} (\bar{\mathbf{f}}_a \times \bar{\mathbf{f}}_b \times \mathfrak{b}^{\ell})$$

by $\mathcal{M}_{1;\ell+2}^{\mathrm{main}}(\beta; \bar{\mathbf{f}}_a \otimes \bar{\mathbf{f}}_b \otimes \mathfrak{b}^{\otimes\ell})$. We recall from Lemma 6.5 [FOOO5] that we have already made a choice of a multisection thereon that is T^n-invariant and transversal.

We consider $\mathcal{M}^{\rm main}_{1;\ell+2}(\beta;\overline{\mathfrak f}_a\otimes\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})$ and focus on its components of (virtual) dimension n: Recall that $\mathfrak b$ lies in $H(X;\Lambda_0)$ and is a formal power series whose summands may have different degrees. For the simplicity of exposition, we will just denote union of these components by $\mathcal{M}^{\rm main}_{1;\ell+2}(\beta;\overline{\mathfrak f}_a\otimes\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})$ as long as there is no danger of confusion.

We consider the forgetful map

$$\mathfrak{forget}:\mathcal{M}^{\rm main}_{1;\ell+2}(\beta;\overline{\mathfrak f}_a\otimes\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})\to\mathcal{M}^{\rm main}_{1;2}. \tag{9.3}$$

This is a composition of the forgetful map $\mathcal{M}^{\rm main}_{1;\ell+2}(\beta;\overline{\mathfrak f}_a\otimes\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})\to\mathcal{M}^{\rm main}_{1;\ell+2}$ which forgets maps and then shrinks the resulting unstable components if any, followed by the forgetful map $\mathcal{M}^{\rm main}_{1;\ell+2}\to\mathcal{M}^{\rm main}_{1;2}$, forgetting 3rd,…,$(\ell+1)$-th marked points.

We remark that $\mathcal{M}^{\rm main}_{1;2}$ is homeomorphic to the disc D^2. In fact, $\mathcal{M}^{\rm main}_{1;1}$ is a one point and the fiber of the forgetful map $\mathcal{M}^{\rm main}_{1;2}\to\mathcal{M}^{\rm main}_{1;1}$ is a disc, which is parameterized by the position of the last interior marked point we forget.

Lemma 9.3. *$\mathcal{M}^{\rm main}_{1;2}$ has a stratification such that the interior of each stratum is described as follows.*

1. $\operatorname{Int} D^2\setminus\{0\}$.
2. *Two copies of $(-1,1)$. We call them $(-1,1)_1$, $(-1,1)_2$*
3. *Three points.* $[\Sigma_0]$, $[\Sigma_{12}]$, $[\Sigma_{21}]$.

The point $0\in D^2$ becomes $[\Sigma_0]$ in the closure of the stratum (1). The two boundary points ±1 in the closure of $(-1,1)_i$ $(i=1,2)$ become $[\Sigma_{12}]$ and $[\Sigma_{21}]$. Thus the union of the four strata $(-1,1)_1$, $(-1,1)_2$, $[\Sigma_{12}]$, $[\Sigma_{21}]$ is a circle, which becomes ∂D^2 in the closure of the stratum (1).

Each of the above strata corresponds to the combinatorial type of the elements of $\mathcal{M}^{\rm main}_{1;2}$

Proof. In addition to the boundary, $\mathcal{M}^{\rm main}_{1;2}$ has one singular point. That is the point where two interior marked points encounter. We denote this point by $[\Sigma_0]$. (Here Σ_0 is the corresponding stable bordered curve.)

The boundary of $\mathcal{M}^{\rm main}_{1;2}$ is identified with two copies of $\mathcal{M}^{\rm main}_{2;1}$. They correspond to the configuration $\Sigma=\Sigma'\cup\Sigma''$ where $[\Sigma']\in\mathcal{M}^{\rm main}_{1;1}$ and $[\Sigma'']\in\mathcal{M}^{\rm main}_{2;1}$. The two components Σ' and Σ'' are glued at their boundary marked points. There are two components depending which interior marked points of $\mathcal{M}^{\rm main}_{1;2}$ becomes the interior marked point of Σ'.

$\mathcal{M}^{\rm main}_{1;1}$ is one point. We see that $\mathcal{M}^{\rm main}_{2;1}$ is an interval as follows: We can normalize an element of it $(z_0,z^{\rm int}_1,z^{\rm int}_2)$ so the boundary marked point is $+1=z_0$ and the real part of the interior marked points $z^{\rm int}_i$ is zero. Moreover we may choose $z^{\rm int}_1+z^{\rm int}_2=0$. Thus $\mathcal{M}^{\rm main}_{2;1}$ is identified with

$$\{(z^{\rm int}_1,z^{\rm int}_2)\mid z^{\rm int}_1=\sqrt{-1}a,\ z^{\rm int}_2=-\sqrt{-1}a,\ a\in[-1,1]\}\cong[-1,1].$$

We write as $(-1,1)_i$ the interior of the components such that i-th interior marked point becomes the interior marked point of $\mathcal{M}^{\rm main}_{1;1}$.

By the above identification, $+1$ of $\partial[-1,1]_2$ corresponds to -1 of $\partial[-1,1]_1$. We write this point Σ_{12}. The other point is Σ_{21}. □



Figure 9.1

The domain of (9.3) also has a stratification according to the combinatorial type of the source. The next lemma is obvious from construction.

Lemma 9.4. *The map (9.3) is continuous and is a stratawise smooth submersion.*

Remark 9.5. When we use the smooth structure we put on $\mathcal{M}^{\text{main}}_{1;\ell+2}(\beta;\bar{\mathfrak{f}}_a\otimes\bar{\mathfrak{f}}_b\otimes\mathfrak{b}^{\otimes\ell})$ in Section A1.4 [FOOO3], then (9.3) will not be smooth in the normal direction to the strata. See [FOOO3] pages 777-778.

We consider the preimages

$$\mathfrak{forget}^{-1}([\Sigma_0])\subset\mathcal{M}^{\text{main}}_{1;\ell+2}(\beta;\bar{\mathfrak{f}}_a\otimes\bar{\mathfrak{f}}_b\otimes\mathfrak{b}^{\otimes\ell}),\quad \mathfrak{forget}^{-1}([\Sigma_{12}])\subset\mathcal{M}^{\text{main}}_{1;\ell+2}(\beta;\bar{\mathfrak{f}}_a\otimes\bar{\mathfrak{f}}_b\otimes\mathfrak{b}^{\otimes\ell}).$$

Our next task is to provide appropriate multisections on some neighborhoods of $\mathfrak{forget}^{-1}([\Sigma_0])$ and $\mathfrak{forget}^{-1}([\Sigma_{12}])$ respectively. For this purpose, we now describe the stratification of our space in these neighborhoods.

We decompose $\mathfrak{b}=\mathfrak{b}_2+\mathfrak{b}_{\text{high}}$ where $\mathfrak{b}_2$ is degree 2 and $\mathfrak{b}_{\text{high}}$ is degree ≥ 4. (We do not need to discuss degree 0 component since it does not affect Jacobian ring.)

For $\alpha \in H_2(X;\mathbb{Z})$, let $\mathcal{M}_\ell(\alpha)$ be the moduli space of stable maps from genus zero closed Riemannian surface with ℓ marked points and of homology class α. Let

$$\mathrm{ev} = (\mathrm{ev}_1, \dots, \mathrm{ev}_\ell) : \mathcal{M}_\ell(\alpha) \to X^\ell$$

be the evaluation map. We define

$$\mathcal{M}_{\ell_1+3}(\alpha; \overline{\mathfrak{f}}_a \otimes \overline{\mathfrak{f}}_b \otimes \mathfrak{b}_{\rm high}^{\otimes \ell_1}) = \mathcal{M}_{\ell_1+3}(\alpha) \ {}_{(\mathrm{ev}_1,\dots,\mathrm{ev}_{\ell_1+2})}\times_{X^{\ell_1+2}} (\overline{\mathfrak{f}}_a \times \overline{\mathfrak{f}}_b \times \mathfrak{b}_{\rm high}^{\ell_1}). \tag{9.4}$$

ev_{ℓ_1+3} defines an evaluation map

$$\mathrm{ev}_{\ell_1+3} : \mathcal{M}_{\ell_1+3}(\alpha; \overline{\mathfrak{f}}_a \otimes \overline{\mathfrak{f}}_b \otimes \mathfrak{b}_{\rm high}^{\otimes \ell_1}) \to X.$$

We next consider $\mathcal{M}^{\rm main}_{k+1;\ell_2}(\beta)$, that is the moduli space of bordered genus zero stable map with $k+1$ boundary and ℓ_2 interior marked points and of homology class $\beta \in H_2(X; L(u); \mathbb{Z})$. We consider

$$\mathcal{M}^{\rm main}_{k+1;\ell_2+1}(\beta; \mathfrak{b}_{\rm high}^{\otimes \ell_2}) = \mathcal{M}^{\rm main}_{k+1;\ell_2+1}(\beta) \ {}_{(\mathrm{ev}_1^{\rm int},\dots,\mathrm{ev}_{\ell_2}^{\rm int})}\times_{X^{\ell_2}} \mathfrak{b}_{\rm high}^{\ell_2}$$

and the evaluation map

$$\mathrm{ev} = (\mathrm{ev}^{\rm int}_{\ell_2+1}, \mathrm{ev}_1, \dots, \mathrm{ev}_k) : \mathcal{M}^{\rm main}_{k+1;\ell_2+1}(\beta; \mathfrak{b}_{\rm high}^{\otimes \ell_2}) \to X \times L(u)^{k+1}.$$

Definition 9.6. Let K be a smooth singular chain of X^2. We define

$$\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K)$$

to be the fiber product (over X^2) of

$$\mathcal{M}_{\ell_1+3}(\alpha; \overline{\mathfrak{f}}_a \otimes \overline{\mathfrak{f}}_b \otimes \mathfrak{b}_{\rm high}^{\otimes \ell_1}) \times \mathcal{M}^{\rm main}_{k+1;\ell_2+1}(\beta; \mathfrak{b}_{\rm high}^{\otimes \ell_2}) \tag{9.5}$$

with K by the map $(\mathrm{ev}_{\ell_1+3}, \mathrm{ev}^{\rm int}_{\ell_2+1})$. Using the evaluation maps at the boundary marked points of the second factor of (9.5), we have an evaluation map

$$\mathrm{ev} = (\mathrm{ev}_0, \dots, \mathrm{ev}_k) : \mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a; \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K) \to L(u)^{k+1}.$$

If K is a manifold with corner, then $\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K)$ has a Kuranishi structure. In general, when K has a triangulation $K = \cup K_i$, we have a family of Kuranishi structures $\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K_i)$ for each of the singular simplices K_i which coincide on the overlapped parts. By an abuse of notation, we simply say that $\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K)$ has a Kuranishi structure.

Lemma 9.7. *As a space with Kuranishi structure, the boundary of the moduli space* $\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; K)$ *is the union of the spaces of one of the following three types:*

(1) $\mathcal{M}_{k+1}(\alpha; \beta; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2; \partial K)$.

(2) $\mathcal{M}_{k'+1}(\alpha; \beta'; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2'; K) \ {}_{\mathrm{ev}_0}\times_{\mathrm{ev}_i} \mathcal{M}^{\rm main}_{k''+1;\ell_2''+1}(\beta''; \mathfrak{b}_{\rm high}^{\otimes \ell_2''})$,
where $k' + k'' = k+1$, $\ell_2' + \ell_2'' = \ell_2$, $\beta' \# \beta'' = \beta$, $i \in \{1, \dots, k''\}$.

(3) $\mathcal{M}^{\rm main}_{k'+1;\ell_2'+1}(\beta'; \mathfrak{b}_{\rm high}^{\otimes \ell_2'}) \ {}_{\mathrm{ev}_0}\times_{\mathrm{ev}_i} \mathcal{M}_{k''+1}(\alpha; \beta''; \overline{\mathfrak{f}}_a, \overline{\mathfrak{f}}_b; \ell_1, \ell_2''; K)$,
where $k' + k'' = k+1$, $\ell_2' + \ell_2'' = \ell_2$, $\beta' \# \beta'' = \beta$, $i \in \{1, \dots, k''\}$.

Proof. Our space $\mathcal{M}_{k+1}(\alpha; \beta; \ell_1, \ell_2; K)$ is defined by taking fiber product. Therefore its boundary is described by taking the boundary of one of the factors. The moduli space of stable maps from a *closed* curve has no boundary. The boundary of $\mathcal{M}^{\rm main}_{k+1;\ell_2}(\beta)$ is union of $\mathcal{M}^{\rm main}_{k'+1;\ell_2'}(\beta') \ {}_{\mathrm{ev}_0}\times_{\mathrm{ev}_i} \mathcal{M}^{\rm main}_{k''+1;\ell_2''}(\beta'')$ where $k' + k'' = k+1$, $\ell_2' + \ell_2'' = \ell_2$, $\beta' \# \beta'' = \beta$, $i \in \{1, \dots, k''\}$. Hence the lemma. $\square$

We now consider the case $K = \Delta$ which is the diagonal

$$\Delta = \{(x,x) \mid x \in L(u)\} \subset L(u)^2,$$

and the moduli space $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta)$.

Lemma 9.8. *There exists a surjective map*

$$\text{Glue} : \bigcup_{\alpha\#\beta'=\beta}\ \bigcup_{\ell_1+\ell_2=\ell} \mathcal{M}_{k+1}(\alpha;\beta';\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta) \to \mathfrak{forget}^{-1}([\Sigma_0]) \tag{9.6}$$

which defines an isomorphism outside codimension 2 *strata as a space with Kuranishi structure. Here we have*

$$\mathfrak{forget}^{-1}([\Sigma_0]) \subset \mathcal{M}^{\text{main}}_{k+1;\ell_1+\ell_2+2}(\beta'\#\alpha;\overline{\mathfrak{f}}_a\otimes\overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\text{high}}^{\otimes\ell}).$$

Proof. Let

$$(w_1;\Sigma^{\text{int}};\vec z^{\text{int}}) \in \mathcal{M}_{\ell_1+3}(\alpha;\overline{\mathfrak{f}}_a\otimes\overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\text{high}}^{\otimes\ell_1}),$$
$$(w_2;\Sigma^{\text{bdy}};\vec z',\vec z'^{\text{int}}) \in \mathcal{M}^{\text{main}}_{k+1;\ell_2+1}(\beta';\mathfrak{b}_{\text{high}}^{\otimes\ell_2}).$$

We assume that $\text{ev}^{\text{int}}_{\ell_1+3}(w_1;\Sigma^{\text{int}};\vec z^{\text{int}}) = \text{ev}^{\text{int}}_{\ell_2+1}(w_2;\Sigma^{\text{bdy}};\vec z',\vec z'^{,\text{int}})$. We then glue Σ^{int} and Σ^{bdy} at $z^{\text{int}}_{\ell_1+3}\in\Sigma^{\text{int}}$ and $z'^{\text{int}}_{\ell_2+1}\in\Sigma^{\text{bdy}}$ to obtain Σ.

We put $w = w_1$ on Σ^{int} and $w = w_2$ on Σ^{bdy}. We then obtain $w : \Sigma \to X$. The marked points on Σ are determined from those on Σ^{int} and Σ^{bdy} in an obvious way. We have thus defined

$$\text{Glue} : \mathcal{M}_{k+1}(\alpha;\beta';\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta) \to \mathfrak{forget}^{-1}([\Sigma_0]). \tag{9.7}$$

The map Glue is an isomorphism onto its image, as a space with Kuranishi structure. In fact, the element $(w;\Sigma;\vec z,\vec z^{\text{int}})$ in $\mathfrak{forget}^{-1}([\Sigma_0])$ is in the image of *unique* point by the map (9.6) if the following two conditions are satisfied.

Condition 9.9.
(1) Either the 1st or the second marked point of Σ^{int} lies in the same irreducible component as the (ℓ_1+3)-th interior marked point.
(2) The (ℓ_2+1)-th interior marked point of Σ^{bdy} is on a disc component, (that is not on the sphere bubble).

Clearly the set of points which does not satisfy Condition 9.9 is of codimension 2 or greater. Hence the lemma. □

Example 9.10. An example for which Condition 9.9 is not satisfied can be described as follows: Consider Σ which has one disc component D^2 and two sphere components S^2_1, S^2_2. We assume $D^2\cap S^2_1 = \{p_1\}$ = one point, $D^2\cap S^2_2=\emptyset$, $S^2_1\cap S^2_2 = \{p_2\}$ = one point. Moreover we assume the first and 2nd interior marked points z^{int}_1, z^{int}_2 are on S^2_2. Let $w : (\Sigma,\partial\Sigma)\to(X,L(u))$ be holomorphic. Then we have

$$(w;\Sigma;z_0,z^{\text{int}}_1,z^{\text{int}}_2) \in \mathfrak{forget}^{-1}([\Sigma_0]) \subset \mathcal{M}^{\text{main}}_{1;2}(\beta).$$

We put $\beta_{(0)} = w_*([D^2])$, $\alpha_1 = w_*[S^2_1]$, $\alpha_2 = w_*[S^2_2]$. ($\beta = \beta_{(0)}\#\alpha_1\#\alpha_2$.) Then we have

$$((w;S^2_1\#S^2_2;z^{\text{int}}_1,z^{\text{int}}_2,p_1),(w;D^2;z_0,p_1)) \in \mathcal{M}_1(\alpha_1\#\alpha_2;\beta_{(0)};\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta)$$

On the other hand,

$$((w;S^2_2;z^{\text{int}}_1,z^{\text{int}}_2,p_2),(w;D^2\#S^2_1;z_0,p_2)) \in \mathcal{M}_1(\alpha_2;\alpha_1\#\beta_{(0)};\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta).$$

Both of them go to $(w;\Sigma;z_0,z^{\text{int}}_1,z^{\text{int}}_2)\in\mathfrak{forget}^{-1}([\Sigma_0])$ by the map Glue.



Figure 9.2

Let us describe $\mathfrak{forget}^{-1}([\Sigma_0])$ in more detail. We consider a point $(w;\Sigma)$ on it. (For simplicity we omit marked points from our notation.) We define the decomposition

$$\Sigma = \Sigma^{\rm int} \cup \Sigma^{\rm glue} \cup \Sigma^{\rm bdy}. \tag{9.8}$$

Let $\Sigma^{\rm int,0}$ be the smallest union of components of Σ which is connected and contains the first and second interior marked points. Let $\Sigma^{\rm bdy,0}$ be the smallest union of components of Σ which is connected and contains all the disc components. Let $\Sigma^{\rm glue,0}$ be the union of sphere components in $\Sigma \setminus \Sigma^{\rm int,0} \setminus \Sigma^{\rm bdy,0}$ such that the following holds:

All the path joining $\Sigma^{\rm int,0}$ with $\Sigma^{\rm bdy,0}$ should intersect this sphere component.

The union $\Sigma^{\rm int,0} \cup \Sigma^{\rm glue,0} \cup \Sigma^{\rm bdy,0}$ is connected. The complement of this union consists of disjoint union of trees of sphere components. We include those trees to one of $\Sigma^{\rm int,0}$, $\Sigma^{\rm glue,0}$, $\Sigma^{\rm bdy,0}$ according to which it is rooted. We have thus obtained $\Sigma^{\rm int}$, $\Sigma^{\rm glue}$, $\Sigma^{\rm bdy}$ and the decomposition (9.8).

Example 9.11. In the case of Σ in Example 9.10, we have $\Sigma^{\rm int} = S^2_2$, $\Sigma^{\rm glue} = S^2_1$, $\Sigma^{\rm bdy} = D^2$.

Lemma 9.12. *$(w;\Sigma)$ satisfies Condition 9.9 if and only if $\Sigma^{\rm glue}$ is empty.*

Suppose $\Sigma^{\rm glue,0}$ is non-empty and contains N sphere components. Then the cardinality of $\text{Glue}^{-1}(w;\Sigma)$ *is $N+1$.*

Moreover the map Glue *in the neighborhood of $\mathbf{x} \in \mathfrak{forget}^{-1}([\Sigma_0])$ is described as follows: There is a Kuranishi neighborhood U of $\mathbf{x}$ in $\mathcal{M}^{\rm main}_{k+1;\ell_1+\ell_2+2}(\beta;\overline{\mathbf{f}}_a\otimes\overline{\mathbf{f}}_b\otimes\mathfrak{b}_{\rm high}^{\otimes\ell})$ such that*

$$U = \overline{U} \times \mathbb{C}^{N+1}$$

where $\mathbf{x} = (o,0)$.

$$\mathbf{x} \in \bigcap_{i=0}^{N} \text{Glue}\left(\mathcal{M}_{k+1}(\alpha_i;\beta_i;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_{i,1},\ell_{i,2};\Delta)\right)$$

if $\mathbf{x} = \mathrm{Glue}(\mathbf{x}_i)$ *with* $\mathbf{x}_i \in \mathcal{M}_{k+1}(\alpha_i;\beta_i;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_{i,1},\ell_{i,2};\Delta)$. *A Kuranishi neighborhood* U_i *of* $\mathbf{x}_i$ *is identified with*

$$U_i = \overline{U} \times (\mathbb{C}^i \times \{0\} \times \mathbb{C}^{N-i}),$$

where U_i *is embedded in* U *by the obvious inclusion.*

The space $\mathfrak{forget}^{-1}([\Sigma_0])$ *is described in our Kuranishi neighborhood by*

$$\left(\left(\bigcup_{i=0}^{N} U_i\right) \cap s_{\mathbf{x}}^{-1}(0)\right)/\Gamma_{\mathbf{x}}.$$

Here $s_{\mathbf{x}}$ *is the Kuranishi map and* $\Gamma_{\mathbf{x}}$ *is the finite group of symmetry of* $(w;\Sigma)$.

Remark 9.13. We can summarize Lemma 9.12 as follows: Our space $\mathfrak{forget}^{-1}([\Sigma_0])$ is a 'normal crossing divisor' and $N+1$ components intersect transversally at $\mathbf{x}$. (We put the phrase 'normal crossing divisor' in quote since there is actually a Kuranishi map and obstruction bundle.)

Proof. Let S^2_i $(i = 1,\dots,N)$ be the components of $\Sigma^{\mathrm{glue},0}$. We include each of the tree of components in $\Sigma^{\mathrm{glue}} \setminus \Sigma^{\mathrm{glue},0}$ to the component S^2_i where this tree is rooted. We then have a decomposition

$$\Sigma^{\mathrm{glue}} = \bigcup_{i=1}^{N} \Sigma^{\mathrm{glue}}_i.$$

We may enumerate them so that

$$\begin{aligned}
&\Sigma^{\mathrm{glue}}_1 \cap \Sigma^{\mathrm{bdy}} = \{z_0\},\\
&\Sigma^{\mathrm{glue}}_i \cap \Sigma^{\mathrm{glue}}_{i+1} = \{z_i\}, \quad i = 1,\dots,N-1\\
&\Sigma^{\mathrm{glue}}_N \cap \Sigma^{\mathrm{int}} = \{z_N\}.
\end{aligned}$$

For $i = 0,\dots,N$, we put

$$\beta_{(i)} = w_*([\Sigma^{\mathrm{bdy}}]) + \sum_{j=1}^{i} w_*([\Sigma^{\mathrm{glue}}_j]),$$

$$\alpha_i = \sum_{j=i+1}^{N} w_*([\Sigma^{\mathrm{glue}}_j]) + w_*([\Sigma^{\mathrm{int}}]).$$

If we cut Σ at z_i, then we obtain an element of $\mathcal{M}_{k+1}(\alpha_i;\beta_i;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_{i,1},\ell_{i,2};\Delta)$. It is clear that those elements are all the elements of $\mathrm{Glue}^{-1}(w;\Sigma)$.

The factor $\overline{U}$ is the product of the Kuranishi neighborhoods of $(w;\Sigma^{\mathrm{bdy}})$, $(w;\Sigma^{\mathrm{glue}}_i)$, $(w;\Sigma^{\mathrm{int}})$ in the moduli spaces in which they are contained. Other than $\overline{U}$, the Kuranishi neighborhood of $\mathbf{x}$ in $\mathcal{M}^{\mathrm{main}}_{k+1;\ell_1+\ell_2+2}(\beta;\overline{\mathfrak{f}}_a\otimes\overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})$ has a parameter. That is the parameter to glue $N+2$ components at the singular points $z_0,\dots,z_N$. They give the factor $\mathbb{C}^{N+1}$.

If we resolve all the singularities z_j $(j = 0,\dots,N)$ other than z_i, we still obtain an element of $\mathfrak{forget}^{-1}([\Sigma_0])$. Moreover it is in the image of $\mathcal{M}_{k+1}(\alpha_i;\beta_i;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_{i,1},\ell_{i,2};\Delta)$. Therefore the lemma follows.

(We remark that our Kuranishi structure is constructed inductively over the strata in the way we have described above. See [FO],[FOOO2].) ☐

Remark 9.14. Actually the T^n-equivariant Kuranishi structure we produced in [FOOO4] Section 15 is *not* compatible with the fiber product description in Lemmas 9.7, 9.6, and 9.12. We will explain the way to go around this point in detail in Section 28.

Lemma 9.15. *There exists a system of multisections on a neighborhood of* $\mathfrak{forget}^{-1}([\Sigma_0])$ *with the following properties:*

1. *It is transversal to* 0 *and is* T^n*-invariant.*
2. *It is compatible at the boundaries described in Lemma 9.7. (Note* $\partial\Delta = \emptyset$ *so* (1) *in Lemma 9.7 does not occur.)*
3. *It is compatible with the forgetful map of the 1st, ..., k-th boundary marked points.*
4. *It is invariant under the arbitrary permutations of the interior marked points.*
5. *It is transversal to the map* (9.7) *in the sense we describe below:*

Let us state Condition (5) precisely. Let U' be a Kuranishi neighborhood of a point $\mathbf{x}$ in $\mathcal{M}_{k+1}(\alpha_i;\beta_i;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_{i,1},\ell_{i,2};\Delta)$. Let $\mathrm{Glue}(\mathbf{x}) = \mathbf{y} \in \mathcal{M}^{\mathrm{main}}_{k+1;\ell_1+\ell_2+2}(\beta;\overline{\mathfrak{f}}_a\otimes \overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})$. Let U be a Kuranishi neighborhood of $\mathbf{y}$ and E an obstruction bundle. Let $\mathfrak{s}_i$ be a branch of our multisection. We require that $\mathfrak{s}_i^{-1}(0)$ is transversal to the map : $U' \to U$, which is induced by Glue.

Proof. The proof can be done by an induction over $\beta\cap\omega$ and the number of boundary marked points. The description in Lemma 9.12 shows that we can perform this construction locally. Then general strategy of the inductive construction of multisection applies. □

We have thus constructed multisection on a neighborhood of $\mathfrak{forget}^{-1}([\Sigma_0])$.

Remark 9.16. The multisection on a neighborhood of $\mathfrak{forget}^{-1}([\Sigma_0])$ which we have just constructed induces a multisection on $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta)$ by pullback under the map Glue. On the other hand, by definition, $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathfrak{f}}_a,\overline{\mathfrak{f}}_b;\ell_1,\ell_2;\Delta)$ is the fiber product of two spaces

$$\mathcal{M}_{\ell_1+3}(\alpha;\overline{\mathfrak{f}}_a\otimes\overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_1}),\quad \mathcal{M}^{\mathrm{main}}_{k+1;\ell_2+1}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_2}) \tag{9.9}$$

over the diagonal. We already have multisections ('$\mathfrak{q}$' multisections) (see Definition 6.6.) on each of (9.9). However, the multisection induced by Glue does not necessarily coincide with the fiber product multisection. In fact, it seems difficult to construct a system of multisections ('$\mathfrak{q}$' multisections) so that the fiber product multisection of (9.9) along the diagonal is transversal. This is related to Remark 11.4 [FOOO4].

We next construct a multisection of a neighborhood of $\mathfrak{forget}^{-1}([\Sigma_{12}])$. Let

$$[w;\Sigma] \in \mathfrak{forget}^{-1}([\Sigma_{12}]) \subset \mathcal{M}^{\mathrm{main}}_{1;\ell+2}(\beta;\overline{\mathfrak{f}}_a\otimes\overline{\mathfrak{f}}_b\otimes\mathfrak{b}_{\mathrm{high}}^{\otimes\ell}).$$

By the definition of the forgetful map, we have the natural map $\Sigma \to \Sigma_{12}$. We recall that Σ_{12} has three components, each of which contains one of the three marked points of Σ_{12}. We call them Σ_{12,z_0}, $\Sigma_{12,z_1^{\mathrm{int}}}$, $\Sigma_{12,z_2^{\mathrm{int}}}$, according to the marked points which they contain. We decompose Σ into the union of three components Σ_{z_0}, $\Sigma_{z_1^{\mathrm{int}}}$, $\Sigma_{z_2^{\mathrm{int}}}$ so that they are mapped to the components Σ_{12,z_0}, $\Sigma_{12,z_1^{\mathrm{int}}}$, $\Sigma_{12,z_2^{\mathrm{int}}}$ of Σ_{12} by the forgetful map, respectively.


z_1^{int}
$\Sigma_{21,z_1^{\mathrm{int}}}$
z_2^{int}
$\Sigma_{12,z_2^{\mathrm{int}}}$
z_0
Σ_{12,z_0}

Figure 9.3

Remark 9.17. The decomposition $\Sigma = \Sigma_{z_0} \cup \Sigma_{z_1^{\mathrm{int}}} \cup \Sigma_{z_2^{\mathrm{int}}}$ is *not* unique.

We consider the conditions

$$w_*([\Sigma_{z_1^{\mathrm{int}}}]) = \beta_{(1)}, \quad w_*([\Sigma_{z_2^{\mathrm{int}}}]) = \beta_{(2)}, \quad w_*([\Sigma_{z_0}]) = \beta_{(0)}. \tag{9.10}$$

Definition 9.18. We denote by $\mathfrak{X}(\beta_{(1)}, \beta_{(2)}, \beta_{(0)})$ the subset of the elements of $\mathfrak{forget}^{-1}([\Sigma_{12}])$ that satisfy (9.10). We thus have a decomposition

$$\mathfrak{forget}^{-1}([\Sigma_{12}]) = \bigcup_{\beta_{(1)}+\beta_{(2)}+\beta_{(0)}=\beta} \mathfrak{X}(\beta_{(1)}, \beta_{(2)}, \beta_{(0)}). \tag{9.11}$$

In other words, we have a map

$$\begin{aligned}\mathrm{Glue} : \Big(\mathcal{M}_{k_1+1;\ell_1+1}(\beta_{(1)}; \bar{\mathfrak{f}}_a \otimes \mathfrak{b}_{\mathrm{high}}^{\otimes \ell_1}) \times \mathcal{M}_{k_2+1;\ell_2+1}(\beta_{(2)}; \bar{\mathfrak{f}}_b \otimes \mathfrak{b}_{\mathrm{high}}^{\otimes \ell_2})\Big) \\ {}_{(\mathrm{ev}_0,\mathrm{ev}_0)} \times_{(\mathrm{ev}_i,\mathrm{ev}_j)} \mathcal{M}_{k_3+3;\ell_3}(\beta_{(0)}; \mathfrak{b}_{\mathrm{high}}^{\otimes \ell_3}) \to \mathfrak{X}\left(\beta_{(1)}, \beta_{(2)}, \beta_{(0)}\right).\end{aligned} \tag{9.12}$$

It is actually an isomorphism to its image.

Since the decomposition $\Sigma = \Sigma_{z_0} \cup \Sigma_{z_1^{\mathrm{int}}} \cup \Sigma_{z_2^{\mathrm{int}}}$ is not unique as we mentioned in Remark 9.17, the images of the maps Glue could overlap.

Example 9.19. Let Σ be a union of 4 disks $D_0, \dots, D_3$. We assume $D_1 \cap D_3 = \{p_{13}\}$, $D_0 \cap D_3 = \{p_{03}\}$, $D_0 \cap D_2 = \{p_{02}\}$ and other intersections are empty. We take $z_1^{\rm int} \in D_1$, $z_2^{\rm int} \in D_2$ and $z_0 \in D_0$. Let $w : \Sigma \to X$ be a holomorphic map. Denote $\beta_{(i)} = w_*([D_i])$. Then

$$\begin{aligned}&(((w; D_1\# D_3; z_1^{\rm int}; p_{03}), (w; D_2; z_2^{\rm int}; p_{02})), (w; D_0; z_0, p_{03}, p_{02}))\\ &\in (\mathcal{M}_{1;1}(\beta_{(1)} + \beta_{(3)}; \overline{\mathbf{f}}_a) \times \mathcal{M}_{1;1}(\beta_{(2)}; \overline{\mathbf{f}}_b))_{(\mathrm{ev}_0,\mathrm{ev}_0)} \times_{(\mathrm{ev}_1,\mathrm{ev}_2)} \mathcal{M}_{3;0}(\beta_{(0)}).\end{aligned} \tag{9.13}$$

On the other hand,

$$\begin{aligned}&(((w; D_1; z_1^{\rm int}; p_{13})), (w; D_2; z_2^{\rm int}; p_{02})), (w; D_3\# D_0; z_0, p_{13}, p_{02}))\\ &\in (\mathcal{M}_{1;1}(\beta_{(1)}; \overline{\mathbf{f}}_a) \times \mathcal{M}_{1;1}(\beta_{(2)}; \overline{\mathbf{f}}_b))_{(\mathrm{ev}_0,\mathrm{ev}_0)} \times_{(\mathrm{ev}_1,\mathrm{ev}_2)} \mathcal{M}_{3;0}(\beta_{(0)} + \beta_{(3)}).\end{aligned} \tag{9.14}$$

They both map to the same element under the map Glue.



Figure 9.4

Here, however, the situation is different from the one for the case of $\mathfrak{forget}^{-1}([\Sigma_0])$. Namely, we can choose our multisection on a neighborhood of $\mathfrak{forget}^{-1}([\Sigma_{12}])$ so that (9.12) preserves the multisection. (Compare Remark 9.16.) In fact, the left hand side of (9.13) is contained in

$$(\mathcal{M}_{1;1}(\beta_1; \overline{\mathbf{f}}_a)_{\mathrm{ev}_0} \times_{\mathrm{ev}_1} \mathcal{M}_{2;0}(\beta_3)) \times \mathcal{M}_{1;1}(\beta_2; \overline{\mathbf{f}}_b))_{(\mathrm{ev}_0,\mathrm{ev}_0)} \times_{(\mathrm{ev}_1,\mathrm{ev}_2)} \mathcal{M}_{3;0}(\beta_0), \tag{9.15}$$

and the left hand side of (9.14) is contained in

$$(\mathcal{M}_{1;1}(\beta_1; \overline{\mathbf{f}}_a)) \times \mathcal{M}_{1;1}(\beta_2; \overline{\mathbf{f}}_b))_{(\mathrm{ev}_0,\mathrm{ev}_0)} \times_{(\mathrm{ev}_1,\mathrm{ev}_2)} (\mathcal{M}_{2;0}(\beta_3)_{\mathrm{ev}_1} \times_{\mathrm{ev}_1} \mathcal{M}_{3;0}(\beta_0)). \tag{9.16}$$

These two spaces are isomorphic together with their Kuranishi structures. (This is because of the associativity of the fiber product.) Moreover the '$\mathfrak{q}$'-multisections on them are compatible. We remark that '$\mathfrak{q}$'-multisections are constructed inductively over the number of *disc* components (and of symplectic area). Such induction was possible because ev_0 (the evaluation map at the boundary marked point) automatically becomes a submersion by T^n equivariance. Since the fiber products appearing in both (9.15) and (9.16) correspond to gluing at the boundary singular point, they are compatible with the '$\mathfrak{q}$'-multisections.

We have the following lemma which corresponds to Lemma 9.12.

Lemma 9.20. *Each of* $\mathfrak{X}(\beta_{(1)}, \beta_{(2)}, \beta_{(0)})$ *has Kuranishi structure with corners. Let* $\mathbf{x} \in \mathfrak{forget}^{-1}([\Sigma_{12}])$. *Then there exist positive integers* N_1, N_2 *such that the following holds.*

1. *A Kuranishi neighborhood* U *of* $\mathbf{x}$ *in* $\mathcal{M}^{\mathrm{main}}_{k+1;\ell+2}(\beta; \overline{\mathbf{f}}_a \otimes \overline{\mathbf{f}}_b \otimes \mathfrak{b}^{\otimes \ell})$ *is* $\overline{U} \times [0,1)^{N_1+1} \times [0,1)^{N_2+1}$.
2. *The space* $\mathfrak{forget}^{-1}([\Sigma_{12}])$ *in this Kuranishi neighborhood is*
$$\left(\left(\overline{U} \times \partial([0,1)^{N_1}) \times \partial([0,1)^{N_2})\right) \cap s^{-1}(0)\right)/\Gamma_{\mathbf{x}}.$$
3. $\mathbf{x}$ *is contained in the union of* $(N_1+1) \times (N_2+1)$ *different strata, that is, in* $\mathfrak{X}(\beta_{(k_1,k_2,1)}, \beta_{(k_1,k_2,2)}, \beta_{(k_1,k_2,0)})$. *(*$k_1 = 1, \dots, N_1$, $k_2 = 1, \dots, N_2$.*)*
4. $\mathbf{x}$ *lies on the codimension* $N_1 + N_2$ *corner on each stratum in* (3), *that is* $\mathfrak{X}(\beta_{(k_1,k_2,1)}, \beta_{(k_1,k_2,2)}, \beta_{(k_1,k_2,0)})$.
5. *The Kuranishi neighborhood of* $\mathbf{x}$ *in* $\mathfrak{X}(\beta_{(k_1,k_2,1)}, \beta_{(k_1,k_2,2)}, \beta_{(k_1,k_2,0)})$ *can be identified with*
$$\overline{U} \times ([0,1)^{k_1} \times \{0\} \times [0,1)^{N_1-k_1}) \times ([0,1)^{k_2} \times \{0\} \times [0,1)^{N_2-k_2}).$$
6. *The obstruction bundle of* $\mathfrak{X}(\beta_{(k_1,k_2,1)}, \beta_{(k_1,k_2,2)}, \beta_{(k_1,k_2,0)})$ *is obtained by restricting the obstruction bundle of* $\mathcal{M}^{\mathrm{main}}_{k+1;\ell+2}(\beta; \overline{\mathbf{f}}_a \otimes \overline{\mathbf{f}}_b \otimes \mathfrak{b}^{\otimes \ell})$.

The proof of this lemma is similar to that of Lemma 9.12 and omitted.

Lemma 9.21. *The multisections on* $\mathfrak{X}(\beta_{(k_1,k_2,1)}, \beta_{(k_1,k_2,2)}, \beta_{(k_1,k_2,0)})$ *induced by* (*9.12*) *from the '*$\mathfrak{q}$*' multisection are compatible on the overlapped part and to the '*$\mathfrak{q}$*'-multisections of* $\mathcal{M}^{\mathrm{main}}_{1;\ell+2}(\beta; \overline{\mathbf{f}}_a \otimes \overline{\mathbf{f}}_b \otimes \mathfrak{b}^{\otimes \ell})$. *Hence it was already extended to the neighborhood of* $\mathfrak{forget}^{-1}([\Sigma_{12}])$.

Proof. As we explained in the paragraph right after Example 9.19, this is a consequence of our construction of '$\mathfrak{q}$'-multisections. ☐

Thus we have defined compatible multisections $\mathfrak{s}$ which we need. We can then define integration along the fiber of the map

$$\mathrm{ev}_0 : \mathcal{M}^{\mathrm{main}}_{1;\ell+2}(\beta; \overline{\mathbf{f}}_a \otimes \overline{\mathbf{f}}_b \otimes \mathfrak{b}^{\otimes \ell}) \to L(u)$$

restricted to

$$\mathfrak{s}^{-1}(0) \cap \mathfrak{forget}^{-1}([\Sigma_0]), \quad \mathfrak{s}^{-1}(0) \cap \mathfrak{forget}^{-1}([\Sigma_{12}]).$$

We use this construction in the next two lemmata. Hereafter we omit $\mathfrak{s}$ from the notation. Namely we will just write as $\mathfrak{forget}^{-1}([\Sigma_0])$ etc. in place of $\mathfrak{s}^{-1}(0) \cap \mathfrak{forget}^{-1}([\Sigma_0])$ etc.

Lemma 9.22. *We have*

$$(\mathrm{ev}_0)_* \left(\mathfrak{forget}^{-1}([\Sigma_0])\right) = (\mathrm{ev}_0)_* \left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right).$$

Here we recall that in this section we consider the case where $\mathfrak{forget}^{-1}([\Sigma_0])$ is n dimensional and so

$$(\mathrm{ev}_0)_* \left(\mathfrak{forget}^{-1}([\Sigma_0])\right) \in H_n(L(u); \Lambda_0) \cong H^0(L(u); \Lambda_0) = \Lambda_0.$$

Similarly $(\mathrm{ev}_0)_* \left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right) \in \Lambda_0$.

Lemma 9.23. *If $a, b \neq 0$, we have:*

$$\begin{aligned}&(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right)\\ &= \sum_{\beta_{(1)}+\beta_{(2)}=\beta}\ \sum_{\ell_1+\ell_2=\ell}(\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell_1+1}(\beta_{(1)};\overline{\mathfrak f}_a\otimes\mathfrak b^{\otimes\ell_1})\right)\\ &\qquad\qquad\times(\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell_2+1}(\beta_{(2)};\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell_2})\right).\end{aligned}$$

Here summation in the right hand side is taken over those $\mathcal{M}^{\mathrm{main}}_{1,\ell_1+1}(\beta_{(1)};\overline{\mathfrak f}_1\otimes\mathfrak b^{\otimes\ell_1})$ which are n-dimensional. (We remark $\overline{\mathfrak f}_0 = PD[X]$.)

If $a = 0$ and $b \neq 0$, then

$$(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right) = (\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell+1}(\beta);\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})\right).$$

The case when $a \neq 0$ and $b = 0$ is similar.

If $a = b = 0$, then

$$(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right) = 1.$$

Remark 9.24. We remark that we have not broken T^n symmetry yet.

Proof of Lemma 9.22. We first take $[\Sigma_2] \in \mathcal{M}^{\mathrm{main}}_{1;2}$ which is in the stratum (1) in Lemma 9.3 and is sufficiently close to $[\Sigma_{12}]$.

Sublemma 9.25.

$$(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_2])\right) = (\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right).$$

Proof. Let $\mathfrak z \in L(u)$ be a generic point. We remark that $(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right)$ may be regarded as a T^n-invariant differential 0 form, that is nothing but a constant function. This implies that the order counted with sign of the moduli space $(\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_{12}])$ is the right hand side of Lemma 9.22. Namely:

$$\#\left((\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_{12}])\right) = (\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_{12}])\right).$$

In the same way we have

$$\#\left((\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_2])\right) = (\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_2])\right).$$

By choosing $\mathfrak z$ generic, we may assume that any element of $(\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_{12}])$ satisfies analogous condition as Condition 9.9. We may also assume that any element of $(\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_{12}])$ is a map from a union of three discs. Then, by a standard gluing argument, we have

$$\#\left((\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_{12}])\right) = \#\left((\mathrm{ev}_0)^{-1}(\mathfrak z)\cap\mathfrak{forget}^{-1}([\Sigma_2])\right).$$

Sublemma 9.25 follows. □

Let $[\Sigma_3] \in \mathcal{M}^{\mathrm{main}}_{1;2}$. We assume that it is sufficiently close to $[\Sigma_0]$ and is in the stratum (1) in Lemma 9.3. In the same way as Sublemma 9.25, we have

$$(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_3])\right) = (\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_0])\right). \tag{9.17}$$

We next take generic smooth path $\gamma : [0,1] \to \mathcal{M}^{\mathrm{main}}_{1;2}$ joining $[\Sigma_2]$ and $[\Sigma_3]$ and contained in the stratum (1) in Lemma 9.3. By Lemma 9.4

$$[0,1]\ {}_\gamma\times_{\mathfrak{forget}}\ \mathcal{M}^{\mathrm{main}}_{1;\ell+2}(\beta;\overline{\mathfrak f}_a\otimes\overline{\mathfrak f}_b\otimes\mathfrak b^{\otimes\ell})$$

has a Kuranishi structure and we may choose $\mathfrak s$ and γ so that we can use Stokes' theorem (Lemma 12.18 [FOOO5]) on $[0,1]\ {}_\gamma\times_{\mathfrak{forget}}\ \mathfrak s^{-1}(0)$ to obtain

$$(\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_3])\right) = (\mathrm{ev}_0)_*\left(\mathfrak{forget}^{-1}([\Sigma_2])\right). \tag{9.18}$$

(9.17), (9.18) and Sublemma 9.25 immediately imply Lemma 9.22. ☐

Proof of Lemma 9.23. Using the decomposition (9.11), we study the contributions of various components of $\mathfrak{X}(\beta_{(1)},\beta_{(2)},\beta_{(0)})$ to the left hand side of Lemma 9.23. We first consider the case $\beta_{(0)}=0$. Using Lemma 9.21 it is easy to see that

$$\begin{aligned}&(\mathrm{ev}_0)_*(\mathfrak{X}(\beta_{(1)},\beta_{(2)},0))\\&=\sum_{\ell_1+\ell_2=\ell}(\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell_1+1}(\beta_{(1)};\bar{\mathbf{f}}_a\otimes\mathfrak{b}^{\otimes\ell_1})\right)\times(\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell_2+1}(\beta_{(2)};\bar{\mathbf{f}}_b\otimes\mathfrak{b}^{\otimes\ell_2})\right),\end{aligned}$$

if $a,b\neq 0$.

If $a=0$, then only the case $\beta_{(1)}=0=\ell_1$ can contribute and

$$(\mathrm{ev}_0)_*(\mathcal{M}^{\mathrm{main}}_{1,1}(\beta_{(1)};\bar{\mathbf{f}}_0\times\mathfrak{b}^{\otimes 0}))=1.$$

In case $a=b=0$, only the case $\beta_{(1)}=\beta_{(2)}=0=\ell_1=\ell_2$ contributes and it gives 1.

We next study the case $\beta_{(0)}\neq 0$. We observe that the differential form

$$(\mathrm{ev}_0)_*\left(\mathcal{M}^{\mathrm{main}}_{1,\ell_1+1}(\beta_{(1)};\bar{\mathbf{f}}_1\otimes\mathfrak{b}^{\otimes\ell_1})\right)\tag{9.19}$$

is degree 0 and constant, (if it is nonzero). This is a consequence of the fact that

$$\mathrm{ev}_0:\mathcal{M}^{\mathrm{main}}_{1,\ell_1+1}(\beta_{(1)};\bar{\mathbf{f}}_a\otimes\mathfrak{b}^{\otimes\ell_1})\to L(u)$$

is a submersion. (We remark again that we did not break T^n symmetry yet.) We write (9.19) as $c_{1,\ell_1,\beta_{(1)}}1_{L(u)}$. We define $c_{2,\ell_2,\beta_{(2)}}$ in a similar way. Now by Lemma 12.20 [FOOO5] we have

$$\begin{aligned}&(\mathrm{ev}_0)_*(\mathfrak{X}(\beta_{(1)},\beta_{(2)},\beta_{(0)}))\\&=\sum_{\ell_0+\ell_1+\ell_2=\ell}c_{1,\ell_1,\beta_{(1)}}c_{2,\ell_2,\beta_{(2)}}(\mathrm{ev}_0)_*(\mathrm{ev}_1,\mathrm{ev}_2)^*(1_{L(u)}\times 1_{L(u)}).\end{aligned}$$

Here

$$(\mathrm{ev}_1,\mathrm{ev}_2):\mathcal{M}^{\mathrm{main}}_{3,\ell_0}(\beta_{(0)};\mathfrak{b}^{\otimes\ell_0})\to L(u)^2.\tag{9.20}$$

Since $1_{L(u)}$ is the strict unit, this is zero unless $\beta_{(0)}=0$. (In other words, (9.20) factors through $\mathcal{M}^{\mathrm{main}}_{3,\ell_0}(\beta_{(0)};\mathfrak{b}^{\otimes\ell_0})\to\mathcal{M}^{\mathrm{main}}_{1,\ell_0}(\beta_{(0)};\mathfrak{b}^{\otimes\ell_0})$. So this is zero unless $\beta_{(0)}=0$.) The proof of Lemma 9.23 is complete. ☐

It is easy to see that an appropriate weighted sum of the right hand side of Lemma 9.23 becomes the first product term in the left hand side of equality (9.2). See (2.10). In the rest of this section we will check that the left hand side of Lemma 9.22 becomes the second term of (9.2) modulo an element of the defining ideal of the Jacobian ring, after taking appropriate weighted sum. The proof is a combination of the proofs of associativity of the quantum cup product and of commutativity of the diagram in Theorem 7.1.

We already defined the moduli space $\mathcal{M}_{k+1}(\alpha;\beta;\bar{\mathbf{f}}_a,\bar{\mathbf{f}}_b;\ell_1,\ell_2;K)$ for a smooth singular chain $K\subset X^2$. So far we considered the case $K=\Delta$, which is T^n-invariant. In the next step, we need to use K which is not T^n-invariant. It prevents us from taking a T^n-invariant multisection on $\mathcal{M}_{k+1}(\alpha;\beta;\bar{\mathbf{f}}_a,\bar{\mathbf{f}}_b;\ell_1,\ell_2;K)$ any more. So by taking a *single* multisection, there is no way to make ev_0 to be a submersion. On the other hand, since we are working on de Rham theory, we need to take integration along the fiber by the map ev_0. For this purpose we use the technique of *continuous* *family of multisections.* We discussed it in detail in Section 12 [FOOO5]. See also Section 30 of the present paper.

Condition 9.26. We consider a system of continuous family of multisections $\mathfrak{s}$ on various $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;K)$ and $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\partial K)$ with the following properties.

(1)
$$\mathrm{ev}_0 : \mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;K)^{\mathfrak{s}} \to L(u)$$
and
$$\mathrm{ev}_0 : \mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\partial K)^{\mathfrak{s}} \to L(u)$$
are submersions. Here $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;K)^{\mathfrak{s}}$ denotes the zero set of the multisection $\mathfrak{s}$ of the Kuranishi structure of the moduli space
$$\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;K).$$

(2) They are compatible at the boundaries described in Lemma 9.7.

(3) They are compatible with the forgetful maps of the boundary marked points.

We remark that since $\partial\partial K = 0$ we do not have (1) in Lemma 9.7 for the case of ∂K.

Lemma 9.27. *If we have a system of continuous families of multisections satisfying Condition 9.26 for* $(\partial K, 0)$*, then we can find a system of continuous families of multisections for* $(K, \partial K)$ *satisfying Condition 9.26 which extends the given one on* ∂K*.*

Proof. By an induction over $\beta \cap \omega$, ℓ_1, ℓ_2, and k, the lemma follows easily from Lemma 12.14 [FOOO5]. □

Now we go back to the proof of Theorem 9.1. Let $\Delta = \{(x,x) \in X^2 \mid x \in X\}$ be the diagonal. Since $\overline{\mathbf{f}}_i$ $(i = 0,\dots,B')$ form a basis of $H(X;\mathbb{C})$, there exists a smooth singular chain R on X^2 such that

$$\partial R = \Delta - \sum_{i,j=0}^{B'} (-1)^{\deg\overline{\mathbf{f}}_i \deg\overline{\mathbf{f}}_j} g^{ij} (\overline{\mathbf{f}}_i \times \overline{\mathbf{f}}_j), \tag{9.21}$$

where $g_{ij} = \langle \overline{\mathbf{f}}_i, \overline{\mathbf{f}}_j \rangle_{\mathrm{PD}_X}$ with $0 \le i,\, j \le B'$, and (g^{ij}) is the inverse of the matrix (g_{ij}). (See Lemma 26.7 for the sign in (9.21).)

Remark 9.28. We remark that R can not be taken to be T^n-invariant.

We consider $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\partial R)$. We remark that we have already fixed a multisection on $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\Delta)$ such that ev_0 is a submersion. (This is a single multisection and is not a continuous family. But it can be regarded as a special case of family of multisections.) By Definition 9.6 we have

$$\begin{aligned}&\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\overline{\mathbf{f}}_i \times \overline{\mathbf{f}}_j)\\ &= \mathcal{M}_{\ell_1+3}(\alpha;\overline{\mathbf{f}}_a \otimes \overline{\mathbf{f}}_b \otimes \overline{\mathbf{f}}_i \otimes \mathfrak{b}_{\mathrm{high}}^{\otimes \ell_1}) \times \mathcal{M}_{k+1;\ell_2+1}(\beta;\overline{\mathbf{f}}_j \otimes \mathfrak{b}_{\mathrm{high}}^{\otimes \ell_2}).\end{aligned} \tag{9.22}$$

Here the product is the *direct* product. On the second factor of (9.22), the (single) multisection (that is the '$\mathfrak{q}$'-multisection) was already chosen in Lemma 6.5 [FOOO5] so that ev_0 is a submersion. We take a (single) multisection on the first factor, such that it becomes transversal to 0.

Therefore we apply Lemma 9.27 to obtain a system of continuous family of multisections $\mathfrak{s}$ on $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;R)$ compatible with other choices we made and such that ev_0 is a submersion on its zero set.

Therefore we can take an integration along the fiber for $\mathcal{M}_{k+1}(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;R)$. (We omit $\mathfrak{s}$ from notation here.) (See Definition 12.11 [FOOO5].)

Definition 9.29. We put

$$\begin{aligned}\widetilde{\mathfrak{X}}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})(b) = \sum_{\ell_1,\ell_2}\sum_{\alpha,\beta}\frac{T^{(\alpha\#\beta)\cap\omega/2\pi}}{(\ell_1+\ell_2)!}\rho^b(\partial\beta)\exp(\mathfrak{b}_2\cap(\alpha\#\beta))\\ (\mathrm{ev}_0)_*(\mathcal{M}_1(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;R)),\end{aligned} \tag{9.23}$$

and

$$\mathfrak{X}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})(b) = \Pi(\widetilde{\mathfrak{X}}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})). \tag{9.24}$$

Here Π is the harmonic projection in (6.14).

We put $b = \sum_{i=1}^n x_i\mathbf{e}_i$ and $y_i = T^{-u_i}e^{x_i}$. We then regard $\widetilde{\mathfrak{X}}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})$, $\mathfrak{X}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})$ as differential form valued formal functions of y_i. More precisely, we have:

Lemma 9.30. $\widetilde{\mathfrak{X}}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})$, $\mathfrak{X}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})$ *are contained in* $\overset{\circ}{\Lambda}{}_0^P\langle\langle y,y^{-1}\rangle\rangle\,\widehat{\otimes}\,\Omega(L(u))$, $H(L(u);\overset{\circ}{\Lambda}{}_0^P\langle\langle y,y^{-1}\rangle\rangle)$, *respectively. They are contained in* $\Lambda_0^P\langle\langle y,y^{-1}\rangle\rangle\,\widehat{\otimes}\,\Omega(L(u))$, $H(L(u);\Lambda_0^P\langle\langle y,y^{-1}\rangle\rangle)$ *if* $\mathfrak{b}\in\mathcal{A}(\Lambda_+)$.

The proof is the same as the proof of Proposition 7.10 and is omitted.

Lemma 9.31.

$$\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_a)\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_b) - \sum_c c^c_{ab}\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_c) = -\,\delta^{\mathfrak{b},b}_{\mathrm{can}}(\mathfrak{X}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})). \tag{9.25}$$

Proof. Using Lemma 9.7, we study the boundary of $\mathcal{M}_1(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;R)$, when its virtual dimension is n. We then use Stokes' theorem (Lemma 12.13 [FOOO5]) to show that the sum vanishes. Then we put a weighted sum as in (9.23), which will give rise to the equality (9.25). The details are in order.

There are three types of components, (1), (2) and (3) in Lemma 9.7. The case (1) that is the space $\mathcal{M}_1(\alpha;\beta;\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2;\partial R)$ becomes the left hand side of Lemma 9.31 after taking integration along the fiber and the weighted sum as in (9.23). This follows from (9.22), Lemmata 9.8, 9.22 and the formula

$$\overline{\mathbf{f}}_a\cup^{\mathfrak{b}}\overline{\mathbf{f}}_b = \sum_{\alpha,\ell,i,j} g^{ij}\frac{T^{\alpha\cap\omega/2\pi}}{\ell!}\exp(\mathfrak{b}_2\cap\alpha)\,\#\,\Big(\mathcal{M}_{\ell+3}(\alpha;\overline{\mathbf{f}}_a\otimes\overline{\mathbf{f}}_b\otimes\overline{\mathbf{f}}_i\otimes\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})\Big)\,\overline{\mathbf{f}}_j.$$

Note those terms appearing for (1) are already harmonic because of its T^n-invariance. So the projection $\Pi^{\mathrm{can}}_{\mathfrak{b},b}$ does not change it.

We next study the other cases, (2), (3) of Lemma 9.7.

For the case $K = R$, the boundary of type (2) becomes

$$\mathfrak{Y}(\alpha,\beta',\beta'',\ell_1,\ell_2',\ell_2'') := \mathcal{M}_1(\alpha;\beta';\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2';R)_{\mathrm{ev}_0}\times_{\mathrm{ev}_1}\mathcal{M}_2(\beta'';\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_2''}). \tag{9.26}$$

We put a (single) multisection on $\mathcal{M}_2(\beta''; \mathfrak{b}_{\rm high}^{\otimes \ell_2''})$ which is T^n-invariant. (Namely the '$\mathfrak{q}$' multisection.) By definition, Theorem 6.18 and Formula (6.9), we have

$$\begin{aligned}\sum_{\alpha,\beta',\beta''} \sum_{\ell_1,\ell_2',\ell_2''} & \frac{T^{(\alpha\#\beta'\#\beta'')\cap\omega/2\pi}}{\ell_1!\ell_2'!\ell_2''!} \rho^b(\partial(\beta'\#\beta''))\exp(\mathfrak{b}_2 \cap (\alpha\#\beta'\#\beta''))\\ & \qquad\qquad \times ({\rm ev}_0)_*(\mathfrak{Y}(\alpha,\beta',\beta'',\ell_1,\ell_2',\ell_2''))\\ &= \delta_{\mathfrak{b},b}(\widetilde{\mathfrak{X}}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b}))\\ &= \delta_{\mathfrak{b},b}^{\rm can}(\mathfrak{X}(\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;R;\mathfrak{b})).\end{aligned}$$

Thus this gives the right hand side of (9.25).

We finally study the boundary of type (3). In the present case this becomes

$$\mathcal{M}^{\rm main}_{1;\ell_2'+1}(\beta';\mathfrak{b}_{\rm high}^{\otimes\ell_2'}) \,{}_{{\rm ev}_0}\times_{{\rm ev}_i} \mathcal{M}_2(\alpha;\beta'';\overline{\mathbf{f}}_a,\overline{\mathbf{f}}_b;\ell_1,\ell_2'';R).$$

We remark that $\mathcal{M}^{\rm main}_{1;\ell_2'+1}(\beta';\mathfrak{b}_{\rm high}^{\otimes\ell_2'})$ is T^n-invariant and we have chosen a T^n-invariant multisection ('$\mathfrak{q}$' multisection) on it. Therefore

$$({\rm ev}_0)_*(\mathcal{M}^{\rm main}_{1;\ell_2'+1}(\beta';\mathfrak{b}_{\rm high}^{\otimes\ell_2'}))$$

is a T^n-invariant differential 0-form on $L(u)$, which is nothing but a constant function. Using Condition 9.26 (3), this implies that the contribution of the boundary of type (3) vanishes.

The proof of Lemma 9.31 is now complete. ☐

Now, Lemma 9.31 and Proposition 7.14 imply

$$\begin{aligned}&\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_a)\widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_b) - \sum_c c_{ab}^c \widetilde{\mathfrak{ks}}_{\mathfrak{b}}(\overline{\mathbf{f}}_c)\\ &\in {\rm Clos}_{d_{\overset{\circ}{P}}}\left(y_i\frac{\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1,\dots,n\right) \subset H(T^n;\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle).\end{aligned}$$

This implies Theorem 9.1. ☐

10. SURJECTIVITY OF KODAIRA-SPENCER MAP

In this section, we will prove the following theorem.

Theorem 10.1. *The Kodaira-Spencer map* $\mathfrak{ks}_{\mathfrak{b}} : H(X;\Lambda_0) \to {\rm Jac}(\mathfrak{PO}_{\mathfrak{b}})$ *is surjective.*

We begin with the corresponding result for the $\mathbb{C}$-reduction of $\mathfrak{ks}_{\mathfrak{b}}$, which is Proposition 10.2 below.

We have the isomorphism

$$H(X;\mathbb{C}) \cong H(X;\Lambda_0)/H(X;\Lambda_+).$$

We denote

$$\overline{\rm Jac}(\mathfrak{PO}_{\mathfrak{b}}) = \frac{{\rm Jac}(\mathfrak{PO}_{\mathfrak{b}})}{\Lambda_+{\rm Jac}(\mathfrak{PO}_{\mathfrak{b}})}.$$

Then $\mathfrak{ks}_{\mathfrak{b}}$ induces a map

$$\overline{\mathfrak{ks}}_{\mathfrak{b}} : H(X;\mathbb{C}) \to \overline{\rm Jac}(\mathfrak{PO}_{\mathfrak{b}}), \tag{10.1}$$

which is a ring homomorphism by Theorem 9.1. (Here $H(X;\mathbb{C})$ is a ring by the classical cup product without quantum corrections.)

Proposition 10.2. (1) *The map $\overline{\mathfrak{ks}}_{\mathfrak b}$ is surjective.*

(2) *If $\mathfrak b \in \mathcal A(\Lambda_+)$, then it is an isomorphism.*

Remark 10.3. Actually (2) holds without the assumption $\mathfrak b \in \mathcal A(\Lambda_+)$. The proof of this will be completed in Section 14.

Definition 10.4. We denote by G_0 the discrete submonoid generated by

$$\{\alpha \cap \omega/2\pi \mid \mathcal M(\alpha) \neq \emptyset\}.$$

For $\mathfrak b \in \mathcal A(\Lambda_0)$, we denote by $G(\mathfrak b)$ the smallest discrete monoid containing G_0 such that $\mathfrak b$ is $G(\mathfrak b)$-gapped in the sense of Definition 1.7.

Proof of Proposition 10.2. By Lemma 3.9 [FOOO5], the elements z_j $(j = 1, \dots, m)$ mod Λ_+ generate $\overline{\mathrm{Jac}}(\mathfrak{PO}_{\mathfrak b})$.

Let $\mathfrak b = \sum_{a=1}^B \mathfrak b_a \mathbf f_a$, $\mathfrak b_a \equiv \mathfrak b_{a,0} \mod \Lambda_+$ with $\mathfrak b_{a,0} \in \mathbb C$.

Lemma 10.5. *There exists $\lambda_1 > 0$ such that*

$$\mathfrak{PO}_{\mathfrak b} \equiv \sum_{j=1}^m e^{\mathfrak b_{j,0}} z_j + \sum_j \frac{1}{\ell_j!} \mathfrak b_{a_1^j,0} \cdots \mathfrak b_{a_{\ell_j}^j,0} z^{\beta_j} (\mathrm{ev}_0)_*(\mathcal M_{1;\ell_j}(\beta_a; \mathbf f_j)) \mod T^{\lambda_1}.$$

Notations appearing in the second term is as follows.

$$\mathbf f_j = \mathbf f_{a_1^j} \otimes \cdots \otimes \mathbf f_{a_{\ell_j}^j},$$

and

$$\mathbf f_{a_i^j} = \mathbf f_{a_i^j(1)} \cup \cdots \cup \mathbf f_{a_i^j(d_{i,j})}$$

with $a_i^j(j) \in \{1, \dots, m\}$, $2d_{i,j} = \deg \mathbf f_{a_i^j}$. We put $d_j = d_{1,j} + \cdots + d_{\ell_j,j}$ and

$$\beta_j = \beta_{c_j(1)} + \cdots + \beta_{c_j(d_j)} \qquad z^{\beta_j} = z_{c_j(1)} \cdots z_{c_j(d_j)}.$$

Also

$$a_1^j = \cdots = a_{\ell_j}^j \in \{1, \dots, m\}$$

never occurs. (This case corresponds to the first term.)

Proof. Let λ_1 be the smallest nonzero element of $G(\mathfrak b)$. We consider $\beta \in H_2(X; L(u); \mathbb Z)$, $\beta \neq 0$ with $\mathcal M_{1;\ell}(\beta) \neq \emptyset$. By Theorem 10.1 [FOOO4], we find

$$\beta = \sum_{j=1}^m k_j \beta_j + \alpha$$

$k_j \geq 0$, $\sum k_j > 0$ and α is realized by a sum of holomorphic spheres. The term induced by $\mathcal M_{1;\ell}(\beta)$ is

$$cT^{\beta \cap \omega/2\pi} y(u)_1^{k'_1} \cdots y(u)_n^{k'_n} = cT^{\alpha \cap \omega/2\pi} z_1^{k_1} \cdots z_m^{k_m},$$

where $c \in \mathbb C$. Therefore it suffices to consider the case $\alpha = 0$. (In fact otherwise $\alpha \cap \omega/2\pi \geq \lambda_1$.) In this case the Maslov index of β is $2\sum_{j=1}^m k_j$. By dimension counting we have $d_a = \sum_{j=1}^m k_j$. Therefore it suffices to consider the case $\beta = \beta_{a(1)} + \cdots + \beta_{a(d)}$. In case $\beta = \beta_j$, we have

$$(\mathrm{ev}_0)_*(\mathcal M_{1;1}(\beta_j; \mathbf f_j)) = 1$$

by Theorem 11.1 [FOOO4]. The lemma follows from (2.10). □

We now prove Proposition 10.2 (2). In case $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, Lemma 10.5 implies

$$\mathfrak{PO}_{\mathfrak{b}} \equiv z_1 + \cdots + z_m \mod \Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle.$$

Therefore we have

$$y_i \frac{\partial \mathfrak{PO}_\beta}{\partial y_i} \equiv \sum_{j=1}^m v_{j,i} z_j \mod \Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle. \tag{10.2}$$

Here $v_{j,i} \in \mathbb{Z}$ is defined by $d\ell_j = (v_{j,1}, \dots, v_{j,n})$.

On the other hand, it is easy to see that z_j satisfies the Stanley-Reisner relation modulo $\Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle$. Furthermore

$$\overline{\mathfrak{ks}}_{\mathfrak{b}}(\mathbf{f}_j) \equiv z_j \mod \Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle.$$

Now it is a consequence of classical result of Stanley that $\overline{\mathfrak{ks}}_{\mathfrak{b}}$ is an isomorphism. (See [Ful] p106 Proposition, for example.)

Next we prove Proposition 10.2 (1). Actually, we will prove a slightly sharper result. We first recall the following:

Convention 10.6. For $R \in \Lambda_0[[Z_1, \dots, Z_m]]$, we substitute z_j into Z_j and denote

$$R(z) = R(z_1, \dots, z_m).$$

This is well-defined by Lemma 2.3.

Proposition 10.7. *Let* $R \in \overset{\circ}{\Lambda}{}_0^P\langle\langle y, y^{-1}\rangle\rangle$. *Then there exist* $\rho \in H(X; \Lambda_0)$ *and* $U \in \overset{\circ}{\Lambda}{}_0^P\langle\langle y, y^{-1}\rangle\rangle$ *such that*

$$R - T^{\lambda_1} U = \mathfrak{ks}_{\mathfrak{b}}(\rho)$$

in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$.

Proof. By Lemma 10.5 we have

$$\mathfrak{ks}_{\mathfrak{b}}(\mathbf{f}_j) \equiv c_j z_j + P_j(z) \mod T^{\lambda_1} \overset{\circ}{\Lambda}{}_0^P\langle\langle y, y^{-1}\rangle\rangle \tag{10.3}$$

for $j = 1, \dots, m$, $c_j \in \mathbb{C} \setminus \{0\}$ and P_j are formal power series of Z_j with Λ_0 coefficient, such that each term of it has degree ≥ 2. (In fact, the second term of Lemma 10.5 has degree ≥ 2 in z_i.)

Using the fact that $\mathfrak{ks}_{\mathfrak{b}}$ is a ring homomorphism and that $\deg \mathbf{f}_j = 2$ and $\prod \mathbf{f}_j^{r_j} = 0$ (where $\prod$ is the ordinary cup product) whenever $\sum r_j \geq \dim X/2 + 1$, we can inductively apply (10.3) to find $\rho \in H(X; \Lambda_0)$, $R' \in \Lambda_0[[Z_1, \dots, Z_m]]$ such that

$$R(z) - \mathfrak{ks}_{\mathfrak{b}}(\rho) = R'(z) + T^{\lambda_1} U_1 \tag{10.4}$$

in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$, $U_1 \in \overset{\circ}{\Lambda}{}_0^P\langle\langle y, y^{-1}\rangle\rangle$ and the degree of each term of R' is greater than $\dim X/2$.

Lemma 10.8. *If the degree of each monomial in* $R' \in \Lambda_0[[Z_1, \dots, Z_m]]$ *is greater than* $\dim X/2$, *then there exists* $U_2 \in \overset{\circ}{\Lambda}{}_0^P\langle\langle y, y^{-1}\rangle\rangle$ *such that the following equality holds in* $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$.

$$R'(z) = T^{\lambda_1} U_2.$$

Proof. We put

$$c_N = \left[\frac{2N}{\dim X + 1}\right]$$

where $[a]$ is the greatest integer less than equal to a. We note that $c_N \to \infty$ as $N \to \infty$.

Sublemma 10.9. *Let N be a positive integer with $N > \dim X/2$. If the degree of each term of $R(N) \in \Lambda_0[[Z_1, \dots, Z_m]]$ is greater than N, then there exist $R(N+1) \in \Lambda_0[[Z_1, \dots, Z_m]]$ and $U(N) \in \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$ such that the degree of each term of $R(N+1)$ is greater than $N+1$ and that the equality*

$$R(N+1)(z) - R(N)(z) = T^{\lambda_1 c_N} U(N)$$

holds in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$.

Proof. We may assume that $R(N)$ is a monomial R_0 of degree N. We can then write

$$R_0 = R_1 R_2 \cdots R_{c_N}$$

so that each of R_i is a monomial of degree $> \dim X/2$.

Then applying (10.3) for each $i = 1, \dots, c_N$, we obtain a polynomial $\mathcal{R}_i(z)$ such that each monomial appearing in $\mathcal{R}_i(z)$ is of degree greater than that of $R_i(z)$ and satisfies

$$\mathfrak{ks}_{\mathfrak b}(R_i(\mathbf{f}_1/c_1, \dots, \mathbf{f}_m/c_m)) - R_i(z) = \mathcal{R}_i(z) + T^{\lambda_1} V_i$$

in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$. Since the ordinary cup product of $1 + (\dim X/2)$ elements vanishes on $H(X;\mathbb{C})$, there exists a polynomial $Q_i(z)$ such that

$$\mathfrak{ks}_{\mathfrak b}(R_i(\mathbf{f}_1/c_1, \dots, \mathbf{f}_m/c_m)) = T^{\lambda_1} Q_i(z).$$

We now set

$$R(N+1)(z) = -\prod_{i=1}^{c_N} (R_i(z) + \mathcal{R}_i(z)) + R(N)(z).$$

Then we find that the degree of each monomial in $R(N+1)(z)$ is bigger than that of $R(N)(z)$ and that

$$R(N+1)(z) - R(N)(z) = -\prod_{i=1}^{c_N} (R_i(z) + \mathcal{R}_i(z)) = -T^{c_N \lambda_1} \prod_{i=1}^{c_N} (V_i(z) + Q_i(z)).$$

By setting $U(N) = -\prod_{i=1}^{c_N} (V_i(z) + Q_i(z))$, we have finished the proof. □

We apply Sublemma 10.9 inductively by putting $R' = R(N_0)$. ($N_0 = \dim X/2 + 1$.) We have

$$R(N)(z) = R(N_0)(z) - \sum_{k=N_0}^{N-1} T^{\lambda_1 c_k} U(k).$$

Furthermore

$$\lim_{N \to \infty} \sum_{k=N_0}^{N} T^{\lambda_1 c_k} U(k)$$

converges. Moreover $R(N)(z)$ converges to 0 since $R(N)$ converges to zero in the formal power series ring. Therefore $R'(z) = R(N_0)(z)$ can be divided by T^{λ_1} in $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak b})$. This finishes the proof of Lemma 10.8. □

We have finished the proof of Proposition 10.7 by applying Lemma 10.8 to (10.4). □

Proposition 10.2 (1) is an immediate consequence of Proposition 10.7. □

Now we are ready to complete the proof of Theorem 10.1.

Proof. Let $R \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$. We use Proposition 10.7 to find $\rho_1 \in H(X;\Lambda_0)$ and R_1 such that $R - \mathfrak{ks}_{\mathfrak{b}}(\rho_1) = T^{\lambda_1}R_1$. We find inductively $\rho_k \in H(X;\Lambda_0)$ and R_k such that $R_{k-1} - \mathfrak{ks}_{\mathfrak{b}}(\rho_k) = T^{\lambda_1}R_k$. Clearly $\rho = \sum_{k=1}^{\infty} T^{\lambda_1(k-1)}\rho_k$ converges and $R = \mathfrak{ks}_{\mathfrak{b}}(\rho)$. □

11. VERSALITY OF THE POTENTIAL FUNCTION WITH BULK

Let z_j $(j = 1, \dots, m)$ be as in Definition 2.2.

Theorem 11.1. *Let $\mathfrak{P} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$. We assume that $\mathfrak{P}$ is G-gapped for a discrete monoid G containing G_0.*

1. *If $\mathfrak{P} \equiv z_1 + \cdots + z_m \mod \Lambda_+^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$, then there exists a G-gapped element $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$ and a G-gapped strict coordinate change y' on* $\operatorname{Int} P$ *such that*
$$\mathfrak{P}(y') = \mathfrak{PO}_{\mathfrak{b}}(y).$$
2. *If $\mathfrak{P} \equiv c_1 z_1 + \cdots + c_m z_m \mod \Lambda_+^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ with $c_i \in \mathbb{C} \setminus \{0\}$, then there exist a G-gapped element $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ and a G-gapped coordinate change y' on* $\operatorname{Int} P$ *such that*
$$\mathfrak{P}(y') = \mathfrak{PO}_{\mathfrak{b}}(y).$$

If $\mathfrak{P} \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$, then y' can be taken to be a strict G-gapped coordinate change on P.

(Note $\Lambda_+^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ is the ideal of $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ which is defined in Definition 2.4 (3).)

Proof. We first prove (1). We focus on the case of $\mathfrak{P} \in \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ since the case of $\mathfrak{P} \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ is similar. We enumerate the elements of G and write $G = \{\lambda_0, \lambda_1, \dots\}$ so that $\lambda_0 = 0$, $\lambda_i < \lambda_{i+1}$. We note that $\lambda_i \to \infty$ as $i \to \infty$.

We will construct a cycle $\mathfrak{b}(k) \in \mathcal{A}(\Lambda_0)$ and a coordinate change $y'(k)$ inductively over k with the following properties:

1. They are G-gapped.
2. $y'(k)$ is a G-gapped and strict coordinate change on $\operatorname{Int} P$.
3. $y'(k)$ satisfies
$$\mathfrak{P}(y'(k)) - \mathfrak{PO}_{\mathfrak{b}(k)}(y) \equiv 0 \mod T^{\lambda_{k+1}}\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle.$$
4. $(y'(k)_i - y'(k-1)_i)/y'(k-1)_i \equiv 0 \mod T^{\lambda_{k+1}}\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.

When $k = 0$, we set $\mathfrak{b}(0) = 0$ and $y'(0) = y$. Then the hypothesis $\mathfrak{B} \equiv z_1 + \cdots + z_m \mod \Lambda_+^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$ implies the above properties for $k = 0$.

Now suppose we have constructed $\mathfrak{b}(k)$ and $y'(k)$ so that

$$\mathfrak{P}(y'(k)) - \mathfrak{PO}_{\mathfrak{b}(k)}(y) \equiv T^{\lambda_{k+1}}\mathcal{P} \mod T^{\lambda_{k+2}}\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$$

holds. By Proposition 10.2 we may take $\mathfrak{z}_{k+1} \in H(X;\mathbb{C})$, $W(k,i) \in \mathbb{C}[[Z_1,\dots,Z_m]]$ $(i = 1,\dots,n)$ so that they satisfy

$$\mathcal{P} - \mathfrak{ks}_{\mathfrak{b}(k)}(\mathfrak{z}_{k+1}) \equiv \sum_{i=1}^{n} W(k,i)(z) y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}(k)}}{\partial y_i} \mod T^{\lambda_1}\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle.$$

(Here we use the fact that everything is G-gapped.) We define

$$\mathfrak{b}(k+1) = \mathfrak{b}(k) + T^{\lambda_{k+1}}\mathfrak{z}_{k+1},$$

and

$$y'(k+1)_i = y'(k)_i - T^{\lambda_{k+1}} W(k,i)(z) y_i, \quad i = 1,\dots,n.$$

It is easy to check Conditions (1) - (4) above for these choices. By construction, the limits $\mathfrak{b} = \lim_{k\to\infty}\mathfrak{b}(k)$, $y' = \lim_{k\to\infty} y'(k)$ exist and satisfy the properties required in the statement (1).

We next prove (2). After applying an explicit coordinate change $\mathfrak{b}(0) = \sum(\log c_i)\mathbf{f}_i$, we have

$$\mathfrak{B}(y'(0)) = \mathfrak{PO}_{\mathfrak{b}(0)}(y) \mod T^{\lambda_1}\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle.$$

Once we have arranged this, we can adapt the same inductive argument as for the case (1). □

Remark 11.2. We do not attempt to remove the gapped-ness from the hypotheses of Theorem 11.1 since non-gapped elements of $\Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle$ do not seem to appear in geometry.

12. ALGEBRAIZATION OF JACOBIAN RING

In this section we prove that there exists a coordinate change transforming the potential function to a Laurent *polynomial*.

Definition 12.1. We denote

$$\Lambda_0^P[y,y^{-1}] = \Lambda[y,y^{-1}] \cap \Lambda_0^P\langle\!\langle y, y^{-1}\rangle\!\rangle = \{w \in \Lambda[y,y^{-1}] \mid \mathfrak{v}_T^P(w) \geq 0\}.$$

We note that $\Lambda_0^P[y,y^{-1}]$ coincides with

$$\Lambda[y,y^{-1}] \cap \Lambda_0^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle.$$

In other words, $\Lambda_0^P[y,y^{-1}]$ is the image of $\Lambda_0[Z_1,\dots,Z_m]$ by the homomorphism in Lemma 2.3.

We also emphasize that $\Lambda_0^P[y,y^{-1}] \neq \Lambda_0[y,y^{-1}]$. For $\epsilon > 0$ we put

$$P_\epsilon = \{u \in \mathbb{R}^n \mid \ell_j(u) \geq \epsilon,\ j = 1,\dots,m\}.$$

See Section 4. (Note $P = \{u \in \mathbb{R}^n \mid \ell_j(u) \geq 0,\ j = 1,\dots,m\}$.)

Theorem 12.2. (1) *If $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$, then for each $\epsilon > 0$, there exists a strict coordinate change y' on P_ϵ such that*

$$\mathfrak{PO}_{\mathfrak{b}}(y') \in \Lambda_0^{P_\epsilon}[y,y^{-1}]. \tag{12.1}$$

(2) *If $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, then y' can be taken to be a strict coordinate change on P and*

$$\mathfrak{PO}_{\mathfrak{b}}(y') \in \Lambda_0^P[y,y^{-1}].$$

If $\mathfrak{b}$ is G-gapped, so is y'.

Proof. We first prove (2). Let $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$ which is G-gapped for a discrete submonoid $G \supseteq G_0$. We put $G = \{\lambda_0, \lambda_1, \dots\}$ as before. We first consider a polynomial

$$\mathfrak{P}_0 = z_1 + \cdots + z_m.$$

This satisfies $\mathfrak{PO}_{\mathfrak{b}} \equiv \mathfrak{P}_0 \mod T^{\lambda_1}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ for $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$. We put

$$\overline{\mathrm{Jac}}(\mathfrak{P}_0) = \frac{\mathrm{Jac}(\mathfrak{P}_0)}{\Lambda_+\mathrm{Jac}(\mathfrak{P}_0)}.$$

Since the property $\mathfrak{PO}_{\mathfrak{b}} \equiv \mathfrak{P}_0 \mod T^{\lambda_1}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ implies the equality of ideals in $\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$

$$\Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle + \left(y_i\frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}; i = 1, \dots, n\right) = \Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle + \left(y_i\frac{\partial\mathfrak{P}_0}{\partial y_i}; i = 1, \dots, n\right),$$

we find

$$\overline{\mathrm{Jac}}(\mathfrak{PO}_{\mathfrak{b}}) \cong \overline{\mathrm{Jac}}(\mathfrak{P}_0).$$

(However, there is no canonical map between $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ and $\mathrm{Jac}(\mathfrak{P}_0)$.) Through this isomorphism, we have a ring isomorphism which we also denote by $\overline{\mathfrak{ks}}$:

$$\overline{\mathfrak{ks}} \ : \ H(X;\mathbb{C}) \to \overline{\mathrm{Jac}}(\mathfrak{PO}_{\mathfrak{b}}) \cong \overline{\mathrm{Jac}}(\mathfrak{P}_0)$$

which sends $\bar{\mathbf{f}}_i$ to $\overline{z}_i$. Then we have

Lemma 12.3. *The polynomials of degree* $\le \dim X/2$ *of* $\overline{z}_1, \dots, \overline{z}_m$ *generate* $\overline{\mathrm{Jac}}(\mathfrak{P}_0)$.

We now inductively construct a coordinate change $y'(k)$ and a polynomial $B_k \in \Lambda_0[Z_1, \dots, Z_m]$ for $k = 0, \dots$ with the following properties:

(1) B_k is a polynomial of degree $\le \dim X/2$. It is G-gapped.

(2) $y'(k)$ is a G-gapped strict coordinate change on P.

(3)
$$\mathfrak{PO}_{\mathfrak{b}}(y'(k)) - (\mathfrak{P}_0 + B_k(z)) \equiv 0 \mod T^{\lambda_{k+1}}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle.$$

(4)
$$\frac{y'(k+1)_i - y'(k)_i}{y_i} \equiv 0 \mod T^{\lambda_{k+1}}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle.$$

(5)
$$B_{k+1} - B_k \equiv 0 \mod T^{\lambda_{k+1}}\Lambda_0[Z_1, \dots, Z_m].$$

Theorem 11.1 enables us to choose $B_0 = 0$. Now suppose we have constructed B_ℓ and $y'(\ell)$ for $\ell = k$. Then

$$\mathfrak{PO}_{\mathfrak{b}}(y'(k)) - (\mathfrak{P}_0 + B_k(z)) \equiv T^{\lambda_{k+1}}\mathcal{P}_k(z) \mod T^{\lambda_{k+2}}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$$

for some $\mathcal{P}_k \in \mathbb{C}[Z_1, \dots, Z_m]$. (Here Lemmas 2.6 and 2.3 show that we can take $\mathcal{P}_k$ as a polynomial.) By Lemma 12.3 there exist polynomials $\Delta B_{k+1} \in \mathbb{C}[Z_1, \dots, Z_m]$ of degree $\le \dim X/2$ and $W_{k+1,i} \in \mathbb{C}[Z_1, \dots, Z_m]$ such that

$$\mathcal{P}_k \equiv \Delta B_{k+1}(z) + \sum_{i=1}^n W_{k+1,i}(z) y_i \frac{\partial\mathfrak{P}_0}{\partial y_i} \mod T^{\lambda_1}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle.$$

We put

$$y'(k+1)_i = y'(k)_i - T^{\lambda_{k+1}}W_{k+1,i}(z)y_i, \qquad B_{k+1} = B_k + T^{\lambda_{k+1}}\Delta B_{k+1}.$$

It is easy to see that they have the required properties and hence we finish the construction of B_k and $y'(k)$ for $k \in \mathbb{Z}_{\ge 0}$.

Now we consider the limits $B = \lim_{k\to\infty} B_k(z)$ and $y'_\infty = \lim_{k\to\infty} y'(k)$ which converge by construction. We also find that $y'_\infty \in \Lambda^P\langle\langle y, y^{-1}\rangle\rangle$.

If we put $\mathfrak{B} = B(z) \in \Lambda_0^P[y, y^{-1}]$, the equality $\mathfrak{PO}_{\mathfrak b}(y'_\infty) = \mathfrak{P}_0 + \mathfrak{B}$ follows. We have thus proved (2).

We next prove (1). By Lemma 10.5, we may put

$$\mathfrak{PO}_{\mathfrak b} \equiv c_1 z_1 + \cdots + c_m z_m + R(z_1, \dots, z_m) \mod \Lambda_+^P[y, y^{-1}] \tag{12.2}$$

where $R \in \Lambda_0[[Z_1, \dots, Z_m]]$ each summand of which has degree ≥ 2, and $c_i \in \mathbb{C}\backslash\{0\}$. We put

$$z_i^\epsilon = T^{-\epsilon} z_i.$$

Note that

$$\mathfrak{v}_T^{P_\epsilon}(z_i^\epsilon) = \inf_{u\in P_\epsilon} \mathfrak{v}_T^u(z_i^\epsilon) = 0.$$

In fact, z_i^ϵ plays a role of z_i for our polytope P_ϵ. There exists a surjective homomorphism

$$\Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle \to \Lambda_0^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle$$

which sends Z_i to z_i^ϵ. We consider $T^{-\epsilon}\mathfrak{PO}_{\mathfrak b} \in \Lambda_0^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle$. From (12.2) it is easy to see that

$$T^{-\epsilon}\mathfrak{PO}_{\mathfrak b} \equiv c_1 z_1^\epsilon + \cdots + c_m z_m^\epsilon \mod \Lambda_+^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle. \tag{12.3}$$

Then we can repeat the proof of (2) and obtained the conclusion. □

Remark 12.4. Theorem 12.2 should be related to the classical results in the deformation theory of isolated singularity. See [Ar], [El].

We next discuss the relationships between several variations of the Jacobian rings.

Condition 12.5. Let $\mathfrak{P} \in \Lambda_0^P[y, y^{-1}]$. We assume that $\mathfrak{P} - \sum_{i=1}^m c_i z_i$ is contained in the ideal $\Lambda_+^P[y, y^{-1}]$.

Remark 12.6. For $\mathfrak b \in \mathcal{A}(\Lambda_+)$, $\mathfrak{PO}_{\mathfrak b}$ satisfies this condition. For $\mathfrak b \in \mathcal{A}(\Lambda_0)$, Formula (12.3) implies that the condition is satisfied for $T^{-\epsilon}\mathfrak{PO}_{\mathfrak b}$ if we replace P and z_i by P_ϵ and z_i^ϵ respectively.

Definition 12.7. Suppose that $\mathfrak{P}$ satisfies Condition 12.5. We put

$$\mathrm{PJaco}(\mathfrak{P}) = \frac{\Lambda_0^P[y, y^{-1}]}{\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)}, \qquad \mathrm{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}) = \frac{\Lambda_0^P[y, y^{-1}]}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)},$$

$$\mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P}) = \frac{\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle}{\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)}, \qquad \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}) = \frac{\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)},$$

$$\mathrm{Jaco}(\mathfrak{P}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle}{\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)}, \qquad \mathrm{PJac}(\mathfrak{P}) = \frac{\Lambda_0^P[y, y^{-1}]}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)}.$$

Here we recall that $\mathrm{Clos}_{\mathfrak{v}_T^P}$ means the closure with respect to the norm $\mathfrak{v}_T^P$ and $\mathrm{Clos}_{d_{\overset{\circ}{P}}}$ means the closure with respect to the metric $d_{\overset{\circ}{P}}$. (See Definition 4.2.) We also note that we already defined

$$\mathrm{Jac}(\mathfrak{P}) = \frac{\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i}\right)}$$

before in Definition 2.8.

In the above notations of various Jacobian rings, the letter 'P' in front of 'Jac' stands for 'polynomial', the letter 'o' after 'Jac' stands for 'taking the quotient *not* by the closure' and the subscript $\mathfrak{v}_T^P$ stands for 'using the topology with respect to the norm $\mathfrak{v}_T^P$'.

There are natural homomorphisms described in the following diagram.

$$\begin{tikzcd}
 & \mathrm{PJaco}(\mathfrak{P}) \arrow[d] \arrow[ddr] & & \\
 & \mathrm{PJac}(\mathfrak{P}) \arrow[rr] & & \mathrm{Jac}(\mathfrak{P}) \\
 & & \mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P}) \arrow[d] & \\
\mathrm{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}) \arrow[uur] \arrow[rr] & & \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}) \arrow[uur] &
\end{tikzcd} \tag{12.4}$$

Lemma 12.8. *Suppose $\mathfrak{P}$ satisfies Condition 12.5. Then we have the following:*

1. *The homomorphism* $\mathrm{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}) \to \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P})$ *is an isomorphism.*
2. *The homomorphism* $\mathrm{PJac}(\mathfrak{P}) \to \mathrm{Jac}(\mathfrak{P})$ *is an isomorphism.*
3. *The homomorphism* $\mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P}) \to \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P})$ *is an isomorphism if* $\mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P})$ *is torsion free.*
4. *The homomorphism* $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}) \to \mathrm{Jac}(\mathfrak{P})$ *is an isomorphism if* $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P})$ *is torsion free.*

Remark 12.9.

1. Theorem 2.32 implies that $\mathrm{Jac}(\mathfrak{P})$ is always torsion free. However, while we are proving Theorem 2.32, we need to check the assumption to use Lemma 12.8 (4).
2. Finally (at the end of Section 14) we have
$$\mathrm{PJac}(\mathfrak{P}) = \mathrm{Jac}(\mathfrak{P}) = \mathrm{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}) = \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}) = \mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P}).$$
On the other hand, $\mathrm{PJaco}(\mathfrak{P}) \neq \mathrm{PJac}(\mathfrak{P})$ in general. (See Example 8.2 [FOOO4].) We will not further study the relationship between $\mathrm{Jaco}(\mathfrak{P})$ and $\mathrm{Jac}(\mathfrak{P})$, since such a study will not be needed in this paper.

Proof of Lemma 12.8. We choose $G \supset G_0$ such that $\mathfrak{P}$ is G-gapped and set $G = \{\lambda_0, \lambda_1, \dots\}$.

We first prove (2). We remark that Lemma 12.3 holds for $\overline{\mathrm{Jac}}(\mathfrak{P})$ also. (We use Theorem 11.1 to prove it.) Moreover Sublemma 10.9 also holds for $\mathrm{Jac}(\mathfrak{P})$.

Injectivity of the map (2) is clear and so we now prove its surjectivity. Let $\mathfrak{B} \in \mathrm{Jac}(\mathfrak{P})$. By an induction over k, we will find $B_k \in \Lambda_0[Z_1, \dots, Z_n]$ such that

1. $\deg B_k \le \dim X/2$.
2. $\mathfrak{B} - B_k(z) \in T^{\lambda_k}\mathrm{Jac}(\mathfrak{P})$.
3. $B_{k+1} - B_k \in T^{\lambda_k}\Lambda_0[[Z_1, \dots, Z_n]]$.

Suppose we have constructed B_k and put

$$\mathfrak{B} - B_k(z) = T^{\lambda_k}F_k(z)$$

with $F_k \in \mathbb{C}[[Z_1, \dots, Z_n]]$. Let F_k' be the sum of the terms of F_k of degree $\le \dim X/2$. Then

$$F_k(z) - F_k'(z) \in T^{\lambda_1}\operatorname{Jac}(\mathfrak{P})$$

by Sublemma 10.9. It is easy to see that

$$B_{k+1} = B_k + T^{\lambda_k}F_k'$$

has the required properties.

Since $\deg B_k \le \dim X/2$ is bounded and $\lambda_k \to \infty$, B_k converges to a *polynomial* in $\Lambda_0[Z_1, \dots, Z_m]$ which we denote by B. Since $B_k(z)$ converges to $\mathfrak{B}$ in $\operatorname{Jac}(\mathfrak{P})$ by Condition (2), we have proved $B(z) = \mathfrak{B}$. By varying $\mathfrak{B}$ in $\operatorname{Jac}(\mathfrak{P})$, the surjectivity of $\operatorname{PJac}(\mathfrak{P}) \to \operatorname{Jac}(\mathfrak{P})$ follows.

The proof of (1) is easier and left for the reader.

We next prove (3). This time the surjectivity is obvious and so it suffices to show the injectivity. Let $x \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ be an element which projects down to zero in $\operatorname{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P})$, i.e., $x \in \operatorname{Clos}_{\mathfrak{v}_T^P}\left(y_i\frac{\partial\mathfrak{P}}{\partial y_i}\right)$. By definition of $\operatorname{Clos}_{\mathfrak{v}_T^P}\left(y_i\frac{\partial\mathfrak{P}}{\partial y_i}\right)$, there exists a sequence $x_\ell \in \left(y_i\frac{\partial\mathfrak{P}}{\partial y_i}\right)$ such that $x - x_\ell$ converges to zero in $\mathfrak{v}_T^P$ topology. Take a monoid $G \supseteq G_0$ such that $\mathfrak{P}$ and x are G-gapped and put $G = \{\lambda_0, \lambda_1, \dots\}$.

We will inductively construct $\mathfrak{Z}_{k,i} \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ which is G-gapped and satisfies

$$\begin{aligned} x - \sum_{i=1}^n \mathfrak{Z}_{k,i} y_i \frac{\partial\mathfrak{P}}{\partial y_i} &\equiv 0 \mod T^{\lambda_k}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle, \\ \mathfrak{Z}_{k+1,i} - \mathfrak{Z}_{k,i} &\equiv 0 \mod T^{\lambda_k}\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle. \end{aligned}$$

Suppose we have $\mathfrak{Z}_{k,i}$. We put

$$x - \sum_{i=1}^n \mathfrak{Z}_{k,i} y_i \frac{\partial\mathfrak{P}}{\partial y_i} = T^{\lambda_k}\mathfrak{Z}(k)$$

for some $\mathfrak{Z}(k) \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$. We take a sufficiently large ℓ so that

$$\mathfrak{v}_T^P(x_\ell - x) > \lambda_{k+1}.$$

Then there exists $\mathfrak{U}_{k+1} \in \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$ such that

$$(x_\ell - x) = T^{\lambda_{k+1}}\mathfrak{U}_{k+1}.$$

Then

$$x_\ell - \sum_{i=1}^n \mathfrak{Z}_{k,i} y_i \frac{\partial\mathfrak{P}}{\partial y_i} = T^{\lambda_k}(\mathfrak{Z}(k) + T^{\lambda_{k+1}-\lambda_k}\mathfrak{U}_{k+1}).$$

Since $x_\ell \in \left(y_i\frac{\partial\mathfrak{P}}{\partial y_i}\right)$, this implies that $T^{\lambda_k}(\mathfrak{Z}(k) + T^{\lambda_{k+1}-\lambda_k}\mathfrak{U}_{k+1})$ lies in $\left(y_i\frac{\partial\mathfrak{P}}{\partial y_i}\right)$ and hence is zero in $\operatorname{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P})$. By the hypothesis that $\operatorname{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P})$ is torsion free, we find that $\mathfrak{Z}(k) + T^{\lambda_{k+1}-\lambda_k}\mathfrak{U}_{k+1}$ is zero in $\operatorname{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P})$. Therefore there exists $Z(k,i) \in \Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle$ such that

$$\mathfrak{Z}(k) + T^{\lambda_{k+1}-\lambda_k}\mathfrak{U}_{k+1} \equiv \sum_{i=1}^n Z(k,i)(z) y_i \frac{\partial\mathfrak{P}}{\partial y_i} \mod T^{\lambda_{k+1}-\lambda_k}\Lambda_+^P\langle\langle y, y^{-1}\rangle\rangle.$$

Hence $\mathfrak{Z}_{k+1,i} := \mathfrak{Z}_{k,i} + T^{\lambda_k}Z(k,i)(z)$, has the required properties for $k+1$.

If we put $\lim_{k\to\infty} \mathfrak{z}_{k,i} = \mathfrak{z}_i$, then we have

$$x = \sum_{i=1}^{n} \mathfrak{z}_i y_i \frac{\partial \mathfrak{P}}{\partial y_i} \in \left(y_i \frac{\partial \mathfrak{P}}{\partial y_i} \right)$$

and hence $x = 0$ in $\mathrm{Jaco}_{\mathfrak{v}_T^P}(\mathfrak{P})$ as required.

Finally we prove (4). Surjectivity of the map immediately follows from (2) and so the injectivity is the only non-obvious part of the proof. By the torsion-freeness assumption of $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P})$, it suffices to show the injectivity after taking the tensor product $\otimes_{\Lambda_0}\Lambda$. Then by the versality given in Theorem 11.1, it is enough to consider the case of the potential function $\mathfrak{P} = \mathfrak{PO}_{\mathfrak{b}}$. For this case, we use the decomposition given Proposition 2.15 for $\mathfrak{PO}_{\mathfrak{b}}$: it is easy to see that $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})$ has the same kind of decomposition such as the one for $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$. Recall if $\mathfrak{y} \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$ and if we set $u := \mathfrak{v}_T(\mathfrak{y})$ u lies in $\mathrm{Int}\, P$. Therefore

$$\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y}) \cong \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y}). \tag{12.5}$$

Once we have this, we can obtain the factorization

$$\frac{\Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}\right)} \otimes_{\Lambda_0} \Lambda \cong \prod_{\mathfrak{y}\in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})} \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\mathfrak{y})$$

as for $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ in Proposition 2.15 in the same way. (Note we take the closure with respect to the norm $\mathfrak{v}_T^P$ in the left hand side of the factorization.) Therefore the injectivity of (4) follows because the last product is isomorphic to $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda$ itself. Now the proof of Lemma 12.8 is complete. □

13. SEIDEL HOMOMORPHISM AND A RESULT BY MCDUFF-TOLMAN

In this section, we use the results by Seidel [Se1] and McDuff-Tolman [MT] to prove Theorem 13.4. To describe the results of [MT], [Se1], we need some notations.

Let $\mathbf{f}_i$ $(i = 1, \dots, m)$ be the divisors which generate $\mathcal{A}^2$. We put

$$D_i = PD(\mathbf{f}_i) \in H_2(X;\mathbb{Z}).$$

Let $\mathcal{P} \subset \{1, \dots, m\}$ be a primitive collection. (See for example Definition 2.4 [FOOO4].) We have $\bigcap_{i\in\mathcal{P}} D_i = \emptyset$ and we have $\mathcal{P}' \subset \{1, \dots, m\} \setminus \mathcal{P}$ and k_i $(i \in \mathcal{P}')$ such that

$$\sum_{i\in\mathcal{P}} d\ell_i = \sum_{i'\in\mathcal{P}'} k_{i'} d\ell_{i'}, \qquad \bigcap_{i'\in\mathcal{P}'} D_{i'} \neq \emptyset.$$

$(k_i \in \mathbb{Z}_{\geq 0}.)$ We define

$$\omega(\mathcal{P}) = \sum_{i\in\mathcal{P}} \ell_i(u) - \sum_{i'\in\mathcal{P}'} k_{i'}\ell_{i'}(u).$$

Note this is independent of the choice of u by Lemma 6.2 [FOOO4]. We take m formal variables Z_i.

Definition 13.1. *The quantum Stanley-Reisner relation associated with* $\mathcal{P}$ *is*

$$\prod_{i\in\mathcal{P}} Z_i - T^{\omega(\mathcal{P})} \prod_{i'\in\mathcal{P}'} Z_{i'} = 0. \tag{13.1}$$

We denote the left hand side of (13.1) by $\mathfrak{I}(\mathcal{P})$. The *quantum Stanley-Reisner ideal* is an ideal of $\Lambda_0^P[Z_1,\dots,Z_m]$ generated by $\mathfrak{I}(\mathcal{P})$ for various $\mathcal{P}$. We denote it by SR_ω.

Theorem 13.2 (McDuff-Tolman [MT])**.** *There exists an element z_i' of $H(X;\Lambda_0)$ with the following properties:*

(1) $z_i' - D_i \equiv 0 \mod H(X;\Lambda_+)$.

(2) *z_i' satisfies the quantum Stanley-Reisner relation in the (small) quantum cohomology ring.*

This theorem is proved in Section 5.1 [MT]. There our z_i' is written as y_i.

Remark 13.3. We take the quantum cup product without bulk deformation in (2) above. We can prove a similar result for the quantum cup product with bulk by using Seidel homomorphism with bulk. We do not prove this here since we do not use it.

Let $d\ell_j = (v_{j,1},\dots,v_{j,n}) \in \mathbb{Z}^n$ for $j = 1,\dots,m$. We put

$$\mathfrak{P}_i^{(0)} = \sum_{j=1}^m v_{j,i} Z_j \in \Lambda_0[Z_1,\dots,Z_m], \qquad i = 1,\dots,n. \tag{13.2}$$

Theorem 13.4. *There exist $\mathfrak{P}_i \in \Lambda_0[Z_1,\dots,Z_m]$ $(i = 1,\dots,n)$ such that*

(1) $\mathfrak{P}_i(z_1',\dots,z_m') = 0$ *in $H(X;\Lambda_0)$. Here we use the quantum cup product with $\mathfrak{b} = \mathbf{0}$ in the left hand side.*

(2) *The ring homomorphism*

$$\psi_1 : \frac{\Lambda_0[Z_1,\dots,Z_m]}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(SR_\omega \cup \{\mathfrak{P}_i \mid i = 1,\dots,n\}\right)} \to QH(X;\Lambda_0)$$

which sends Z_j to z_j' is an isomorphism.

(3) $\mathfrak{P}_i \equiv \mathfrak{P}_i^{(0)} \mod \Lambda_+[Z_1,\dots,Z_m]$.

Remark 13.5. There is a similar discussion in Section 5.1 [MT]. We give the detail of its proof below for completeness' sake since in [MT] the process of taking the closure of the *ideal* is not explicitly discussed.

Proof. We begin with the following proposition.

Proposition 13.6. *There exists an isomorphism*

$$\psi_2 : \frac{\Lambda_0\langle\langle Z_1,\dots,Z_m\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^P}(SR_\omega)} \to \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$$

such that $\psi_2(Z_j) = z_j$.

Proof. By Lemma 2.3 we have a surjective and continuous homomorphism

$$\tilde\psi_2 : \Lambda_0\langle\langle Z_1,\dots,Z_m\rangle\rangle \to \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle$$

such that $\tilde\psi_2(Z_j) = z_j$. Since z_j satisfies the quantum Stanley-Reisner relation, $\tilde\psi_2$ induces a continuous and surjective homomorphism ψ_2. It suffices to show the injectivity of ψ_2. Let $Z^w = Z_1^{w_1}\cdots Z_m^{w_m}$ be a monomial with $w = (w_1,\dots,w_m) \in \mathbb{Z}^m_{\ge 0}$. We say Z^w is *in minimal expression* if $\mathfrak{v}_T^P(z^w) = 0$.

Lemma 13.7. *For any Z^w there exists a minimal expression $Z^{w'}$ and $\lambda \in \mathbb{R}$ such that*

(1) $Z^w \equiv T^{\lambda} Z^{w'} \mod SR_{\omega}$.
(2) $\lambda \geq 0$ *and the equality holds only when* $w' = w$.

Proof. See the proof of Proposition 6.9 [FOOO4]. $\square$

Now let $P \in \Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle$. Then by the definition of $\Lambda_0\langle\langle Z_1, \dots, Z_m\rangle\rangle$ which is the completion with respect to the norm $\mathfrak{v}_T$, we can write P as the sum $P = \sum_{j=1}^{\infty} a_j Z^{w_{(j)}}$ with $w_{(j)} = (w_{(j),1}, \dots, w_{(j),m})$ and $\lim \mathfrak{v}_T(a_j) = \infty$.

Lemma 13.8. *Suppose* $P = \sum_{j=1}^{\infty} a_j Z^{w_{(j)}}$ *is in the kernel of* ψ_2. *Choose* $w'_{(j)}$ *and* λ'_j *as in Lemma 13.7 so that* $P \equiv \sum_j a_j T^{\lambda'_j} Z^{w'_{(j)}} \mod SR_{\omega}$ *and that* $Z^{w'_{(j)}}$ *are in minimal expression. Then we have*

$$\sum_j a_j T^{\lambda'_j} Z^{w'_{(j)}} = 0.$$

Proof. Suppose to the contrary that we have $\sum_j a_j T^{\lambda'_j} Z^{w'_{(j)}} \neq 0$.

We can rewrite the sum $\sum_j a_j T^{\lambda'_j} Z^{w'_{(j)}}$ as $\sum_j b_j T^{\lambda'_j} Z^{w''_{(j)}}$ so that $w''_{(j)} \neq w''_{(j')}$ if $j \neq j'$. We put $c = \mathfrak{v}_T\left(\sum_j b_j T^{\lambda'_j} Z^{w''_{(j)}}\right)$. (Note that we use the norm $\mathfrak{v}_T$ in $\Lambda_0[[Z_1, \dots, Z_m]]$ here.) Take $J = \{j \mid \mathfrak{v}_T(b_j) + \lambda'_j = c\}$. Then J is nonempty by the condition (2) in Lemma 13.7. For $j \in J$, we write $Z^{w''_{(j)}} = \prod_{i=1}^m Z_i^{k_{i,j}}$. We put $\{i_{1,j}, \dots, i_{\ell(j),j}\} = \{i \mid k_{i,j} \neq 0\}$. By definition of the minimal expression, we find that $P_{I(j)} = \bigcap_{a=1}^{\ell(j)} P_{i_{a,j}}$ is nonempty. Let $u_j = \mathfrak{v}_T^{u_j}(z^{w''_{(j)}})$ be an interior point of $P_{I(j)}$. We recall $\mathfrak{v}_T^{u_j}(z^{w''_{(j)}}) = 0$ since $u_j \in P_{I(j)} \subset \partial P$.

We choose j so that $P_{I(j)}$ is minimal in that $P_{I(j)} \supseteq P_{I(j')}$ does not hold for any $j' \neq j$. Then, we have $\mathfrak{v}_T^{u_j}(z^{w''_{j'}}) > 0$ for any $j' \neq j$. In fact there is i such that $k_{i,j'} > 0$ and $k_{i,j} = 0$. Since $\mathfrak{v}_T^{u_j}(z_i) > 0$, it follows that $\mathfrak{v}_T^{u_j}(z^{w''_{(j')}}) > 0$.

Therefore $i_u\psi_2(\sum_j b_j T^{\lambda'_j} Z^{w''_{(j)}}) \equiv b_j z^{w''_{(j)}}$ modulo $T^c \Lambda_0^{u_j}\langle\langle y, y^{-1}\rangle\rangle$. Here

$$i_u : \Lambda_0^P\langle\langle y, y^{-1}\rangle\rangle \to \Lambda_0^{u_j}\langle\langle y, y^{-1}\rangle\rangle$$

is the obvious inclusion. This contradicts to the hypothesis $P \in \operatorname{Ker}\psi_2$. $\square$

The proof of Proposition 13.6 is now complete. $\square$

Construction of $\mathfrak{P}_i$ required in Theorem 13.4 will be finished by the following lemma.

Lemma 13.9. *Let* $G \supset G_0$ *such that* z'_j *are* G *gapped. Then there exist* $\mathfrak{P}_i^{(k)} \in \Lambda_0[Z_1, \dots, Z_m]$ *with the following properties:*

(1) $\deg \mathfrak{P}_i^{(k)} \leq \dim X/2$ *and* $\mathfrak{P}_i^{(k)}$ *is* G*-gapped.*
(2) $\mathfrak{P}_i^{(k+1)} \equiv \mathfrak{P}_i^{(k)} \mod T^{\lambda_{k+1}}\Lambda_0[Z_1, \dots, Z_m]$.
(3) $\mathfrak{P}_i^{(0)}$ *is as in (13.2).*
(4) $\mathfrak{P}_i^{(k)}(z'_1, \dots, z'_m) \equiv 0 \mod T^{\lambda_{k+1}} H(X; \Lambda_0)$.

Proof. Let $\mathfrak{P}_i^{(0)}$ be as in (13.2). Theorem 13.2 implies

$$\mathfrak{P}_i^{(0)}(z'_1, \dots, z'_m) = \sum_j v_{j,i} D_j \equiv 0 \mod T^{\lambda_1} H(X; \Lambda_0).$$

Hence we can start the induction. We put

$$\mathfrak{P}_i^{(k)}(z'_1,\dots,z'_m) \equiv T^{\lambda_{k+1}} c_i \mod T^{\lambda_{k+2}} H(X;\Lambda_0)$$

with $c_i \in H(X;\mathbb{C})$. Since D_i generate $H(X;\mathbb{C})$ as a ring, we have $P_{i,k} \in \mathbb{C}[Z_1,\dots,Z_m]$ such that

$$c_i = P_{i,k}(D_1,\dots,D_m).$$

We put $P_i^{(k+1)} = P_i^{(k)} - T^{\lambda_{k+1}} P_{i,k}$. It is easy to check that this has the required properties. □

We put $\mathfrak{P}_i = \lim_{k\to\infty} \mathfrak{P}_i^{(k)}$. Properties (1) and (3) are satisfied by Lemma 13.9. The proof that ψ_1 is an isomorphism is the same as the proof of Lemma 12.8 (2) and (3). (We remark that $H(X;\Lambda_0)$ is a free Λ_0 module. Therefore the assumption of Lemma 12.8 (3) is satisfied in our case.) Thus the next lemma complete the proof of Theorem 13.4. □

Lemma 13.10. *We have*

$$\frac{\Lambda_0[Z_1,\dots,Z_m]}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(SR_\omega \cup \{\mathfrak{P}_i \mid i=1,\dots,n\}\right)} \cong \frac{\Lambda_0[[Z_1,\dots,Z_m]]}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(SR_\omega \cup \{\mathfrak{P}_i \mid i=1,\dots,n\}\right)}.$$

Proof. We remark that $\psi_1(Z_i) = z'_i$. Since $z'_i \equiv PD([D_i]) \mod T^{\lambda_1}$ and quantum cohomology is the same as singular cohomology ring modulo T^{λ_1} we have

$$\psi_1(Z_1^{k_1}\cdots Z_m^{k_m}) \equiv 0 \mod T^{\lambda_1(k_1+\cdots+k_m-n)/n} H(X;\Lambda_0). \tag{13.3}$$

Using (13.3) in place of Sublemma 10.9 we can repeat the proof of Lemma 12.8 (2) to prove Lemma 13.10. □

14. Injectivity of Kodaira-Spencer map

14.1. **The case $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$.** In this subsection, we prove the injectivity for the case $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$. We discuss the case $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ in the next subsection.

Theorem 14.1. *If $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, then*

$$\mathfrak{ks}_{\mathfrak{b}} : H(X;\Lambda_0) \to \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$$

is injective.

The main part of the proof is the following proposition. We put

$$\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda = \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda,$$
$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda = \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda.$$

Proposition 14.2.

$$\dim_{\mathbb{C}} H(X;\mathbb{C}) \le \dim_\Lambda \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda.$$

Proposition 14.2 ⇒ Theorem 14.1. Let $\bar{\mathbf{f}}_i$ $(i=1,\dots,B')$ be the basis of $H(X;\Lambda_0)$. By Theorem 10.1, $\mathfrak{ks}_{\mathfrak{b}}(\bar{\mathbf{f}}_i)$ generates $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})$. Therefore by Proposition 14.2 they are linearly independent in $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda$. Thus

$$\mathfrak{ks}_{\mathfrak{b}}^P : H(X;\Lambda_0) \to \mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})$$

is injective. We can also prove that $\mathfrak{ks}_{\mathfrak{b}}^P$ is surjective in the same way as Theorem 10.1. It follows that $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}})$ is torsion free. Therefore $\mathrm{Jac}_{\mathfrak{v}_T^P}(\mathfrak{PO}_{\mathfrak{b}}) \cong \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ by Lemma 12.8 (4). Hence Theorem 14.1. □

The proof of Proposition 14.2 occupies the rest of this section. Let y' be as in Theorem 12.2 (2). We put

$$\mathcal{P} = \mathfrak{PO}_{\mathfrak{b}}(y') \in \Lambda_0^P[y, y^{-1}].$$

Since y' is strict and $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, we have

$$\mathcal{P} \equiv z_1 + \cdots + z_m \mod \Lambda_+^P[y, y^{-1}].$$

Let $\mathfrak{P}_i \in \Lambda_0[Z_1, \dots, Z_m]$ be as in Theorem 13.4. We have $\mathfrak{P}_i(z) \in \Lambda_0^P[y, y^{-1}]$.

Definition 14.3.

$$\mathfrak{B}_i = s\mathfrak{P}_i(z) + (1-s)y_i \frac{\partial \mathcal{P}}{\partial y_i} \in \Lambda_0^P[s, y, y^{-1}]. \tag{14.1}$$

$$\mathcal{R} = \frac{\Lambda[s, y, y^{-1}]}{(\mathfrak{B}_i : i = 1, \dots, n)}, \qquad \underline{\mathcal{R}} = \frac{\Lambda[s, y, y^{-1}]}{\operatorname{Clos}_{\mathfrak{v}_T^P}(\mathfrak{B}_i : i = 1, \dots, n)}. \tag{14.2}$$

There are morphisms of schemes over Λ:

$$Spec(\underline{\mathcal{R}}) \longrightarrow Spec(\mathcal{R}) \xrightarrow{\pi} Spec(\Lambda[s]) = \Lambda.$$

We denote $I = (\mathfrak{B}_1, \dots, \mathfrak{B}_n)$ and by $\overline{I}$ the closure of I with respect to $\mathfrak{v}_T^P$. We take irredundant primary decompositions of the ideals I and $\overline{I}$:

$$I = \bigcap_{i \in \mathfrak{I}_+} \mathfrak{q}_i, \qquad \overline{I} = \bigcap_{j \in \mathfrak{I}} \mathfrak{Q}_j. \tag{14.3}$$

Here an irredundant primary decomposition represents the given ideal as the intersection of primary ideals such that any $\mathfrak{q}_i$ (resp. $\mathfrak{Q}_j$) cannot be eliminated and the set of associated primes $\{\sqrt{\mathfrak{q}_i}\}$ (resp. $\{\sqrt{\mathfrak{Q}_j}\}$) consists of mutually distinct prime ideals. Note that, for an irredundant primary decomposition, primary ideals belonging to minimal associated primes are uniquely determined. We decompose $Spec(\mathcal{R})$ (resp. $Spec(\underline{\mathcal{R}})$) to irreducible components. Namely

$$|Spec(\mathcal{R})| = \bigcup_{i \in \mathfrak{I}_+} \left| Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{q}_i}\right) \right|, \qquad |Spec(\underline{\mathcal{R}})| = \bigcup_{j \in \mathfrak{I}} \left| Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{Q}_j}\right) \right|.$$

Here and hereafter in this section, we put $|Spec\, A|$ for the set of Λ-valued points of a Λ-scheme $Spec\, A$. We put

$$\mathfrak{I} = \left\{ i \in \mathfrak{I}_+ \;\middle|\; \left| Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{q}_i}\right) \right| \cap |Spec(\underline{\mathcal{R}})| \ne \emptyset \right\} \tag{14.4}$$

and

$$\mathfrak{X}_i = Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{q}_i}\right), \qquad \mathfrak{Y}_j = Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{Q}_j}\right).$$

Let $(\mathfrak{X}_i)_{\mathrm{red}}$ (resp. $(\mathfrak{Y}_j)_{\mathrm{red}}$) be the associated reduced schemes to $\mathfrak{X}_i$ (resp. $\mathfrak{Y}_j$). Namely

$$(\mathfrak{X}_i)_{\mathrm{red}} = Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\sqrt{\mathfrak{q}_i}}\right), \qquad (\mathfrak{Y}_j)_{\mathrm{red}} = Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\sqrt{\mathfrak{Q}_j}}\right).$$

We also put

$$I_0 = \bigcap_{i \in \mathfrak{I}} \mathfrak{q}_i \tag{14.5}$$

and

$$\mathfrak{X} = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{I_0}\right), \qquad \mathfrak{Y} = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\bigcap_{j\in\mathfrak{I}}\mathfrak{Q}_j}\right).$$

Then we have

$$|\mathfrak{X}| = \bigcup_{i\in\mathfrak{I}}|\mathfrak{X}_i|, \qquad |\mathfrak{Y}| = \bigcup_{j\in\mathfrak{I}}|\mathfrak{Y}_j|.$$

Lemma 14.4. *We have*

$$\underline{\mathcal{R}} \cong \frac{\Lambda^P\langle\!\langle y,y^{-1}\rangle\!\rangle[s]}{\mathrm{Clos}_{\mathfrak{v}_T^P}(\mathfrak{B}_i : i=1,\dots,n)}.$$

Proof. We use the following:

Sublemma 14.5. *Let z_i be as in (2.7). Then we have*

$$z_1^{k_1}\cdots z_m^{k_m} \equiv 0 \mod T^{\lambda_1(k_1+\cdots+k_m-n)/n}\frac{\Lambda_0^P\langle\!\langle y,y^{-1}\rangle\!\rangle[s]}{\mathrm{Clos}_{\mathfrak{v}_T^P}(\mathfrak{B}_i : i=1,\dots,n)}.$$

Proof. We remark that z_i satisfies the Stanley-Reisner relation and linear relation (that is $\sum_{j=1}^m v_{ij}z_j = 0$, $i=1,\dots,n$ see (13.2)) modulo $T^{\lambda_1}H(X;\Lambda_0)$. The cohomology ring $H(X;\mathbb{Q})$ is a quotient of the polynomial ring over z_i by the Stanley-Reisner relation and linear relation. (See Proposition in p.106 of [Ful], for example.) On $H(X;\mathbb{Q})$ the product of $(n+1)$ of z_i's are zero. Therefore we have

$$z_1^{k_1}\cdots z_m^{k_m} \equiv 0 \mod T^{\lambda_1(k_1+\cdots+k_n-n)/n}H(X;\Lambda_0)$$

if $k_1+\cdots+k_m > n$. The sublemma follows easily. □

Once we have Sublemma 14.5, the rest of the proof is the same as the proof of Lemma 13.10. □

Lemma 14.6. *If $(\mathfrak{y}_1,\dots,\mathfrak{y}_n,\mathfrak{s}) \in |Spec(\underline{\mathcal{R}})|$, then*

$$(\mathfrak{v}_T(\mathfrak{y}_1),\dots,\mathfrak{v}_T(\mathfrak{y}_n)) \in P. \tag{14.6}$$

If $(\mathfrak{y}_1,\dots,\mathfrak{y}_n,\mathfrak{s}) \in |Spec(\mathcal{R})|$ and (14.6) is satisfied, then $(\mathfrak{y}_1,\dots,\mathfrak{y}_n,\mathfrak{s}) \in |Spec(\underline{\mathcal{R}})|$.

Proof. The first half is a consequence of Lemmas 2.13 (B) and 14.4. The second half follows from the fact that $y_i \mapsto \mathfrak{y}_i$ induces continuous map from $(\Lambda[y,y^{-1}],\mathfrak{v}_T^P)$ to Λ if $(\mathfrak{v}_T(\mathfrak{y}_1),\dots,\mathfrak{v}_T(\mathfrak{y}_n)) \in P$. □

Corollary 14.7. *We have*

$$\overline{I}_0 = \overline{I},$$

where $\overline{I}_0$ is the closure of I_0.

Proof. Let $i \notin \mathfrak{I}$. We will prove $\overline{\mathfrak{q}}_i = \Lambda[s,y,y^{-1}]$.

In fact, since $i \notin \mathfrak{I}$, we have

$$\left|Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\overline{\mathfrak{q}}_i}\right)\right| \cap |Spec(\underline{\mathcal{R}})| \subseteq \left|Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i}\right)\right| \cap |Spec(\underline{\mathcal{R}})| = \emptyset.$$

On the other hand, since $\overline{\mathfrak{q}}_i \supseteq \overline{I}$, it follows from Lemma 14.6 that

$$\left|Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\overline{\mathfrak{q}}_i}\right)\right| \subseteq |Spec(\underline{\mathcal{R}})|.$$

Therefore

$$\left|Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\overline{\mathfrak{q}}_i}\right)\right| = \emptyset.$$

Namely $\overline{\mathfrak{q}}_i = \Lambda[s,y,y^{-1}]$ as claimed.

Therefore $1 \in \overline{\mathfrak{q}}_i$ for $i \notin \mathfrak{I}$. The corollary follows easily. ☐

Proposition 14.8. *For each $i \in \mathfrak{I}$ there exists $j \in \mathfrak{J}$ such that $|\mathfrak{X}_i| \subseteq |\mathfrak{Y}_j|$.*

Proof. Let $\mathfrak{Z}$ be one of the irreducible components of $\mathfrak{X}_i \cap Spec(\underline{\mathcal{R}})$. We will prove the coincidence of geometric points in $\mathfrak{X}_i$ and $\mathfrak{Z}$. The proposition then will follow.

Lemma 14.9. *Let $\mathfrak{Z}$, $\mathfrak{X}_i$ be as above, and $|\mathfrak{Z}| \neq |\mathfrak{X}_i|$. Then, there exists a commutative diagram of Λ-schemes:*

$$\begin{tikzcd}
(\mathfrak{X}_i)_{\mathrm{red}} & \mathfrak{Z}_{\mathrm{red}} \arrow[l] \\
Spec(B) \arrow[u, "\mathfrak{f}"'] & Spec(\Lambda) \arrow[l] \arrow[u, "\bar{\mathfrak{f}}"']
\end{tikzcd} \tag{14.7}$$

with the following properties. Let $\tilde{\mathfrak{y}} \in |Spec(B)|$ be the geometric point corresponding to the lower horizontal arrow.

1. *$Spec(B)$ is smooth at $\tilde{\mathfrak{y}}$.*
2. *$\mathfrak{f}$ is an embedding.*
3. *$|Spec(B)| \cap |Spec(\underline{\mathcal{R}})|$ is zero dimensional. In particular, $\mathfrak{f}$ cannot be lifted to a morphism $Spec(B) \to \mathfrak{Z}_{\mathrm{red}}$.*

Proof. We may take a 1-dimensional irreducible and reduced scheme $\mathfrak{C} = Spec(A) \subset (\mathfrak{X}_i)_{\mathrm{red}}$, which is a subscheme of $Spec(\mathcal{R})$ with $\dim(\mathfrak{C} \cap \mathfrak{Z}_{\mathrm{red}}) = 0$. Let $\mathfrak{y}$ be the intersection point. Then we normalize $\mathfrak{C}$ at $\mathfrak{y}$. We pick a point $\tilde{\mathfrak{y}}$ in the inverse image of $\mathfrak{y}$ under the normalization map. Its appropriate open subscheme is the required $Spec(B)$. ☐

Let $\mathfrak{m}_{\tilde{\mathfrak{y}}}$ be a maximal ideal of B corresponding to $\tilde{\mathfrak{y}}$. The localization $B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$ is a one dimensional regular local ring. Let S be a generator of the ideal $\mathfrak{m}_{\tilde{\mathfrak{y}}}$ in $B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$. The inclusion $\Lambda[S] \to B$ can be regarded as a map $Spec(B) \to Spec(\Lambda[S])$. Its differential at $\tilde{\mathfrak{y}}$ is nonzero. We next want to apply 'inverse mapping theorem' for this map at $\tilde{\mathfrak{y}}$. Of course, in algebraic geometry we do not have one. In other words, the inverse does not exist on a Zariski open subset. However, if we consider the infinitesimal neighborhood of $S = 0$ in $Spec(\Lambda[S])$, we have an inverse. The detail will be as follows.

We regard $y_i, s \in B$ as Λ-valued functions on $|Spec(B)|$. We write $s = y_0$ and define $\mathfrak{y}_i = y_i(\tilde{\mathfrak{y}})$ $(i = 0, \dots, n)$. Here $\mathfrak{y}_i \neq 0$ for $i \neq 0$.

Now we take generators $z_1, \dots, z_h$ of B as a Λ-algebra such that $z_i(\tilde{\mathfrak{y}}) = 0$ $(i = 1, \dots, h)$. Since $z_i \in \mathfrak{m}_{\tilde{\mathfrak{y}}} \subset B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$, we have

$$z_i = S^k \frac{P_i(z_1, \dots, z_h)}{Q_i(z_1, \dots, z_h)} \tag{14.8}$$

with $P_i(0, \dots, 0) \neq 0$, $Q_i(0, \dots, 0) \neq 0$, $k \in \mathbb{Z}_{>0}$. We put

$$\begin{aligned} F_i(Z_1, \dots, Z_h, S) &= Z_i - S^k \frac{P_i(Z_1, \dots, Z_h)}{Q_i(Z_1, \dots, Z_h)} \\ &= Z_i - S^k U_i(Z_1, \dots, Z_h). \end{aligned} \tag{14.9}$$

Lemma 14.10. *Under the situation above, we have* $K, C > 0$ *such that if*

$$\mathfrak{v}_T(a_i) \geq K \quad (i = 1, \dots, h), \tag{14.10}$$

then $Q_i(a_1, \dots, a_h) \neq 0$ *and the following inequalities hold*

$$\mathfrak{v}_T\left(\frac{\partial U_i}{\partial Z_j}(a_1, \dots, a_h)\right) > -C, \tag{14.11}$$

$$\mathfrak{v}_T\left(\frac{\partial^2 U_i}{\partial Z_j \partial Z_k}(a_1, \dots, a_h)\right) > -C. \tag{14.12}$$

The proof is easy and is left to the reader. We remark that if both of $(a_1, \dots, a_h)$ and $(a'_1, \dots, a'_h)$ satisfy (14.10), then (14.12) yields

$$\begin{aligned} &\mathfrak{v}_T\left(\frac{\partial U_i}{\partial Z_j}(a_1, \dots, a_h) - \frac{\partial U_i}{\partial Z_j}(a'_1, \dots, a'_h)\right) \\ &> \inf_{i=1,\dots,h} \mathfrak{v}_T(a_i - a'_i) - C. \end{aligned} \tag{14.13}$$

We choose $K > 2C$. Now we consider $\mathfrak{S}$ with $\mathfrak{v}_T(\mathfrak{S}) \geq K$. Then we have

$$\frac{\partial F_i}{\partial Z_j}(0, \dots, 0, \mathfrak{S}) \equiv \begin{cases} 1 \mod T^{K-C} & \text{if } i = j \\ 0 \mod T^{K-C} & \text{if } i \neq j. \end{cases} \tag{14.14}$$

This is a consequence of (14.11). We also have

$$F_i(0, \dots, 0, \mathfrak{S}) \equiv 0 \mod T^{K-C}.$$

Now we iteratively define a_i^ℓ $(i = 1, \dots, h)$ by induction on $\ell = 0, 1, \dots$ such that

$$(\Delta_{\ell+1,i})_{i=1}^h = -\left(\left[\frac{\partial F_i}{\partial Z_j}(a_1^\ell, \dots, a_h^\ell, \mathfrak{S})\right]_{i=1,\ j=1}^{i=h,\ j=h}\right)^{-1} (F_i(a_1^\ell, \dots, a_h^\ell, \mathfrak{S}))_{i=1}^h$$

where $a_i^0 = 0$, $a_i^\ell = \Delta_{1,i} + \dots + \Delta_{\ell,i}$.

We can prove by induction on ℓ that

$$F_i(a_1^\ell, \dots, a_h^\ell, \mathfrak{S}) \equiv 0 \mod T^{(\ell+1)(K-C)}$$

and

$$\Delta_{\ell+1,i} \equiv 0 \mod T^{(\ell+1)(K-C)}.$$

We define

$$a_i(\mathfrak{S}) = \lim_{\ell \to \infty} a_i^\ell. \tag{14.15}$$

Then they satisfy the equations

$$F_i(a_1(\mathfrak{S}), \dots, a_h(\mathfrak{S}), \mathfrak{S}) = 0, \quad i = 1, \dots, h.$$

On the other hand, we can write

$$y_i = R_i(z_1, \dots, z_h), \quad i = 0, \dots, n \tag{14.16}$$

for $R_i \in \Lambda[Z_1, \dots, Z_h]$. We define

$$y_i(\mathfrak{S}) = R_i(a_1(\mathfrak{S}), \dots, a_h(\mathfrak{S})).$$

Lemma 14.11. *For any* $\mathfrak{S}$ *with sufficiently large* $\mathfrak{v}_T(\mathfrak{S})$, $y(\mathfrak{S})$ *is a geometric point of Spec*(B).

We will prove it later.

We consider $\mathfrak{S}$ with sufficiently large $\mathfrak{v}_T(\mathfrak{S})$ as in Lemma 14.11. Note $\mathfrak{v}_T(y_i(\mathfrak{S})) = \mathfrak{v}_T(y_i(0)) = \mathfrak{v}_T(\mathfrak{y}_i)$ holds if $\mathfrak{v}_T(\mathfrak{S})$ is sufficiently large.

On the other hand, since $\mathfrak{y}$ is in $|Spec(\underline{\mathcal{R}})|$, Lemma 14.6 implies

$$(\mathfrak{v}_T(\mathfrak{y}_1), \dots, \mathfrak{v}_T(\mathfrak{y}_n)) \in P.$$

Therefore $y(\mathfrak{S})$ is also a geometric point of $|Spec(\underline{\mathcal{R}})|$. Moreover, from $U_i(0, \dots, 0) \neq 0$, we have $z_i(\mathfrak{S}) \neq 0$ for at least one i. Since $Spec(B)$ is a normalization of $\mathfrak{C} = Spec(A)$, B is a finite A-module and so the mapping

$$(a_1, \dots a_h) \mapsto (R_0(a_1, \dots, a_h), \dots, R_n(a_1, \dots, a_h))$$

is finite to one. Thus not only $z(\mathfrak{S})$ but also $y(\mathfrak{S})$ are non-constant.

Hence $Spec(B)$ has infinitely many geometric points in $|Spec(\underline{\mathcal{R}})|$. This is a contradiction. The proof of Proposition 14.8 is completed assuming Lemma 14.11. □

Proof of Lemma 14.11. We may identify B with

$$\frac{\Lambda[Z_1, \dots, Z_h, S]}{\mathfrak{p}}$$

where Z_i corresponds to z_i in B. We denote by $B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$ the localization of B by the maximal ideal $\mathfrak{m}_{\tilde{\mathfrak{y}}}$ corresponding to the point $\tilde{\mathfrak{y}}$. We next take the completion $B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$ of $B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$ by the ideal $\mathfrak{m}_{\tilde{\mathfrak{y}}}$. It is a complete discrete valuation ring over the field Λ (with respect to the $\mathfrak{m}_{\tilde{\mathfrak{y}}}$ topology), hence isomorphic to the formal power series ring $\Lambda[[t]]$. (See for example, [Ei] Proposition 10.16, [Mat] Theorem 37 in (17.H).) (Here we used the fact that Λ is algebraically closed in order to show $\Lambda \cong B/\mathfrak{m}_{\tilde{\mathfrak{y}}} \cong B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}/\mathfrak{m}_{\tilde{\mathfrak{y}}} B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}$. (See [FOOO4] Lemma A.1.)) Note that we may take $S \in \Lambda[[t]]$ as the generator S of $\mathfrak{m}_{\tilde{\mathfrak{y}}}$ we took before. Thus we may take $t = S$.

We have the following diagram.

$$\begin{tikzcd}
B \cong \Lambda[Z_1, \dots, Z_h, S]/\mathfrak{p} \arrow[r] \arrow[d] & \Lambda[Z_1, \dots, Z_h, S]/\mathfrak{p}' \arrow[d] \\
B_{\mathfrak{m}_{\tilde{\mathfrak{y}}}} \arrow[r] \arrow[d] & B/\mathfrak{m}_{\tilde{\mathfrak{y}}} \arrow[d, "\tilde{\mathfrak{f}}"] \\
B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}} \cong \Lambda[[S]] \arrow[r] & B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}}/\mathfrak{m}_{\tilde{\mathfrak{y}}} B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}} \cong \Lambda
\end{tikzcd} \tag{14.17}$$

The element $z_i \in B$ induces $[z_i] \in B^{\wedge}_{\mathfrak{m}_{\tilde{\mathfrak{y}}}} \cong \Lambda[[S]]$. We denote this element by $\mathfrak{f}_i(S)$. For any $W \in \mathfrak{p} \subset \Lambda[Z_1, \dots, Z_h, S]$ we have

$$W(\mathfrak{f}_1(S), \dots, \mathfrak{f}_h(S), S) = 0 \in \Lambda[[S]]. \tag{14.18}$$

We remark that

$$Q_i(Z_1, \dots, Z_h) F_i(Z_1, \dots, Z_h, S) \in \mathfrak{p}.$$

Therefore by (14.18) we have

$$Q_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S)) F_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S), T^k S) = 0.$$

Since $Q_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S)) \neq 0$ for sufficiently large K, we obtain

$$F_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S), T^k S) = 0.$$

In other words, by (14.9) we have

$$\begin{aligned}\mathfrak{f}_i(T^K S) &= T^{kK} S^k \frac{P_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S))}{Q_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S))} \\ &= T^{kK} S^k U_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S)).\end{aligned} \tag{14.19}$$

We observe that Q_i is nonzero at $Z_1 = \dots = Z_h = 0$, therefore we may regard

$$U_i \in \Lambda[[Z_1, \dots, Z_h]].$$

Note $\mathfrak{f}_i(S) \in S\Lambda[[S]]$. Therefore the substitution

$$U_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S))$$

makes sense using S-adic topology. (Namely as a formal power series of S.) Now we observe that the equation

$$\mathfrak{f}_i(T^K S) = T^{kK} S^k U_i(\mathfrak{f}_1(T^K S), \dots, \mathfrak{f}_h(T^K S)) \tag{14.20}$$

with $\mathfrak{f}_i(0) = 0$, *uniquely* characterizes $\mathfrak{f}_i(S) \in \Lambda[[S]]$.

In fact, we define $X_{i,k}$ $(i = 1, \dots, h)$ inductively by $X_{i,0} = 0$ and

$$X_{i,k+1} = T^{kK} S^k U_i(X_{1,k}, \dots, X_{h,k}). \tag{14.21}$$

Then $\{X_{i,k}\}_{k=0,1,\dots}$ converges to $\mathfrak{f}_i(T^K S)$ in S-adic topology and is necessary the unique solution of (14.20).

Now we consider the same inductive construction of the solution of (14.20), but this time with T-adic topology. Namely we use $\mathfrak{v}_T$ with $\mathfrak{v}_T(S) = 0$. Then for sufficient large K, we have a solution of (14.21) in the way we described during the iterative argument of the proof of Proposition 14.8. It gives $a_i(T^K S)$. Namely, if we substitute S with element c of Λ_0, we obtain $a_i(cT^K)$ which we defined before in (14.15) with $\mathfrak{S} = cT^K$ satisfying $\mathfrak{v}_T(\mathfrak{S}) \ge K$.

Now by the uniqueness of the solution of the recursive equation (14.21) we have

$$a_i(T^K S) = \mathfrak{f}_i(T^K S).$$

In particular,

$$W(a_1(T^K S), \dots, a_h(T^K S), T^k S) = 0$$

for all $W \in \mathfrak{p}$. This proves the lemma. □

Remark 14.12. Suppose that $\mathfrak{f}_i(S) \in T^{-C}\Lambda_0[[S]]$ for some C. (We do not know whether this is true or not.) In that case we can discuss as follows. We put $\mathfrak{g}_i = R_i(\mathfrak{f}_1(S), \dots, \mathfrak{f}_h(S))$, where $R_i \in \Lambda[Z_1, \dots, Z_h]$ as in (14.16). We then can write

$$\mathfrak{g}_i(S) = \sum_{j=0}^{\infty} T^{\lambda_{i,j}} \mathfrak{g}_{i,j}(S).$$

Here $\mathfrak{g}_{i,j}(S) \in \mathbb{C}[[S]] \setminus \{0\}$ and $\lambda_{i,j} \in \mathbb{R}$. We may assume $\lambda_{i,j} < \lambda_{i,j+1}$.

We put

$$j_i^0 = \inf\{j \mid \mathfrak{f}_{i,j}(0) \neq 0\}.$$

It is easy to see that $\mathfrak{y}_i = \mathfrak{f}_i(0)$. Therefore we have

$$\mathfrak{v}_T(\mathfrak{y}) = (\lambda_{1,j_1^0}, \dots, \lambda_{n,j_n^0}) = u \in P. \tag{14.22}$$

Now we write

$$f_{i,j}(S) = d_{i,j} S^{\rho_{i,j}} + \text{higher order terms}$$

with $d_{i,j} \in \mathbb{C} \setminus 0$ and $\rho_{i,j} > 0$. For $\kappa > 0$ we define

$$\lambda_i(\kappa) = \inf\{\lambda_{i,j} + \rho_{i,j}\kappa \mid j = 0, \dots, j_i^0\}.$$

We take a complex number $c \in \mathbb{C}$ which satisfies

$$\sum_{\substack{j=0,\dots,j_i^0;\\ \lambda_{i,j}+\rho_{i,j}\kappa=\lambda_i(\kappa)}} d_{i,j}c^{\rho_{i,j}} \neq 0.$$

We note that we can choose c independent of κ, because it is enough to take c so that $\sum_{j\in\mathcal{J}} d_{i,j}c^{\rho_{i,j}} \neq 0$ for all $\mathcal{J} \subseteq \{0,\dots,j_i^0\}$. Then we have

$$\vec{\mathfrak{v}}_T(\mathfrak{f}(cT^{\kappa})) = (\lambda_1(\kappa), \dots, \lambda_n(\kappa)),$$

which we denote by $\lambda(\kappa)$. The function $\lambda(\kappa)$ is piecewise affine. In particular, it is continuous. (14.22) implies that $\lambda(\kappa) \in P$ for sufficiently large κ. Indeed, $\lambda(\kappa)$ is constant for sufficiently large κ. It actually imlies $\lambda(\kappa) \in P$ for all $\kappa > 0$. In fact if $\lambda(\kappa) \notin P$ for some κ then there exists κ_0 such that $\lambda(c\kappa_0) \in \partial P$. This implies that $(\mathfrak{y}'_1, \dots, \mathfrak{y}'_n) = \mathfrak{f}(cT^{\kappa_0})$ is a solution of $\mathfrak{B}_i = 0$ and its valuation lies on the boundary of moment polytope. This is impossible since the leading order equation of $\mathfrak{B}_i$ is the same as the leading order equation of $y_i\frac{\partial\mathfrak{PO}_{\mathfrak{b}}}{\partial y_i}$.

In a similar situation, [FOOO5] Section 5 provides an example where $\kappa \mapsto (\lambda_1(\kappa), \dots, \lambda_n(\kappa))$ is a non-constant affine map. We can also find an example where it is piecewise affine in a similar way.

The following is a consequence of Proposition 14.8.

Corollary 14.13. *If $i \in \mathfrak{I}$, then for any $\mathfrak{s} \in \Lambda$ we have:*

$$\operatorname{rank}_{\Lambda} \frac{\Lambda[s, y, y^{-1}]}{\mathfrak{q}_i(s - \mathfrak{s})} < \infty. \tag{14.23}$$

We also have

$$\dim \mathfrak{X} \leq 1. \tag{14.24}$$

Proof. Theorems 10.1 and 11.1 imply that the fiber $\pi^{-1}(\mathfrak{s}) \cap \mathfrak{Y}$ is 0 dimensional. Then Proposition 14.8 implies that $\pi^{-1}(\mathfrak{s}) \cap \mathfrak{X}$ is 0 dimensional. This implies (14.23). (14.24) is its immediate consequence. □

Proposition 14.14. *$\pi : \mathfrak{X} \to Spec(\Lambda[s])$ is projective.*

Proof. (The argument below is of similar flavor to the valuative criterion of properness (See Theorem 4.7 Chapter II [Ha]).)

We denote by $\mathbb{P}^1_{y_i}$ the projective line $Proj(\Lambda[y_0, y_i])$. We have $\prod_{i=1}^n \mathbb{P}^1_{y_i} \times Spec(\Lambda[s]) \supset \mathfrak{X}$. (Here the product is a fiber product over $Spec(\Lambda)$.) Let $\overline{\mathfrak{X}}$ be the Zariski closure of $\mathfrak{X}$ in $\prod_{i=1}^n \mathbb{P}^1_{y_i} \times Spec(\Lambda[s])$. It suffices to show $\overline{\mathfrak{X}} = \mathfrak{X}$. We will prove this by contradiction.

(14.24) implies that

$$\#(|\overline{\mathfrak{X}}| \setminus |\mathfrak{X}|) < \infty. \tag{14.25}$$

Let $(\mathfrak{y}, \mathfrak{s}) \in \overline{\mathfrak{X}} \setminus \mathfrak{X}$. We may replace the coordinate y_i by y_i^{-1} and may assume that $\mathfrak{y} \in Spec(\Lambda[y_1, \dots, y_n])$. (The map $y_i \mapsto y_i^{-1}$ is an automorphism of $\mathbb{P}^1_{y_i}$.)

We put $\mathfrak{y} = (\mathfrak{y}_1, \dots, \mathfrak{y}_n)$. Since $\mathfrak{y} \notin \mathfrak{X}$, there exists i such that $\mathfrak{y}_i = 0$. We take one dimensional scheme $\mathfrak{C} \subset \overline{\mathfrak{X}}$ such that $\mathfrak{y} \in \mathfrak{C}$ and $\mathfrak{C} \setminus \{\mathfrak{y}\} \subset \mathfrak{X}$. Replacing $\mathfrak{C}$ by its normalization $Spec(B)$ we may assume that $\mathfrak{y}$ is a smooth point of $Spec(B)$. (In the proof of Proposition 14.8 we denote it by $\tilde{\mathfrak{y}}$.) Let $\mathfrak{m}_{\mathfrak{y}}$ be the maximal ideal of B corresponding to $\mathfrak{y}$ and let S be a generator of $\mathfrak{m}_{\mathfrak{y}} B_{\mathfrak{m}_{\mathfrak{y}}}$.

In the same way as the proof of Proposition 14.8 we can find K such that if $\kappa > K$ then there exists a geometric point $\mathfrak{y}(T^{\kappa}) = (\mathfrak{y}_1(T^{\kappa}), \dots, \mathfrak{y}_n(T^{\kappa}))$ of B such that $\mathfrak{y} - \mathfrak{y}(T^{\kappa}) \equiv 0 \mod T^{\kappa}$. If $\mathfrak{y}_i \neq 0$, then $\mathfrak{v}_T(\mathfrak{y}_i(T^{\kappa})) = \mathfrak{v}_T(\mathfrak{y}_i)$ for sufficiently large κ. If $\mathfrak{y}_i = 0$, then $\mathfrak{v}_T(\mathfrak{y}_i(T^{\kappa})) \geq \kappa$. Since $\mathfrak{y}_i = 0$ for at least one i, it implies that $\vec{\mathfrak{v}}_T(\mathfrak{y}(T^{\kappa})) \notin P$ for sufficiently large κ. Therefore B has infinitely many geometric points outside $\mathfrak{X}$. This contradicts to $B \subset \overline{\mathfrak{X}}$ and (14.25). ☐

Proposition 14.15. *We have*

(1) $\dim \mathfrak{X} = 1$.
(2) $\pi : \mathfrak{X} \to Spec(\Lambda[s])$ *is flat.*
(3) *Let* $\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}$ *be the maximal ideal corresponding to* $(\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}|$. *Then* $\mathfrak{B}_1, \dots, \mathfrak{B}_n$ *is a regular sequence in* $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}}$.

Proof. (3) Let $\mathfrak{x} = (\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}|$ and $\mathfrak{p}_{\mathfrak{x}}$ or $\mathfrak{m}_{\mathfrak{x}}$ the ideals of $\mathcal{R}$ or $\Lambda[s, y, y^{-1}]$ corresponding to $\mathfrak{x}$, respectively. By Corollary 14.13, we have

$$0 = \dim \frac{\mathcal{R}_{\mathfrak{p}_{\mathfrak{x}}}}{(s - \mathfrak{s})} = \dim \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{(\mathfrak{B}_1, \dots, \mathfrak{B}_n, s - \mathfrak{s})}.$$

It follows that $(\mathfrak{B}_1, \dots, \mathfrak{B}_n, s - \mathfrak{s})$ is a system of parameter of a Cohen-Macaulay ring $\Lambda[y, y^{-1}, s]_{\mathfrak{m}_{\mathfrak{x}}}$. (See 12.J [Mat].) Therefore it is a regular sequence. (See Theorem 31 [Mat].) (3) follows.

(2) Note that the flatness of π is equivalent to the flatness for all localizations at all maximal ideals of $Spec(\Lambda[s])$. Since the fibers of $\pi : \mathfrak{X} \to Spec(\Lambda[s])$ are zero dimensional, it suffices to show that $s - \mathfrak{s}$ is not a zero divisor in $\mathcal{R}_{\mathfrak{p}_{\mathfrak{x}}}$ for any $\mathfrak{x} = (\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}|$. This follows from the fact that $\mathfrak{B}_1, \dots, \mathfrak{B}_n, s - \mathfrak{s}$ is a regular sequence in $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$.

(1) By (3) $\mathfrak{B}_1, \dots, \mathfrak{B}_n$ is a regular seqence in $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$, where $\mathfrak{m}_{\mathfrak{x}}$ is any maximal ieal of $\mathfrak{X}$. Therefore $\dim \mathfrak{X} = \dim \Lambda[s, y, y^{-1}] - n = 1$. (See Theorem 29 [Mat].) ☐

Proposition 14.16. *For any* $i \in \mathfrak{I}$ *we have the following.*

(1) $\dim \mathfrak{X}_i = 1$.
(2) $\pi : \mathfrak{X}_i \to Spec(\Lambda[s])$ *is flat.*

Proof. (1) : By Proposition 14.15 (1), $\dim \mathfrak{X}_i$ is either 1 or 0. Suppose $\dim \mathfrak{X}_i = 0$. Let $\mathfrak{x} \in |\mathfrak{X}_i|$. By Proposition 14.15 (3), $\mathfrak{B}_1, \dots, \mathfrak{B}_n, s - \mathfrak{s}$ is a regular sequence in $\Lambda[y, y^{-1}, s]_{\mathfrak{m}_{\mathfrak{x}}}$. By Lemma 14.17 below, $\mathfrak{q}_i \Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$ appears in the irredundant primaly decomposition of $(\mathfrak{B}_1, \dots, \mathfrak{B}_n)\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$. Then by Theorem 14.19 below, the height of $\mathfrak{q}_i$ in $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$ is n. Namely

$$\dim \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{\mathfrak{q}_i \Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}} = 1.$$

This contradicts to $\dim \mathfrak{X}_i = 0$. Thus we obtain (1).

To prove (2) we prepare some elementary lemmas and a basic fact.

Lemma 14.17. *Let* R *be an integral domain and* $\mathfrak{q}$, $\mathfrak{Q}$ *primaly ideals of* R. *Suppose that* $\mathfrak{p}$ *is a prime ideal containing both* $\mathfrak{q}$ *and* $\mathfrak{Q}$. *If* $\mathfrak{q} R_{\mathfrak{p}} \subseteq \mathfrak{Q} R_{\mathfrak{p}}$, *then* $\mathfrak{q} \subseteq \mathfrak{Q}$.

Proof. Let $x \in \mathfrak{q}$. Then by assumption we have $a \in R$, $b \in R \setminus \mathfrak{p}$ and $c \in \mathfrak{Q}$ such that $x = ca/b$ in $R_{\mathfrak{p}}$. It follows that $xb = ca$ in R. (We use the fact that R is an integral domain here.) Therefore $xb \in \mathfrak{Q}$. Since $b \in R \setminus \mathfrak{p}$ and $\mathfrak{p} \supseteq \mathfrak{Q}$ is a prime ideal, it follows that $x \in \mathfrak{Q}$. We thus prove $\mathfrak{q} \subseteq \mathfrak{Q}$. □

Lemma 14.18. *Let R be a commutative ring with unit, A a commutative R-algebra and J_0 an ideal in A. Suppose that $J_0 = \cap_{i \in \mathfrak{I}} \mathfrak{q}_i$ is an irredundant primary decomposition without embedding primes. Then the following two conditions are equivalent:*

1. *A/J_0 is torsion-free as an R-module.*
2. *For any i, $A/\mathfrak{q}_i$ is torsion-free as an R-module.*

Proof. Suppose that $A/\mathfrak{q}_i$ has a torsion, i.e., there exist $a \in R \setminus \{0\}$ and $f \in A \setminus \mathfrak{q}_i$ such that $f \cdot a \in \mathfrak{q}_i$. Since the decomposition $J_0 = \cap_i \mathfrak{q}_i$ has no embedded primes, we can pick $g_j \in \mathfrak{q}_j \setminus \sqrt{\mathfrak{q}_i}$ for each $j \neq i$. Set $h = f \prod_{j \neq i} g_j$. Then we find that $h \notin \mathfrak{q}_i$, hence $h \notin J_0$, and $ah = (af) \prod_{j \notin i} g_j \in J_0$. Namely, h is a torsion element in A/J_0. The converse implication is clear. □

Theorem 14.19. (Macauley's unmixedness theorem. cf. Corollary 18.14 [Ei]) *Let R be a Cohen-Macauley ring and I an ideal generated by an R-regular sequence $(x_1, \dots, x_k)$, then I has no embedded primes and the height of I is k.*

Proof of Proposition 14.16 (2)*:* Let $\mathfrak{x} = (\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}_i|$. By definition (14.4) we have

$$\begin{aligned}\mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}} &= \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{(\mathfrak{B}_1, \dots, \mathfrak{B}_n)_{\mathfrak{m}_{\mathfrak{x}}}} = \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{\bigcap_{i \in \mathfrak{I}_+} (\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}} \\ &= \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{\bigcap_{i \in \mathfrak{I}} (\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}} = \frac{\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}}{\bigcap_{\mathfrak{q}_i \subset \mathfrak{m}_{\mathfrak{x}}} (\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}},\end{aligned}$$

because for $i \in \mathfrak{I}_+ \setminus \mathfrak{I}$ we have $(\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}} = (1)$, i.e., $\mathfrak{q}_i \subset \mathfrak{m}_{\mathfrak{x}}$. By Proposition 14.15 (3), $\mathfrak{B}_1, \dots, \mathfrak{B}_n$ is a regular sequence in $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{\mathfrak{x}}}$. Therefore, Proposition 14.15 (2), Theorem 14.19 and Lemma 14.18 above imply the flatness of $\pi : \mathfrak{X}_i \to Spec(\Lambda[s])$. □

During the proof of Proposition 14.16, we have showed that irredundant primary decompositions (14.5) has no embedded primes.

Lemma 14.20. *In the situation of Proposition 14.8 we have $|\mathfrak{X}_i| = |\mathfrak{Y}_j|$.*

Proof. By Corollary 14.7 we have

$$|Spec(\underline{\mathcal{R}})| \subseteq |\mathfrak{X}|. \tag{14.26}$$

Therefore $\dim \mathfrak{Y}_j \leq 1$. On the other hand, $\dim \mathfrak{X}_i = 1$ by Proposition 14.16. Therefore the lemma follows from Proposition 14.8. □

For each $i \in \mathfrak{I}$ we take $j(i)$ such that $\mathfrak{X}_i$ and $\mathfrak{Y}_{j(i)}$ are as in Proposition 14.8.

Proposition 14.21. *We have $\mathfrak{X}_i = \mathfrak{Y}_{j(i)}$, i.e., $\mathfrak{q}_i = \mathfrak{Q}_{j(i)}$.*

Proof. By Corollary 14.7 we have $I_0 \subseteq \overline{I}$. It follows from Lemma 14.20 that

$$|\mathfrak{X}| = |Spec(\underline{\mathcal{R}})|.$$

Therefore by Lemma 14.20 again we have $\dim \mathfrak{Y}_j = 0$, i.e.,the associated prime of $\mathfrak{Q}_j$ is an embedded prime, if $j \notin \{j(i) \mid i \in \mathfrak{I}\}$.

We put

$$B_1 = \bigcup_{j \notin \{j(i) | i \in \mathfrak{I}\}} \pi(|\mathfrak{Y}_j|).$$

$B_1 \subset \Lambda$ is a finite set. We put

$$B_2 = \bigcup_{i_1, \neq i_2; i_1, i_2 \in \mathfrak{I}} \pi(|\mathfrak{X}_{i_1}| \cap |\mathfrak{X}_{i_j}|).$$

$B_2 \subset \Lambda$ is also a finite set. Let $B = B_1 \cup B_2$. We take $c \in \Lambda \setminus \pi(B)$. Let $\mathfrak{p}$ be the prime ideal $\sqrt{\mathfrak{q}_i} = \sqrt{\mathfrak{Q}_{j(i)}}$. Pick a maximal ideal $\mathfrak{m}$ containing $\mathfrak{p}$ and $s - c$. In other words, the maximal ideal $\mathfrak{m} = \mathfrak{m}_{\mathfrak{x}}$ defines a point

$$\mathfrak{x} = (c, \mathfrak{y}) \in |\mathfrak{X}_i| \cap \pi^{-1}(c) = |\mathfrak{Y}_{j(i)}| \cap \pi^{-1}(c).$$

We denote by $\mathfrak{m}_{\mathfrak{y}}$ the maximal ideal in $\Lambda[y, y^{-1}]$, which defines $\mathfrak{y} \in (\Lambda^{\times})^n$.

Now we compare the rings $\{\mathcal{R}/(s-c)\mathcal{R}\}_{\mathfrak{m}_{\mathfrak{y}}}$ and $\{\underline{\mathcal{R}}/(s-c)\underline{\mathcal{R}}\}_{\mathfrak{m}_{\mathfrak{y}}}$. By Lemma 14.6 and Proposition 14.8, we have $\vec{\mathfrak{v}}_T(\mathfrak{y}) \in P$. In the same way as in Section 7 in [FOOO4], these rings are characterized as the common generalized eigenspaces of the multiplication by y_i with eigenvalues $\mathfrak{y}_i$. Then by the same way in the proof of Proposition 8.6 in [FOOO4] we can show that the natural projection $\mathcal{R} \to \underline{\mathcal{R}}$ induces an isomorphism

$$\left(\mathcal{R}/(s-c)\mathcal{R}\right)_{\mathfrak{m}_{\mathfrak{y}}} \cong \left(\underline{\mathcal{R}}/(s-c)\underline{\mathcal{R}}\right)_{\mathfrak{m}_{\mathfrak{y}}}.$$

We first consider the kernel of the projection $\mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}} \to \underline{\mathcal{R}}_{\mathfrak{m}_{\mathfrak{x}}}$ which is $\overline{I}_{\mathfrak{m}_{\mathfrak{x}}}/I_{\mathfrak{m}_{\mathfrak{x}}}$. We have the following:

Lemma 14.22. *We have*

$$I_{\mathfrak{m}_{\mathfrak{x}}} = (\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}, \qquad \overline{I}_{\mathfrak{m}_{\mathfrak{x}}} = (\mathfrak{Q}_{j(i)})_{\mathfrak{m}_{\mathfrak{x}}}.$$

In particular,

$$\overline{I}_{\mathfrak{m}_{\mathfrak{x}}}/I_{\mathfrak{m}_{\mathfrak{x}}} \cong (\mathfrak{Q}_{j(i)})_{\mathfrak{m}_{\mathfrak{x}}}/(\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}.$$

Proof. Recall from (14.3) that $I = \bigcap_{i \in \mathfrak{I}_+} \mathfrak{q}_i$ and $\overline{I} = \bigcap_{j \in \mathfrak{I}} \mathfrak{Q}_j$. We note that $\mathfrak{q}_k$ for $k \in \mathfrak{I}_+ \setminus \mathfrak{I}$ cannot contain $\mathfrak{m}_{\mathfrak{x}}$, because $|\mathfrak{X}_k| \cap |Spec(\underline{\mathcal{R}})| = \emptyset$. Therefore we have

$$(I_0)_{\mathfrak{m}_{\mathfrak{x}}} = I_{\mathfrak{m}_{\mathfrak{x}}} = \bigcap_{\mathfrak{q}_j \subset \mathfrak{m}_{\mathfrak{x}}} (q_j)_{\mathfrak{m}_{\mathfrak{x}}}.$$

We also have

$$\overline{I}_{\mathfrak{m}_{\mathfrak{x}}} = \bigcap_{\mathfrak{Q}_k \subset \mathfrak{m}_{\mathfrak{x}}} (\mathfrak{Q}_k)_{\mathfrak{m}_{\mathfrak{x}}}.$$

By the choice of c, we find that $\mathfrak{X}_i$, resp. $\mathfrak{Y}_{j(i)}$, is the only irreducible component of $|\mathfrak{X}|$, resp. $|\mathfrak{Y}|$ containing the Λ-point $\mathfrak{x}$, and $\mathfrak{q}_i$, resp. $\mathfrak{Q}_{j(i)}$, is the unique primary ideal in the irredundant primary decompositions (14.3) of I, resp. $\overline{I}$, which is contained in $\mathfrak{m}_{\mathfrak{x}}$. Hence we obtain

$$I_{\mathfrak{m}_{\mathfrak{x}}} = (\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}} \subset \overline{I}_{\mathfrak{m}_{\mathfrak{x}}} = (\mathfrak{Q}_{j(i)})_{\mathfrak{m}_{\mathfrak{x}}}.$$

□

The rings $\mathcal{R}/(s-c)\mathcal{R}$ and $\underline{\mathcal{R}}/(s-c)\underline{\mathcal{R}}$ are $\Lambda[s, y, y^{-1}]$-algebras. By substituting c to s, we can also regard them as $\Lambda[y, y^{-1}]$-algebras. Then we have

$$\left(\mathcal{R}/(s-c)\mathcal{R}\right)_{\mathfrak{m}_{\mathfrak{y}}} \cong \mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}}/(s-c)\mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}}$$

and

$$(\underline{\mathcal{R}}/(s-c)\underline{\mathcal{R}})_{\mathfrak{m}_{\mathfrak{y}}} \cong \underline{\mathcal{R}}_{\mathfrak{m}_{\mathfrak{x}}}/(s-c)\underline{\mathcal{R}}_{\mathfrak{m}_{\mathfrak{x}}}.$$

By utilizing Lemma 14.22, we have the following commutative diagram such that the horizontal and vertical lines are exact sequences.

$$\begin{tikzcd}
 & 0 \arrow[r] \arrow[d] & 0 \arrow[r] \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] \arrow[d] & \operatorname{Ker} q \arrow[r] \arrow[d] & (\mathfrak{Q}_{j(i)}/\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & 0 \arrow[r] \arrow[d] & 0 \arrow[d] \\
0 \arrow[r] \arrow[d] & (s-c)\mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d, "q"'] & \mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & (\mathcal{R}/(s-c)\mathcal{R})_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & 0 \arrow[d] \\
0 \arrow[r] & (s-c)\underline{\mathcal{R}}_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & \underline{\mathcal{R}}_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & (\underline{\mathcal{R}}/(s-c)\underline{\mathcal{R}})_{\mathfrak{m}_{\mathfrak{x}}} \arrow[r] \arrow[d] & 0 \\
 & 0 \arrow[r] & 0 \arrow[r] & 0 &
\end{tikzcd}$$

Note that

$$\operatorname{Ker} q = (s-c)\mathcal{R}_{\mathfrak{m}_{\mathfrak{x}}} \cap \big(\mathfrak{Q}_{j(i)}/\mathfrak{q}_i\big)_{\mathfrak{m}_{\mathfrak{x}}} \cong \big(\mathfrak{q}_i + (s-c)\Lambda[s,y,y^{-1}] \cap \mathfrak{Q}_{j(i)}\big)_{\mathfrak{m}_{\mathfrak{x}}}/(\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}.$$

Since $\mathfrak{Q}_{j(i)}$ is a primary ideal and $s-c \notin \sqrt{\mathfrak{Q}_{j(i)}}$, we have

$$(s-c)\Lambda[s,y,y^{-1}] \cap \mathfrak{Q}_{j(i)} = (s-c)\mathfrak{Q}_{j(i)}.$$

Therefore we have

$$\operatorname{Ker} q \cong (s-c)(\mathfrak{Q}_{j(i)}/\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}.$$

It follows from the above diagram that

$$(s-c)(\mathfrak{Q}_{(j(i)}/\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}} \cong (\mathfrak{Q}_{(j(i)}/\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}}.$$

Now we set $M = \mathfrak{Q}_{j(i)}/\mathfrak{q}_i$. What we have showed is

$$M_{\mathfrak{m}_{\mathfrak{x}}}/(s-c)M_{\mathfrak{m}_{\mathfrak{x}}} = (M/(s-c)M)_{\mathfrak{m}_{\mathfrak{x}}} = 0.$$

Then we obtain

$$M_{\mathfrak{m}_{\mathfrak{y}}}/(s-c)M_{\mathfrak{m}_{\mathfrak{y}}} = (M/(s-c)M)_{\mathfrak{m}_{\mathfrak{y}}} \cong (M/(s-c)M)_{\mathfrak{m}_{\mathfrak{x}}} = 0. \tag{14.27}$$

When we localize $M/(s-c)M$ at the maximal ideal $\mathfrak{m}_{\mathfrak{x}}$ in $\Lambda[s,y,y^{-1}]$, we regard it as a $\Lambda[s,y,y^{-1}]$-module such that s acts on it by the multiplication by c. When we localize $M/(s-c)M$ at the maximal ideal $\mathfrak{m}_{\mathfrak{y}}$ in $\Lambda[y,y^{-1}]$, we regard it as a $\Lambda[y,y^{-1}]$-module. Therefore we obtain the isomorphism of the second equality in (14.27).

To apply Nakayama's lemma, we show the following lemma whose proof is given at the end of the proof of Porposition 14.21.

Lemma 14.23. *$M_{\mathfrak{m}_{\mathfrak{y}}}$ is a finitiely generated module over $\Lambda[s]$.*

We localize $\Lambda[s]$ and $M_{\mathfrak{m}_{\mathfrak{y}}}$ at the maximal ideal $(s-c) \subset \Lambda[s]$ to obtain

$$(M_{\mathfrak{m}_{\mathfrak{y}}})_{(s-c)} = (s-c)(M_{\mathfrak{m}_{\mathfrak{y}}})_{(s-c)}$$

by (14.27). Then Nakayama's lemma implies that $(M_{\mathfrak{m}_{\mathfrak{y}}})_{(s-c)} = 0$.

We can regard $(M_{\mathfrak{m}_{\mathfrak{y}}})_{(s-c)}$ as a $\Lambda[s,y,y^{-1}]$-module and localize it at the maximal ideal $\mathfrak{m}_{\mathfrak{x}}$ in $\Lambda[s,y,y^{-1}]$. Then by noticing the fact that $\mathfrak{m}_{\mathfrak{y}} = \mathfrak{m}_{\mathfrak{x}} \cap \Lambda[y,y^{-1}]$ and $(s-c)\Lambda[s] = \mathfrak{m}_{\mathfrak{x}} \cap \Lambda[s]$, we find that

$$M_{\mathfrak{m}_{\mathfrak{x}}} = \left((M_{\mathfrak{m}_{\mathfrak{y}}})_{(s-c)}\right)_{\mathfrak{m}_{\mathfrak{x}}}.$$

Therefore we have $M_{\mathfrak{m}_{\mathfrak{x}}} = 0$, i.e., $(\mathfrak{q}_i)_{\mathfrak{m}_{\mathfrak{x}}} = (\mathfrak{Q}_{j(i)})_{\mathfrak{m}_{\mathfrak{x}}}$. Then Lemma 14.17 implies that $\mathfrak{q}_i = \mathfrak{Q}_{j(i)}$. Hence we obtain $\mathfrak{X}_i = \mathfrak{Y}_{j(i)}$ as schemes. This finishes the proof of Proposition 14.21 assuming Lemma 14.23.

Proof of Lemma 14.23. By Lemma 14.13, $\dim_\Lambda \frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i + (s-\mathfrak{s})}$ is finite. Since y_i acts on this ring by the multiplication by $\mathfrak{y}_i$, we have

$$\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i + (s-\mathfrak{s})}\right)_{\mathfrak{m}_{\mathfrak{y}}} = \frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i + (s-\mathfrak{s})}.$$

Hence the Krull dimension of this ring is 0. Since $\{\mathfrak{B}_1, \dots, \mathfrak{B}_n, s-c\}$ is a regular sequence, the Krull dimension of $\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i}\right)_{\mathfrak{m}_{\mathfrak{y}}}$ is 1. Since s is transcendental over Λ, we find that $\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i}\right)_{\mathfrak{m}_{\mathfrak{y}}}$ is a finitely generated $\Lambda[s]$-module. From the exact sequence

$$0 \to \left(\mathfrak{Q}_{j(i)}/\mathfrak{q}_i\right)_{\mathfrak{m}_{\mathfrak{y}}} \to \left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i}\right)_{\mathfrak{m}_{\mathfrak{y}}} \to \left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{Q}_j(i)}\right)_{\mathfrak{m}_{\mathfrak{y}}} \to 0,$$

we find that $M_{\mathfrak{m}_{\mathfrak{y}}} = \left(\mathfrak{Q}_{j(i)}/\mathfrak{q}_i\right)_{\mathfrak{m}_{\mathfrak{y}}}$ is a finitely generated $\Lambda[s]$-module. □

□

Corollary 14.24. *$I_0 = \overline{I}_0 = \overline{I}$. $\mathfrak{X} = Spec(\underline{\mathcal{R}})$.*

Proof. By Proposition 14.21 $I_0 = \bigcap_{j(i):i\in\mathfrak{I}} \mathfrak{Q}_{j(i)}$. On the other hand we have $\bigcap_{j(i):i\in\mathfrak{I}} \mathfrak{Q}_{j(i)} \supseteq \overline{I} = \overline{I}_0 \supseteq I_0$. Hence $\overline{I} = \bigcap_{j(i):i\in\mathfrak{I}} \mathfrak{Q}_{j(i)} = I_0$, as required. □

We are now in the position to complete the proof of Proposition 14.2. By Proposition 14.14, the coordinate ring $\underline{\mathcal{R}}$ of $\mathfrak{X}$ is a finitely generated $\Lambda[s]$ module. On the other hand, since $\Lambda[s]$ is a principal ideal domain, Proposition 14.15 implies that $\underline{\mathcal{R}}$ is a finitely generated free module over $\Lambda[s]$. In particular, we find that $\dim_\Lambda \underline{\mathcal{R}}/(s-0) = \dim_\Lambda \underline{\mathcal{R}}/(s-1)$. Therefore

$$\begin{aligned}\dim_\Lambda H(X;\Lambda) &= \dim_\Lambda \underline{\mathcal{R}}/(s-1) = \dim \underline{\mathcal{R}}/(s-0) \\ &= \dim_\Lambda \operatorname{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}\mathfrak{O}_{\mathfrak{b}})_\Lambda = \dim_\Lambda \operatorname{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}\mathfrak{O}_{\mathfrak{b}})_\Lambda.\end{aligned}$$

Here we use Lemma 12.8 to prove the last equality. □

In geometric terms, we may also put it as follows. We have (Theorems 9.9 III and 12.8 III [Ha])

$$\begin{aligned}\dim_\Lambda H(X;\Lambda) &= \dim_\Lambda H^0(\pi^{-1}(1);\mathfrak{O}_{\pi^{-1}(1)}) = \dim H^0_\Lambda(\pi^{-1}(0);\mathfrak{O}_{\pi^{-1}(0)}) \\ &= \dim_\Lambda \operatorname{PJac}_{\mathfrak{v}_T^P}(\mathfrak{P}\mathfrak{O}_{\mathfrak{b}})_\Lambda = \dim_\Lambda \operatorname{Jac}_{\mathfrak{v}_T^P}(\mathfrak{P}\mathfrak{O}_{\mathfrak{b}})_\Lambda.\end{aligned}$$

14.2. **The case $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$.** In this subsection, we generalize Theorem 14.1 to the case $\mathfrak{b} \in \mathcal{A}(\Lambda_0)$ and complete the proof of Theorem 2.32 (1). We choose any $\mathfrak{P}' \in \Lambda_0^P[y, y^{-1}]$ such that $\mathfrak{P}'$ is transformed to an element $\mathfrak{PO}_{\mathfrak{b}'}$ for some $\mathfrak{b}' \in \mathcal{A}(\Lambda_+)$. (Theorem 12.2 (2).) We have

$$\dim_\Lambda \mathrm{Jac}(\mathfrak{P}')_\Lambda = \sum_k \dim_\Lambda H^k(X;\Lambda)$$

by Theorem 14.1. Hence the next proposition completes the proof of Theorem 2.32 (1). □

Proposition 14.25.

$$\dim_\Lambda \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda = \dim_\Lambda \mathrm{Jac}(\mathfrak{P}')_\Lambda.$$

Remark 14.26. Before we start the proof of this proposition, we explain the idea of the argument. The idea is to reduce the problem for the Jacobian rings associated to the polytope P_ϵ, which is defined by

$$P_\epsilon = \{u \in \mathbb{R}^n \mid \ell_j(u) \geq \epsilon\}$$

as in Section 12. We explain the restriction of the potential function $\mathfrak{PO}_{\mathfrak{b}}$ from the interior of the polytope $\overset{\circ}{P}$ to the polytope P_ϵ.

Consider the following diagram:

$$\begin{tikzcd}
\Lambda_0[[Z_1,\dots,Z_m]] \arrow[r, "\phi"] \arrow[d, "\widetilde{j}"'] & \Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle \arrow[d, "j"'] \\
\Lambda_0\langle\langle Z_1^{(\epsilon)},\dots,Z_m^{(\epsilon)}\rangle\rangle \arrow[r, "\phi^{(\epsilon)}"] & \Lambda_0^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle
\end{tikzcd}$$

where ϕ and $\phi^{(\epsilon)}$ are the homomorphisms we have used:

$$\phi(Z_j) = T^{-\lambda_j} y_1^{v_{j,1}} \cdots y_n^{v_{j,n}}, \quad \phi^{(\epsilon)}(Z_j^{(\epsilon)}) = T^{-(\lambda_j+\epsilon)} y_1^{v_{j,1}} \cdots y_n^{v_{j,n}},$$

j is the inclusion, i.e., $j(y_i) = y_i$, and $\widetilde{j}(Z_i) = T^\epsilon Z_i^{(\epsilon)}$. It is straightforward to check the commutativity of the above diagram. We write $z_j = \phi(Z_j)$ and $z_j^{(\epsilon)} = \phi^{(\epsilon)}(Z_j^{(\epsilon)})$. Then we have

$$j(\mathcal{P}(z_1,\dots,z_m)) = \mathcal{P}(T^\epsilon z_1^{(\epsilon)},\dots,T^\epsilon z_m^{(\epsilon)}),$$

which belongs to $T^\epsilon \Lambda_0^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle$.

Proof of Proposition 14.25. By Lemma 10.5, we may put

$$\mathfrak{PO}_{\mathfrak{b}} \equiv c_1 z_1 + \cdots + c_m z_m + R(z_1,\dots,z_m) \mod \Lambda_+^P[y, y^{-1}] \tag{14.28}$$

where $R \in \Lambda_0[[Z_1,\dots,Z_m]]$ each summand of which has degree ≥ 2, and $c_i \in \mathbb{C} \setminus \{0\}$. We firstly take $\varepsilon > 0$ so that there is no critical point $\mathfrak{y}$ of $\mathfrak{PO}_{\mathfrak{b}}$ such that $\vec{\mathfrak{v}}_T(\mathfrak{y}) \in \mathrm{Int}P \setminus P_\epsilon$. By Theorem 12.2 we have a strict coordinate change $y'(y)$ which converge on P_ϵ such that

$$\mathfrak{PO}_{\mathfrak{b}}(y') = \mathfrak{P}(y) \in \Lambda_0^{P_\epsilon}[y, y^{-1}].$$

(Namely it is a Laurent polynomial.) Then by Proposition 2.15, we have

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})_\Lambda \cong \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}}\left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1,\dots,n\right)} \cong \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}}\left(y_i \frac{\partial \mathfrak{P}}{\partial y_i} : i = 1,\dots,n\right)}.$$

(Note we already proved Lemma 5.1 when we proved the surjectivity of Kodaira-Spencer map. Therefore the proof of Proposition 2.15 is already completed by now.) Moreover

$$\mathrm{Jac}(\mathfrak{P}')_\Lambda \cong \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}}\left(y_i\frac{\partial\mathfrak{P}'}{\partial y_i} : i = 1,\dots,n\right)}.$$

Thus it suffices to show that

$$\dim_\Lambda \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}}\left(y_i\frac{\partial\mathfrak{P}}{\partial y_i} : i = 1,\dots,n\right)} = \dim_\Lambda \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1}\rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}}\left(y_i\frac{\partial\mathfrak{P}'}{\partial y_i} : i = 1,\dots,n\right)}.$$

We define

$$\mathfrak{P}_{\mathrm{para}} = \alpha s\mathfrak{P} + (1-s)\mathfrak{P}' \in \Lambda_0^{P_\epsilon}[s, y, y^{-1}].$$

(Here $\alpha \in \mathbb{C} \setminus \{c_1,\dots,c_m,0\}$ is a constant.) In fact, we have

$$T^{-\epsilon}\mathfrak{P}_{\mathrm{para}} \in \Lambda_0^{P_\epsilon}[s, y, y^{-1}].$$

From now on we choose and take $\epsilon > 0$ so small so that the following holds: For each $s \in \mathbb{C}$ with $\alpha s c_i + (1-s) \neq 0$, $(i = 1,\dots,m)$ there is no solution of leading term equation of $\mathfrak{P}_{\mathrm{para}}$ on $\mathrm{Int}P \backslash P_\epsilon$. (See [FOOO5] Definitin 4.3 for the definition of leading term equation.) We also assume that the leading term equation of $\alpha\mathfrak{P} - \mathfrak{P}'$ also has no solution on $\mathrm{Int}P \setminus P_\epsilon$. (We use the assumption $\alpha \in \mathbb{C} \setminus \{c_1,\dots,c_m,0\}$ here.)

We put

$$\mathfrak{R} = \frac{\Lambda[s, y, y^{-1}]}{\left(y_i\frac{\partial\mathfrak{P}_{\mathrm{para}}}{\partial y_i} : i = 1,\dots,n\right)}, \qquad \underline{\mathfrak{R}} = \frac{\Lambda[s, y, y^{-1}]}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\varepsilon}}\left(y_i\frac{\partial\mathfrak{P}_{\mathrm{para}}}{\partial y_i} : i = 1,\dots,n\right)}.$$

We have natural morphisms

$$Spec(\underline{\mathfrak{R}}) \longrightarrow Spec(\mathfrak{R}) \xrightarrow{\pi} Spec(\Lambda[s]) = \Lambda.$$

If we put

$$z_i^\epsilon = T^{-\epsilon} z_i,$$

we derive from (14.28) that

$$T^{-\epsilon}\mathfrak{P}_{\mathrm{para}} \equiv \sum_{j=1}^m (\alpha s c_i + (1-s)) z_i^\epsilon \mod \Lambda_+^{P_\epsilon}[y, y^{-1}]. \tag{14.29}$$

Denote

$$\begin{aligned} B &= \{b_1,\dots,b_A\} = \{s \mid \exists s\ s\alpha c_i + (1-s) = 0\} \subset \mathbb{C},\\ B_+ &= \{b \in \Lambda \mid \exists i\ b \equiv b_i, \quad \mathrm{mod}\ \Lambda_+, i = 1,\dots,m\}, \quad B_+^c = \Lambda \setminus B_+. \end{aligned}$$

Let

$$I = \left(y_i\frac{\partial\mathfrak{P}_{\mathrm{para}}}{\partial y_i} : i = 1,\dots,n\right)$$

and let $\overline{I}$ be its closure with respect to the $\mathfrak{v}_T^{P_\epsilon}$-norm. We take their irredundant primary decompositions:

$$I = \bigcap_{i\in\mathfrak{I}_+} \mathfrak{q}_i, \qquad \overline{I} = \bigcap_{j\in\mathfrak{I}_+} \mathfrak{Q}_j. \tag{14.30}$$

We define

$$\begin{aligned} \mathfrak{I} &= \left\{ i \in \mathfrak{I}_+ \;\middle|\; \left| Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i} \right) \right| \cap |Spec(\underline{\mathfrak{R}})| \cap \pi^{-1}(B_+^c) \ne \emptyset \right\}, \\ \mathfrak{J} &= \left\{ j \in \mathfrak{J}_+ \;\middle|\; \left| Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{Q}_j} \right) \right| \cap \pi^{-1}(B_+^c) \ne \emptyset \right\}. \end{aligned} \tag{14.31}$$

We also put

$$\mathfrak{X}_i = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i} \right), \qquad \mathfrak{Y}_j = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{\mathfrak{Q}_j} \right).$$

$$I_0 = \bigcap_{i \in \mathfrak{I}} \mathfrak{q}_i, \qquad J_0 = \bigcap_{j \in \mathfrak{J}} \mathfrak{Q}_j, \tag{14.32}$$

and

$$\mathfrak{X} = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{I_0} \right), \qquad \mathfrak{Y} = Spec\left(\frac{\Lambda[s,y,y^{-1}]}{J_0} \right).$$

$\mathfrak{X}_i$ and $\mathfrak{Y}_j$ are irreducible components of $\mathfrak{X}$ and $\mathfrak{Y}$, respectively.

Lemma 14.27. *If* $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |\mathfrak{Y}|$ *and* $\mathfrak{s} \in B_+^c$, *then*

$$(\mathfrak{v}_T(\mathfrak{y}_1), \dots, \mathfrak{v}_T(\mathfrak{y}_n)) \in P_\varepsilon. \tag{14.33}$$

On the contrary if $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |\mathfrak{X}|$, $\mathfrak{s} \in B_+^c$ *and (14.33) is satisfied, then* $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |\mathfrak{Y}|$.

Proof. Let $\mathfrak{s} \in B_+^c$. Then we can show

$$\frac{\underline{\mathfrak{R}}}{(s - \mathfrak{s})} \cong \frac{\Lambda^{P_\epsilon}\langle\langle y, y^{-1} \rangle\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^{P_\epsilon}} \left(y_i \frac{\partial(\mathfrak{P}_{\mathrm{para}}|_{s=\mathfrak{s}})}{\partial y_i} : i = 1, \dots, \right)}, \tag{14.34}$$

in the same way as the proof of Lemma 14.4.

In fact, $\mathfrak{P}_{\mathrm{para}}|_{s=\mathfrak{s}} \equiv \sum(\alpha\mathfrak{s} - c_i)z_i \mod \Lambda_+$ and $\alpha s - c_i \notin \Lambda_+$ if $\mathfrak{s} \in B_+^c$. (In case $\mathfrak{s} \in \Lambda \setminus \Lambda_0$ we have $\mathfrak{P}_{\mathrm{para}}|_{s=\mathfrak{s}} \equiv T^{-\rho} \sum(\alpha - c_i)z_i \mod T^{-\rho}\Lambda_+^{P_\epsilon}\langle\langle y, y^{-1} \rangle\rangle$ and $\alpha \ne c_i$.)

The rest of the proof of the first half is the same as the proof of Lemma 14.6 using Lemma 12.8 (2) in place of Lemma 14.4. (See (14.29).)

Let $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |\mathfrak{X}|$, $\mathfrak{s} \in B_+^c$. Then $y_i \mapsto \mathfrak{y}_i$ and $s \mapsto \mathfrak{s}$ define a continuous homomorphism from $(\Lambda[y, y^-1, s], \mathfrak{v}_T^{P_\epsilon})$. Therefore $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |Spec(\underline{\mathfrak{R}})|$. Let $Spec(\Lambda[s,y,y^{-1}]/\mathfrak{Q}_j)$ be an irreducible component containing $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s})$. Since $\mathfrak{s} \in B_+^c$, $j \in \mathfrak{J}$. Therefore $(\mathfrak{y}_1, \dots, \mathfrak{y}_n, \mathfrak{s}) \in |\mathfrak{Y}|$ as required. □

We define a finite set $\mathcal{B}_1 \subset B_+$ as follows. Let $i \in \mathfrak{I} \setminus \mathfrak{I}_0$. Then,

$$|Spec(\Lambda[s,y,y^{-1}]/\mathfrak{q}_i)| \cap \pi^{-1}(B_+^c) \cap (\vec{\mathfrak{v}}_T)^{-1}(P_\epsilon) = \emptyset.$$

Therefore, by Lemma 14.33,

$$|Spec(\Lambda[s,y,y^{-1}]/\overline{\mathfrak{q}}_i)| \subset \pi^{-1}(B_+).$$

Since

$$\pi\left(|Spec(\Lambda[s,y,y^{-1}]/\overline{\mathfrak{q}}_i)|\right) \tag{14.35}$$

is a constructible set contained in B_+ by Chevalley's theorem (See page 42 [Mat]), it is easy to see that (14.35) is a finite set.

In the same way we find that if $j \in \mathfrak{J}_+ \setminus \mathfrak{J}$ then $\pi\left(|Spec(\Lambda[s,y,y^{-1}]/\mathfrak{Q}_j)|\right)$ is a finite subset of B_+.

We define

$$\mathcal{B}_1 = \bigcup_{i \in \mathfrak{I}_+ \setminus \mathfrak{I}} \pi\left(|Spec(\Lambda[s,y,y^{-1}]/\overline{\mathfrak{q}}_i)|\right) \cup \bigcup_{j \in \mathfrak{I}_+ \setminus \mathfrak{I}} \pi\left(|Spec(\Lambda[s,y,y^{-1}]/\mathfrak{Q}_j)|\right). \tag{14.36}$$

Hereafter we denote by $A_{\mathcal{B}_1}$ the localization of A by the multiplicative set generated by $\{s - \mathfrak{s} \mid \mathfrak{s} \in \mathcal{B}_1\}$.

Corollary 14.28. *We have*

$$(\overline{I}_0)_{\mathcal{B}_1} = (\overline{I})_{\mathcal{B}_1} = (J_0)_{\mathcal{B}_1},$$

where $\overline{I}_0$ is the closure of I_0.

Proof. The proof is similar to the proof of Corollary 14.7. If $j \in \mathfrak{I}_+ \setminus \mathfrak{I}$, then $1 \in (\mathfrak{Q}_j)_{\mathcal{B}_1}$. (This is because $|Spec(\Lambda[s,y,y^{-1}]/\mathfrak{Q}_j)| \cap \pi^{-1}(\Lambda \backslash \mathcal{B}_1) = \emptyset$.) This implies $(\overline{I})_{\mathcal{B}_1} = (J_0)_{\mathcal{B}_1}$. If $i \in \mathfrak{I}_+ \setminus \mathfrak{I}$, then $1 \in (\overline{\mathfrak{q}}_i)_{\mathcal{B}_1}$. This implies $(\overline{I}_0)_{\mathcal{B}_1} = (\overline{I})_{\mathcal{B}_1}$. □

Proposition 14.29. *For each $i \in \mathfrak{I}$ there exists $j \in \mathfrak{I}$ such that $|\mathfrak{X}_i| \subseteq |\mathfrak{Y}_j|$.*

Proof. Let $\mathfrak{Z}$ be one of the irreducible components of $\mathfrak{X}_i \cap \underline{\mathfrak{Y}}$. We may assume

$$|\mathfrak{Z}| \cap (\vec{\mathfrak{v}}_T)^{-1}(P_\epsilon) \cap \pi^{-1}(B_+^c) \neq \emptyset.$$

In fact,

$$|\mathfrak{X}_i| \cap (\vec{\mathfrak{v}}_T)^{-1}(P_\epsilon) \cap \pi^{-1}(B_+^c) \neq \emptyset, \tag{14.37}$$

by definition. The point on (14.37) is also contained in $\mathfrak{Y}$ by Lemma 14.27. Thus we take $\mathfrak{Z}$ which contains it.

We will prove the coincidence of geometric points in $\mathfrak{X}_i$ and $\mathfrak{Z}$. To prove this we prove an analogue of Lemma 14.9 with an additional condition that the point corresponding to $\overline{\mathfrak{f}}$ is contained in $\pi^{-1}(B_+^c)$. We can prove it in the following way. Assume $|\mathfrak{Z}| \neq |\mathfrak{X}_i|$. We consider the set of points $\mathfrak{y} \in |\mathfrak{Z}|$ such that there exists a 1-dimensional irreducible and reduced scheme $\mathfrak{C} \subset (\mathfrak{X}_i)_{\mathrm{red}}$, which is a subscheme of $Spec(\mathfrak{R})$, with $\dim(\mathfrak{C} \cap \mathfrak{Z}_{\mathrm{red}}) = 0$ and $\mathfrak{y} \in |\mathfrak{C} \cap \mathfrak{Z}_{\mathrm{red}}|$. We can easily see that this set is Zariski open in $|\mathfrak{Z}|$. Therefore there exists such $\mathfrak{y}$ in $\pi^{-1}(B_+^c)$. The rest of the proof is the same as the proof of Lemma 14.9.

Now we observe that for $\mathfrak{s} = \pi(\mathfrak{y})$ where $\mathfrak{y}$ corresponds to $\overline{\mathfrak{f}}$ we have

$$T^{-\epsilon}\mathfrak{P}_{\mathfrak{s}} \equiv \sum_{j=1}^{m} (\alpha\mathfrak{s}c_i + (1-\mathfrak{s}))z_i^{\epsilon} \mod \Lambda_+^{P_\epsilon}[y,y^{-1}]$$

and $\alpha\mathfrak{s}c_i + (1-\mathfrak{s}) \notin \Lambda_+$. We can use this fact and proceed in the same way as the proof of Proposition 14.8 to prove Proposition 14.29. □

Lemma 14.30. *If $i \in \mathfrak{I}_0$, then for any $\mathfrak{s} \in B_+^c$ we have:*

$$\operatorname{rank}_\Lambda \frac{\Lambda[s,y,y^{-1}]}{\mathfrak{q}_i(s-\mathfrak{s})} < \infty.$$

Proof. By Proposition 14.29 it suffices to show that $\mathfrak{Y} \cap \pi^{-1}(\mathfrak{s})$ is zero dimensional. For this purpose it suffices to show that

$$\dim_\Lambda \mathrm{Jac}(T^{-\epsilon}\mathfrak{P}_{\mathfrak{s}}) < \infty.$$

This follows from (14.29), Theorem 10.1 and Theorem 11.1 (2). □

Corollary 14.31. *If $i \in \mathfrak{I}_0$, then $\dim \mathfrak{X}_i \leq 1$.*

This is immediate from Lemma 14.30.

Proposition 14.32. *There exists a finite subset* $\mathcal{B}_2 \supset \mathcal{B}_1$ *of* B_+ *such that*

$$\mathfrak{X} \cap \pi^{-1}(\Lambda \setminus \mathcal{B}_2) \to \Lambda \setminus \mathcal{B}_2$$

is projective.

Proof. We consider the scheme $\prod_{i=1}^n \mathbb{P}^1_{y_i} \times Spec(\Lambda[s])$. Let $\overline{\mathfrak{X}}$ be the closure of $\mathfrak{X}$ in $\prod_{i=1}^n \mathbb{P}^1_{y_i} \times Spec(\Lambda[s])$. We put

$$D = \left\{ (y_1, \cdots : y_n, s) \in \left| \prod_{i=1}^n \mathbb{P}^1_{y_i} \times Spec(\Lambda[s]) \right| \middle| \exists i \ y_i \in \{0, \infty\} \right\}.$$

By Corollary 14.31 the intersection $|\overline{\mathfrak{X}}| \cap D$ is a finite set. We put

$$\mathcal{B}_2 = \pi(|\overline{\mathfrak{X}}| \cap D) \cup \mathcal{B}_1 \subset \Lambda.$$

We can show that $\mathcal{B}_2 \subset B_+$ in the same way as the proof of Proposition 14.14. □

Proposition 14.33. *There exists a finite subset* $\mathcal{B}_3 \supset \mathcal{B}_2$ *of* B_+ *such that the following holds. We denote* $\mathcal{B}_3^c = \Lambda \setminus \mathcal{B}_3$.

(1) $\dim \mathfrak{X} = 1$.
(2) $\pi : \mathfrak{X} \cap \pi^{-1}(\mathcal{B}_3^c) \to \mathcal{B}_3^c$ *is flat.*
(3) *Let* $\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}$ *be the maximal ideal corresponding to* $(\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}| \cap \pi^{-1}(\mathcal{B}_3^c)$. *Then* $y_1 \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_1}, \dots, y_n \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_n}$ *is a regular sequence in* $\Lambda[s, y, y^{-1}]_{\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}}$.

Proof. We put

$$|\mathfrak{X}'| = \bigcup_{i \in \mathfrak{I}_+ \setminus \mathfrak{I}} \left| Spec\left(\frac{\Lambda[s, y, y^{-1}]}{\mathfrak{q}_i} \right) \right|.$$

Since $\dim \mathfrak{X} \le 1$, the intersection $|\mathfrak{X}'| \cap |\mathfrak{X}|$ is a finite set. We put

$$\mathcal{B}_3 = \mathcal{B}_2 \cup \pi(|\mathfrak{X}'| \cap |\mathfrak{X}|).$$

Then we can prove (3) in the same way as the proof of Proposition 14.15 (3) as follows. Let $(\mathfrak{y}, \mathfrak{s}) \in |\mathfrak{X}| \cap \pi^{-1}(\mathcal{B}_3^c)$, and $\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}$ the corresponding ideal of $\Lambda[s, y, y^{-1}]$. We find that $I\Lambda[s, y, y^{-1}]_{\mathfrak{m}_\mathfrak{x}} = I_0\Lambda[s, y, y^{-1}]_{\mathfrak{m}_\mathfrak{x}}$. Therefore Lemma 14.30 implies that $y_1 \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_1}, \dots, y_n \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_n}, s - \mathfrak{s}$ is a regular sequence of $\mathfrak{m}_{(\mathfrak{y},\mathfrak{s})}$. (3) follows.

We can then prove (2) and (1) in the same way as Proposition 14.15. □

Proposition 14.34. *For any* $i \in \mathfrak{I}$ *we have the following.*

(1) $\dim \mathfrak{X}_i = 1$.
(2) $\pi : \mathfrak{X}_i \cap \pi^{-1}(\mathcal{B}_3^c) \to \mathcal{B}_3^c$ *is flat.*

Proof. Proposition 14.34 follows from Proposition 14.33, Lemmas 14.17, 14.18 and Theorem 14.19 in the same way as the proof of Proposition 14.16. □

Lemma 14.35. *In the situation of Proposition 14.29, we have* $|\mathfrak{X}_i| = |\mathfrak{Y}_j|$.

The proof is by the same argument as in Lemma 14.20. For each $i \in \mathfrak{I}$ we take $j(i)$ such that $\mathfrak{X}_i$ and $\mathfrak{Y}_{j(i)}$ are as in Proposition 14.29.

Proposition 14.36. *We have* $\mathfrak{X}_i = \mathfrak{Y}_{j(i)}$, *i.e.,* $\mathfrak{q}_i = \mathfrak{Q}_{j(i)}$.

Proof. The proof is similar to the proof of Proposition 14.21.

We put

$$B_1 = \bigcup_{j\in\mathfrak{J}_+\setminus\{j(i)\mid i\in\mathfrak{I}\}} \pi(|\mathfrak{Y}_j|)$$

and

$$B_2 = \bigcup_{i_1,\neq i_2; i_1,i_2\in\mathfrak{I}} \pi(|\mathfrak{X}_{i_1}|\cap|\mathfrak{X}_{i_j}|).$$

They are finite sets. We take $c \in \Lambda \setminus \pi(B_1 \cup B_2) \setminus B_+$.

We take $\mathfrak{x} = (c, \mathfrak{y}) \in |\mathfrak{X}_i| \cap \pi^{-1}(c)$. Let $\mathfrak{m}_{\mathfrak{y}}$ be the maximal ideal of $\Lambda[y, y^{-1}]$ corresponding to $\mathfrak{y}$. By the same argument as in Proposition 14.21 we can show

$$(\mathfrak{R}/(s-c))_{\mathfrak{m}_y} \cong (\underline{\mathfrak{R}}/(s-c))_{\mathfrak{m}_y}.$$

We can now apply the proof of Proposition 14.21 to obtain

$$\mathfrak{R}_{\mathfrak{m}_{\mathfrak{x}}} = \underline{\mathfrak{R}}_{\mathfrak{m}_{\mathfrak{x}}}. \tag{14.38}$$

Here $\mathfrak{m}_{\mathfrak{x}}$ is the maximal ideal of $\Lambda[s, y, y^{-1}]$ corresponding to $\mathfrak{x}$.

In fact we remark that (14.38) is an equality of the ideals which are localized at $\mathfrak{x}$. Also the argument of this equality in the proof of Proposition 14.21 works only on those local rings and their quotients. Therefore the proof goes without change, using Propositions 14.33 and 14.34.

Once (14.38) is proved, we have $\mathfrak{q}_i = \mathfrak{Q}_{j(i)}$ by Lemma 14.17. Proposition 14.36 follows. □

Corollary 14.37. *$\mathfrak{X} = \mathfrak{Y}$.*

The proof is the same as one of Corollary 14.24 using Proposition 14.36. Now we obtain

$$\begin{aligned}\dim_\Lambda \mathrm{PJac}(\mathfrak{P}')_\Lambda &= \dim_\Lambda \mathrm{Jac}(\mathfrak{P}')_\Lambda = \dim_\Lambda H^0(\pi^{-1}(1); \mathfrak{O}_{\pi^{-1}(1)})\\ &= \dim H^0_\Lambda(\pi^{-1}(0); \mathfrak{O}_{\pi^{-1}(0)}) = \dim_\Lambda \mathrm{PJac}(\mathfrak{P})_\Lambda = \dim_\Lambda \mathrm{Jac}(\mathfrak{P})_\Lambda.\end{aligned}$$

Hence the proof of Proposition 14.25. □

Remark 14.38. Comparing with the argument for the case of $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$, we replace P by P_ϵ in the above argument. In fact, in our situation, there may be a solution of $y_k \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_k} = 0$ whose valuation lies on ∂P. Moreover $\mathfrak{PO}_{\mathfrak{b}} \equiv z_1 + \cdots + z_m$ mod $T^{\lambda_1}\Lambda_0\langle\!\langle s, y, y(u)^{-1}\rangle\!\rangle$ may not be true. (See Example 14.39.) By this reason we prove algebraization of $\mathfrak{PO}_{\mathfrak{b}}$ by a coordiante transformaion converging on P_ε only. We can easily prove that there exists a neighborhood U of ∂P such that there exists no solution of $y_k \frac{\partial \mathfrak{P}_{\rm para}}{\partial y_k} = 0$ whose valuation lies on $U \cap \mathrm{Int}P$. In fact, if $u \in U \cap \mathrm{Int}P$, then $\mathfrak{P}_{\rm para} \equiv z_1 + \cdots + z_m \mod T^{\lambda_1}\Lambda_0[s, y(u), y(u)^{-1}]$. We used this fact to find P_ϵ.

Example 14.39. Let $X = \mathbb{C}P^2$ and $D = \{(0,0)\} \subset \mathbb{C}^2 \subset \mathbb{C}P^2$. We put $\mathfrak{b} = cPD[D]$, $c \in \mathbb{C} \setminus \{0\}$. Then we have

$$\mathfrak{PO}_{\mathfrak{b}} = y_1 + y_2 + cy_1y_2 + Ty_1^{-1}y_2^{-1}.$$

Because of the third term $\mathfrak{PO}_{\mathfrak{b}} \equiv z_1 + z_2 + z_3 \mod T^{\lambda_1}\Lambda_0^P[y, y^{-1}]$ is *not* true.

In this example, there is a critical point $(\mathfrak{y}_1, \mathfrak{y}_2) = (-c, -c)$ whose valuation is $(0, 0) \in \partial P$.

We remark that $y_i \mapsto \mathfrak{y}_i$ defines a homomorphism

$$\frac{\Lambda^P\langle\!\langle y, y^{-1}\rangle\!\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(y_1\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_1}, y_2\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_2}\right)} \to \Lambda.$$

It follows that

$$\mathrm{rank}_{\Lambda}\frac{\Lambda^P\langle\!\langle y, y^{-1}\rangle\!\rangle}{\mathrm{Clos}_{\mathfrak{v}_T^P}\left(y_1\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_1}, y_2\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_2}\right)} = 4 > 3 = \mathrm{rank}_{\mathbb{Q}} H(\mathbb{C}P^2;\mathbb{Q}).$$

On the other hand,

$$\mathrm{rank}_{\Lambda}\frac{\Lambda^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_1\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_1}, y_2\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_2}\right)} = 3.$$

In fact, $\sum_{k=1}^{\infty} y_1^k$ converges in $d^{\overset{\circ}{P}}$-topology. Therefore $y_i \mapsto \mathfrak{y}_i$ does not define a homomorphism

$$\frac{\Lambda^{\overset{\circ}{P}}\langle\!\langle y, y^{-1}\rangle\!\rangle}{\mathrm{Clos}_{d_{\overset{\circ}{P}}}\left(y_1\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_1}, y_2\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_2}\right)} \to \Lambda.$$

15. THE CHERN CLASS c_1 AND CRITICAL VALUES OF $\mathfrak{PO}_{\mathfrak{b}}$

In this section we prove Theorem 1.4. We regard $\mathfrak{PO}_{\mathfrak{b}}$ as an element in $\Lambda_0\langle\!\langle y, y^{-1}\rangle\!\rangle$.

Proposition 15.1. *If we assume* $\mathfrak{b} \in H^2(X;\Lambda_0)$*, then we have*

$$\mathfrak{ks}_{\mathfrak{b}}(c_1(X)) \equiv \mathfrak{PO}_{\mathfrak{b}} \mod \left(y_i\frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1, \dots, n\right).$$

Proof. We decompose $b = b_0 + x \in H^1(L(u);\Lambda_0)$ where $\mathfrak{v}_T(b_0) = 0$ and $\mathfrak{v}_T(x) > 0$. Then, by definition, we have

$$\begin{aligned} &\mathfrak{PO}_{\mathfrak{b}}(y_1, \dots, y_n) PD[L(u)] \\ = &\sum_{k=0}^{\infty}\sum_{\beta} T^{\omega\cap\beta/2\pi}\mathfrak{m}_{k,\beta}^{\mathfrak{b},\rho}(b, \dots, b) \\ = &\sum_{k=0}^{\infty}\sum_{\beta} \exp(\mathfrak{b}\cap\beta) T^{\omega\cap\beta/2\pi}\rho^{b_0}(\partial\beta)\mathfrak{m}_{k,\beta}(x, \dots, x). \end{aligned} \tag{15.1}$$

Here we use the assumption $\mathfrak{b} \in H^2(X;\Lambda_0)$. It is well known (see the Lemma in p. 109 of [Ful]) that the first Chern class $c_1(X)$ can be expressed as

$$c_1(X) = \sum_{i=1}^{m} PD[D_{\mathbf{p}_i}].$$

By a dimension counting argument using the choice $\mathfrak{b} \in H^2(X;\Lambda_0)$, we derive that only the classes $\beta \in H_2(X, L(u);\mathbb{Z})$ in (15.1) satisfying $\mu(\beta) = 2$ nontrivially contribute in the sum of the right hand side. Therefore, by the Maslov index formula and the classification theorem of holomorphic discs stated in p.781 of [CO], we obtain $c_1(X) \cap \beta = 1$.

Regard $c_1(X)$ as a constant vector field on $H(X;\Lambda_0)$. Then, by the definition of $\mathfrak{ks}_{\mathfrak{b}}$ given in Definition 2.30, we have

$$\mathfrak{ks}_{\mathfrak{b}}(c_1(X)) = c_1(X)[\mathfrak{PO}_{\mathfrak{b}}] \mod \left(y_i \frac{\partial \mathfrak{PO}_{\mathfrak{b}}}{\partial y_i} : i = 1, \dots, n \right) \tag{15.2}$$

where $c_1(X)[\mathfrak{PO}_{\mathfrak{b}}]$ denotes the directional derivative of the potential function $\mathfrak{PO}_{\mathfrak{b}}$ with respect to the vector field $c_1(X)$. Noting that only the term $\exp(\mathfrak{b}\cap\beta)$ depends on the variable $\mathfrak{b}$ and

$$c_1(X)[\exp(\mathfrak{b}\cap\beta)] = \sum_{i=1}^m PD[D_{\mathbf{p}_i}][\exp(\mathfrak{b}\cap\beta)] = \left(\sum_{i=1}^m PD[D_{\mathbf{p}_i}][\mathfrak{b}\cap\beta]\right)\exp(\mathfrak{b}\cap\beta).$$

A straightforward computation, or the formula

$$(\vec{v}[x_1], \dots, \vec{v}[x_n]) = (v_1, \dots, v_n) \quad \text{for } \vec{v} = (v_1, \dots, v_n),$$

gives rise to

$$PD[D_{\mathbf{p}_i}][\mathfrak{b}\cap\beta] = [D_{\mathbf{p}_i}]\cap\beta$$

and hence

$$c_1(X)[\exp(\mathfrak{b}\cap\beta)] = (c_1(X)\cap\beta)\exp(\mathfrak{b}\cap\beta) = \exp(\mathfrak{b}\cap\beta).$$

Substituting this into (15.2) and (15.1), the proposition follows. □

Theorem 1.4 follows from Proposition 15.1 and Theorem 1.1 (1) by standard commutative algebra. □

16. Hirzebruch surface F_2: an example

In this section, we study an example using Theorem 1.4. Consider the Hirzebruch surface $F_2(\alpha)$ whose moment polytope is given by

$$P_\alpha = \{(u_1, u_2) \mid 0 < u_1, u_2,\ u_1 + 2u_1 < 2,\ u_2 < 1 - \alpha\}. \tag{16.1}$$

We put

$$\begin{aligned}&\partial_1 P = \{(u_1,u_2)\in P \mid u_1 = 0\}, \quad \partial_2 P = \{(u_1,u_2)\in P \mid u_2 = 0\},\\ &\partial_3 P = \{(u_1,u_2)\in P \mid u_1 + 2u_2 = 2\}, \quad \partial_4 P = \{(u_1,u_2)\in P \mid u_2 = 1-\alpha\},\end{aligned}$$

and $D_i = \pi^{-1}(\partial_i P)$, $i = 1,2,3,4$. We put

$$\mathfrak{b} = \sum_{i=1}^4 w_i PD([D_i]) \in H^2(F_2(\alpha);\Lambda_0). \tag{16.2}$$

Theorem 16.1. *The potential function of $F_2(\alpha)$ is given by*

$$\begin{aligned}\mathfrak{PO}^u_{\mathfrak{b}} = T^{u_1}e^{w_1}y_1 + T^{u_2}e^{w_2}y_2 + T^{2-u_1-2u_2}e^{w_3}y_1^{-1}y_2^{-2}&\\ + T^{1-\alpha-u_2}e^{w_4}(1+T^{2\alpha})y_2^{-1}.&\end{aligned} \tag{16.3}$$

Remark 16.2. For $\mathfrak{b} = 0$, the leading order potential function, which is defined and denoted by $\mathfrak{PO}_0$ in [FOOO4] (4.9) is given by

$$T^{u_1}y_1 + T^{u_2}y_2 + T^{2-u_1-2u_2}y_1^{-1}y_2^{-2} + T^{1-\alpha-u_2}y_2^{-1}.$$

We note that this is different from the full potential function $\mathfrak{PO}^u_{\mathfrak{b}}$ with $\mathfrak{b} = \mathbf{0}$ for the case of $F_2(\alpha)$.

Remark 16.3. Theorem 16.1 was previously proved by D. Auroux [Aur2] Proposition 3.1 by a different method.

Theorem 16.1 is [FOOO7] Theorem 2.2. We reproduce the proof below for reader's convenience.

Proof. We first prove the following lemma.

Lemma 16.4. *There exists $c \in \Lambda_+$ such that*

$$\begin{aligned}\mathfrak{PO}^u_{\mathfrak b} = T^{u_1}e^{w_1}y_1 + T^{u_2}e^{w_2}y_2 + T^{2-u_1-2u_2}e^{w_3}y_1^{-1}y_2^{-2}\\ + T^{1-\alpha-u_2}e^{w_4}(1+c)y_2^{-1}.\end{aligned} \tag{16.4}$$

Proof. Since $\deg \mathfrak b = 2$, only the moduli spaces $\mathcal{M}_{k+1}(\beta)$ with $\mu(\beta) = 2$ contribute to $\mathfrak{PO}^u_{\mathfrak b}$. (See (6.9) [FOOO5].) We remark that $c_1(D_i) > 0$ for $i = 1, 2, 3$ and $c_1(D_4) = 0$. Suppose $\mathcal{M}_{k+1}(\beta) \neq \emptyset$. Then by [FOOO4] Theorem 11.1 there exist $k_i \geq 0$ and $\ell_j \geq 0$ with $\sum k_i > 0$ such that

$$\beta = \sum_{i=1}^{4} k_i\beta_i + \sum_{j=1}^{4} \ell_j D_j.$$

Here $\beta_i \in H_2(X, L(u); \mathbb{Z})$ with $\beta_i \cap D_j = 1$ $(i = j)$, $\beta_i \cap D_j = 0$ $(i \neq j)$. We remark $\mu(\beta_i) = 2$. Therefore, using $\mu(\beta) = 2$, we find that $\sum k_i = 1$ and $\ell_j = 0$ for $j \neq 4$. Since β must be represented by a connected genus zero bordered curve, it is easy to see that $k_4 = 1$ if $\ell_4 \neq 0$. Thus only the classes β_i $(i = 1, 2, 3, 4)$ and $\beta_4 + \ell D_4$ contribute. The lemma follows easily. □

Thus it suffices to show $c = T^{2\alpha}$. (We remark that $2\alpha = D_4 \cap \omega/2\pi$. See [CO] Theorem 8.1 or rather its proof.)

Let $S^2(A)$ be a symplectic 2 sphere with area A.

Lemma 16.5. *$F_2(\alpha)$ is symplectomorphic to $S^2(1-\alpha) \times S^2(1+\alpha)$.*

This is proved as Proposition 4.1 in [FOOO7].

The moment polytope of $S^2(1-\alpha) \times S^2(1+\alpha)$ is

$$P'(\alpha) = \{(u_1, u_2) \mid 0 < u_1 < 1-\alpha,\ 0 < u_2 < 1+\alpha\}.$$

There is a unique balanced fiber $u = ((1-\alpha)/2, (1+\alpha)/2)$, whose potential function is

$$T^{(1-\alpha)/2}(y_1 + y_1^{-1}) + T^{(1+\alpha)/2}(y_2 + y_2^{-1}). \tag{16.5}$$

It has 4 critical points $y_1, y_2 = \pm 1$ with the corresponding critical values given by

$$\pm 2T^{(1-\alpha)/2}(1 \pm T^{\alpha}) \tag{16.6}$$

respectively.

On the other hand, the balanced fiber of $F_2(\alpha)$ is located at $((1+\alpha)/2, (1-\alpha)/2)$, which has the potential function

$$\mathfrak{PO}^u = T^{(1-\alpha)/2}(y_2 + (1+c)y_2^{-1}) + T^{(1+\alpha)/2}(y_1 + y_1^{-1}y_2^{-2}). \tag{16.7}$$

The critical point equation for $\mathfrak{PO}^u$ is

$$0 = 1 - y_1^{-2}y_2^{-2}. \tag{16.8}$$

$$0 = 1 - 2T^{\alpha}y_1^{-1}y_2^{-3} - (1+c)y_2^{-2}. \tag{16.9}$$

The equation (16.8) implies $y_1y_2 = \pm 1$. We consider the two cases separately.

Case 1: $y_1y_2 = -1$. Equation (16.9) implies $y_2^2 = 1 + c - 2T^{\alpha}$. Then the corresponding critical values are

$$\pm\, 2T^{(1-\alpha)/2}\sqrt{1+c-2T^{\alpha}}. \tag{16.10}$$

Case 2: $y_1y_2 = 1$. (16.9) implies $y_2^2 = 1 + c + 2T^{\alpha}$. Then the critical values are

$$\pm\, 2T^{(1-\alpha)/2}\sqrt{1+c+2T^{\alpha}}. \tag{16.11}$$

Thus we must have

$$1 \pm T^{\alpha} = \sqrt{1+c\pm 2T^{\alpha}}$$

from which $c = T^{2\alpha}$ immediately follows. The proof of Theorem 16.1 is complete.

$\square$

We have thus calculated the potential function of $F_2(\alpha)$ using Theorem 1.4. (We use Theorem 1.1 (1) but not (2) in the proof of Theorem 1.4.) Below we check Theorem 1.1 (2) for the case of $F_2(\alpha)$. For simplicity we consider the case $\mathfrak{b} = \mathbf{0}$. We remark that $F_2(\alpha)$ and $S^2(1-\alpha)\times S^2(1+\alpha)$ are both nef. Therefore it follows from Theorem 2.24 (3) that the residue pairing is given by the reciprocal of the determinant of the Hessian matrix.

We first calculate the Hessian determinant of the potential function (16.5) of $S^2(1-\alpha)\times S^2(1+\alpha)$ at the critical points $y_1 = \pm 1$, $y_2 = \pm 1$. (We take $x_i = \log y_i$ as the coordinate.) It is easy to see that the Hessian matrix is diagonal with entries

$$\left(y_1\frac{\partial}{\partial y_1}\right)^2(\mathfrak{PO}) = T^{(1-\alpha)/2}(y_1+y_1^{-1}),\left(y_2\frac{\partial}{\partial y_2}\right)^2(\mathfrak{PO}) = T^{(1+\alpha)/2}(y_2+y_2^{-1}).$$

Therefore its determinant is:

$$\pm\, 4T. \tag{16.12}$$

We next consider the case of $F_2(\alpha)$. The Hessian matrix of (16.3) is

$$\begin{bmatrix} T^{(1+\alpha)/2}(y_1+y_1^{-1}y_2^{-2}) & 2T^{(1+\alpha)/2}y_1^{-1}y_2^{-2} \\ 2T^{(1+\alpha)/2}y_1^{-1}y_2^{-2} & T^{(1-\alpha)/2}(y_2+(1+T^{2\alpha})y_2^{-1}) + 4T^{(1+\alpha)/2}y_1^{-1}y_2^{-2} \end{bmatrix}$$

For the case $y_1y_2 = -1$, $y_2 = \pm(1-T^{\alpha})$, the determinant of this matrix is

$$-\frac{2T^{(1+\alpha)/2}}{y_2}\cdot\frac{2T^{(1-\alpha)/2}(1-3T^{\alpha}+T^{2\alpha})}{y_2} - \frac{4T^{1+\alpha}}{y_2^2} = \mp 4T.$$

For the case $y_1y_2 = 1$, $y_2 = \pm(1+T^{\alpha})$, the determinant of this matrix is

$$\frac{2T^{(1+\alpha)/2}}{y_2}\cdot\frac{2T^{(1-\alpha)/2}(1+3T^{\alpha}+T^{2\alpha})}{y_2} - \frac{4T^{1+\alpha}}{y_2^2} = \pm 4T.$$

Thus they coincide with (16.12).

In other words, the Poincaré duality pairing between the units of the factors of the decomposition of $QH(F_2(\alpha);\Lambda) \cong \Lambda^4$ gives rise to $1/4T, 1/4T, -1/4T$ and $-1/4T$ respectively.

We have thus illustrated Theorem 1.1 (2) in this case.

Part 2. Coincidence of pairings

17. OPERATOR $\mathfrak{p}$ AND POINCARÉ DUALITY

Sections 17 - 21 are devoted to the proof of Theorem 2.32 (2). In Sections 17, 19 we construct a homomorphism

$$i_{\#,\mathrm{qm},(\mathfrak{b},b,u)} : HF((L(u),\mathfrak{b},b);(L(u),\mathfrak{b},b);\Lambda_0) \to H(X;\Lambda_0). \tag{17.1}$$

We now explain how the $\mathfrak{ks}_{\mathfrak{b}}$ is related to a homomorphism

$$i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)} : H(X;\Lambda_0) \to HF((L(u),\mathfrak{b},b);(L(u),\mathfrak{b},b);\Lambda_0). \tag{17.2}$$

In case $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function, we have a splitting

$$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda \cong \prod_{\mathfrak{y}\in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})} \Lambda 1_{\mathfrak{y}}.$$

See Proposition 2.15 and Definition 2.16. We remark that the proof of Proposition 2.15 is completed at this stage since the proof of Theorem 2.32 (1) is completed in Section 14. Each critical point $\mathfrak{y}$ corresponds to a pair (u,b) such that $HF((L(u),\mathfrak{b},b);(L(u),\mathfrak{b},b);\Lambda_0) \cong H(T^n;\Lambda_0)$. (Theorem 3.12 [FOOO5].) So we identify $\mathfrak{y}$ to such (u,b) and regard $\mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$ as the set of such (u,b)'s. We write $1_{(u,b)}$ in place of $1_{\mathfrak{y}}$. We now decompose $\mathfrak{ks}_{\mathfrak{b}}(Q) \in \Lambda$ by

$$\mathfrak{ks}_{\mathfrak{b}}(Q) = \sum_{(u,b)\in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})} \mathfrak{ks}_{\mathfrak{b};(u,b)}(Q) 1_{(u,b)}. \tag{17.3}$$

Lemma 17.1.

$$i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q) = \mathfrak{ks}_{\mathfrak{b};(u,b)}(Q)\cdot PD([L(u)]). \tag{17.4}$$

This lemma actually is a direct consequence of the definition. We do not give its proof here since we do not need it. The reader may regard the right hand side of (17.4) as the definition of $i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}$. We are going to construct the adjoint operator $i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}$ of $i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}$, i.e., the operator satisfying

$$\langle i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q), P\rangle_{\mathrm{PD}_{L(u)}} = \langle Q, i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(P)\rangle_{\mathrm{PD}_X}. \tag{17.5}$$

Here $\langle\cdot,\cdot\rangle_{\mathrm{PD}_{L(u)}}$ and $\langle\cdot,\cdot\rangle_{\mathrm{PD}_X}$ are Poincaré duality pairings on $L(u)$ and X respectively, and $Q \in H(X;\Lambda_0)$ and $P \in H(L(u);\Lambda_0) \cong HF((L(u),\mathfrak{b},b);(L(u),\mathfrak{b},b);\Lambda_0)$.

For the proof of Theorem 2.32 (2) we need to realize the operator $i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}$ geometrically. We use the operator $\mathfrak{p}$, which was introduced in Sections 3.8 and 7.4 [FOOO3], for this purpose. We review it in this section. Let C be a graded free $\Lambda_{0,\mathrm{nov}}$ module. Here $\Lambda_{0,\mathrm{nov}}$ is the universal Novikov ring:

$$\Lambda_{0,\mathrm{nov}} = \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} e^{n_i} \in \Lambda_{\mathrm{nov}} \;\middle|\; \lambda_i \ge 0, \lim_{i\to\infty} \lambda_i = \infty, a_i \in R \right\}.$$

($\deg e = 2$.) We define its degree shift $C[1]$ by $C[1]^k = C^{k+1}$. We define $B_k(C[1])$, $\widehat{B}(C[1])$ as in Section 6. We also consider a map $\Delta^{k-1} : BC \to (BC)^{\otimes k}$ as in Section 6 and use the expression:

$$\Delta^{k-1}(\mathbf{x}) = \sum_c \mathbf{x}_c^{k;1} \otimes \cdots \otimes \mathbf{x}_c^{k;k}.$$

We define $\mathrm{cyc} : B_k(C[1]) \to B_k(C[1])$ by

$$\mathrm{cyc}(x_1 \otimes \cdots \otimes x_k) = (-1)^{\deg' x_k \times (\sum_{i=1}^{k-1} \deg' x_i)} x_k \otimes x_1 \otimes \cdots \otimes x_{k-1}. \tag{17.6}$$

Here and hereafter we put $\deg' x = \deg x - 1$. We also put $\deg'(x_1 \otimes \cdots \otimes x_k) = \sum_{i=1}^{k} \deg' x_i$.

Definition 17.2. $B_k^{\mathrm{cyc}}(C[1])$ is the quotient of $B_k(C[1])$ by the submodule generated by $\mathrm{cyc}(\mathbf{x}) - \mathbf{x}$ for $\mathbf{x} \in B_k(C[1])$. We put:

$$\widehat{B}^{\mathrm{cyc}}(C[1]) = \widehat{\bigoplus_{k=0}^{\infty}} B_k^{\mathrm{cyc}}(C[1])$$

the completed direct sum of them.

For a filtered A_∞ algebra $(C, \{\mathfrak{m}_k\}_{k=0}^{\infty})$ we define $\widehat{d} : \widehat{B}(C[1]) \to \widehat{B}(C[1])$ by

$$\widehat{d}(\mathbf{x}) = \sum_c (-1)^{\deg' \mathbf{x}^{3;1}} \mathbf{x}_c^{3;1} \otimes \mathfrak{m}_{|\mathbf{x}_c^{3;2}|}(\mathbf{x}_c^{3;2}) \otimes \mathbf{x}_c^{3;3}, \tag{17.7}$$

where $\mathbf{x}_c^{3;2} \in B_{|\mathbf{x}_c^{3;2}|}(C[1])$.

The equation $\widehat{d} \circ \widehat{d} = 0$ holds by definition of filtered A_∞ algebra.

We also define

$$\begin{aligned} \delta^{\mathrm{cyc}}(\mathbf{x}) = & \sum_c (-1)^{\deg' \mathbf{x}^{3;1}} \mathbf{x}_c^{3;1} \otimes \mathfrak{m}_{|\mathbf{x}_c^{3;2}|}(\mathbf{x}_c^{3;2}) \otimes \mathbf{x}_c^{3;3} \\ & + \sum_{c : \mathbf{x}^{3;1} \neq 1} (-1)^{(\deg' \mathbf{x}^{3;1} + \deg' \mathbf{x}^{3;2}) \deg' \mathbf{x}^{3;3}} \\ & \qquad \mathfrak{m}_{|\mathbf{x}_c^{3;1}| + |\mathbf{x}_c^{3;3}|}(\mathbf{x}_c^{3;3} \otimes \mathbf{x}_c^{3;1}) \otimes \mathbf{x}_c^{3;2} \\ & - \mathfrak{m}_0(1) \otimes \mathbf{x}. \end{aligned} \tag{17.8a}$$

In other words[3]

$$\begin{aligned} & \delta^{\mathrm{cyc}}(x_1 \otimes \cdots \otimes x_k) \\ & = \sum_{1 \le i < j \le k} (-1)^{*} x_1 \otimes \cdots \otimes \mathfrak{m}_{j-i+1}(x_i, \dots, x_j) \otimes \cdots \otimes x_k \\ & \quad + \sum_{1 \le i < j \le k} (-1)^{**} \mathfrak{m}_{i+k-j+1}(x_j, \dots, x_k, x_1, \dots, x_i) \otimes x_{i+1} \otimes \cdots \otimes x_{j-1} \\ & \quad - \mathfrak{m}_0(1) \otimes (x_1 \otimes \cdots \otimes x_k) \end{aligned} \tag{17.8b}$$

where

$$* = \deg' x_1 + \cdots + \deg' x_{i-1}, \quad ** = (\deg' x_j + \cdots + \deg' x_k)(\deg' x_1 + \cdots + \deg' x_i).$$

Lemma 17.3. $\delta^{\mathrm{cyc}} \circ \delta^{\mathrm{cyc}} = 0$.

The proof is a straightforward calculation which we omit.

We can prove easily that δ^{cyc} induces an operator on $\widehat{B}^{\mathrm{cyc}}(C[1])$, which we denote by the same symbol. Therefore $(\widehat{B}^{\mathrm{cyc}}(C[1]), \delta^{\mathrm{cyc}})$ is a chain complex.

Remark 17.4. We do not need any kind of cyclic symmetry of $\mathfrak{m}$ in order to prove the well-defined-ness of $\widehat{B}^{\mathrm{cyc}}(C[1])$ by δ^{cyc}.

---

[3] The authors thank to Dr.Ganatra who explained that one needs to subtract the term $\mathfrak{m}_0(1) \otimes \mathbf{x}$ in the differential δ^{cyc} during the conference on Cyclic symmetry in symplectic geometry at American Institute of Math. 2009 October.

Remark 17.5. In [FOOO3] Lemma & Definition 3.8.4, the differential on $\widehat{B}^{\rm cyc}(C[1])$ is defined by restricting $\widehat{d}$. In this article we defined $\widehat{B}^{\rm cyc}(C[1])$ as a quotient of $\widehat{B}(C[1])$. So we need to define differential in a different way. (See Remark 6.8.)

We remark that on the submodule of cyc invariant elements $\widehat{d}$ is proportional to $\delta^{\rm cyc}$. More precisely, the Diagram below commutes.

$$\begin{tikzcd}
(B_kC[1])^{\mathbb{Z}_k} \arrow[r, "\times 1/k"'] \arrow[d, "\widehat{\mathfrak m}_{k-\ell}"'] & B_kC[1] \arrow[r] & B_k^{\rm cyc}C[1] \arrow[d, "{\rm pro}\circ\widehat{\delta}^{\rm cyc}"] \\
(B_\ell C[1])^{\mathbb{Z}_\ell} \arrow[r, "\times 1/\ell"'] & B_\ell C[1] \arrow[r] & B_\ell^{\rm cyc}C[1]
\end{tikzcd}$$

where the 1st column is the invariant part of the action of symmetric group. The first horizontal map is the $1/k$ times the inclusion homomorphism (or $1/\ell$ times the inclusion homomorphism), respectively. The second horizontal map is the projection.

The left vertical map is a coderivation induced by $\mathfrak m_{k-\ell}$ and the right vertical map is the composition of $\delta^{\rm cyc}$ and the projection.

We can generalize the construction of $\widehat{d}$ as follows. Let H be a graded free $\Lambda_{0,\rm nov}$ module. We define $E_k(H[2])$, $\widehat{E}(H[2])$ as in Section 6.

Using a sequence of operators

$$\mathfrak q_{\ell;k} : E_\ell(H[2]) \otimes B_k(H^*(L;R)[1]) \to H^*(L;R)[1] \tag{17.9}$$

$k, \ell = 0, 1, 2, \dots$ introduced in Section 6, we obtain $\widehat{\mathfrak q} : E(H[2]) \otimes \widehat{B}(C[1]) \to \widehat{B}(C[1])$ by

$$\widehat{\mathfrak q}_\ell(\mathbf y;\mathbf x) = \sum_c (-1)^{\deg' \mathbf x^{3;1}(\deg' \mathbf y+1)} \mathbf x_c^{3;1} \otimes \mathfrak q_{\ell;|\mathbf x_c^{3;2}|}(\mathbf y;\mathbf x_c^{3;2}) \otimes \mathbf x_c^{3;3}. \tag{17.10}$$

We also define $\widehat{\widehat{\mathfrak q}}$ by

$$\begin{aligned}\widehat{\widehat{\mathfrak q}}_\ell(\mathbf y;\mathbf x) = &\sum_c (-1)^{\deg' \mathbf x^{3;1}(\deg' \mathbf y+1)} \mathbf x_c^{3;1} \otimes \mathfrak q_{\ell;|\mathbf x_c^{3;2}|}(\mathbf y;\mathbf x_c^{3;2}) \otimes \mathbf x_c^{3;3} \\ &+ \sum_{c:\mathbf x^{3;1}\ne 1} (-1)^{(\deg' \mathbf x^{3;1}+\deg' \mathbf x^{3;2})\deg' \mathbf x^{3;3}} \\ &\qquad \mathfrak q_{|\mathbf x_c^{3;1}|+|\mathbf x_c^{3;3}|}(\mathbf y;\mathbf x_c^{3;3}\otimes \mathbf x_c^{3;1}) \otimes \mathbf x_c^{3;2}.\end{aligned} \tag{17.11}$$

It induces $\widehat{\mathfrak q}^{\rm cyc} : E(H[2]) \otimes \widehat{B}^{\rm cyc}(C[1]) \to \widehat{B}^{\rm cyc}(C[1])$. (We need to subtract a term similar to the last term of (17.8a).)

Now we consider $C = H(L;\Lambda_{0,\rm nov})$, $H = H(X;\Lambda_{0,\rm nov})$. Here L is a relatively spin Lagrangian submanifold of a symplectic manifold X. (We do *not* assume that X is a toric manifold or L is a torus in this section.) In this case, the operator $\mathfrak q$ is defined in Sections 3.8 and 7.4 of [FOOO3]. It was explained in Section 2 [FOOO5]. In fact, we have

$$\mathfrak{PO}_{\mathfrak b}(b) = \sum_{\ell=0}^\infty\sum_{k=0}^\infty \frac{1}{\ell!}\mathfrak q_{\ell;k}(\mathfrak b^{\otimes \ell}; b^{\otimes k})$$

and

$$i^{\#}_{{\rm qm},(\mathfrak b,b)}(Q) = (-1)^{\deg Q}\sum_{\ell=0}^\infty\sum_{k=0}^\infty \frac{1}{\ell!}\mathfrak q_{\ell+1;k}([Q\otimes \mathfrak b^{\otimes \ell}]; b^{\otimes k}).$$

We define the operator $GW_{0,\ell+1} : E_{\ell+1}(H(X;\Lambda_0)) \to H(X;\Lambda_0)$ by

$$\langle \mathfrak{a}_0, GW_{\ell+1}(\mathfrak{a}_1 \otimes \cdots \otimes \mathfrak{a}_{\ell+1})\rangle_{PD_X} = GW_{\ell+2}(\mathfrak{a}_0 \otimes \mathfrak{a}_1 \otimes \cdots \otimes \mathfrak{a}_{\ell+1})$$

where the right hand side is explained in Section 2. Let $[x_1 \otimes \cdots \otimes x_k]$ be the equivalence class of $x_1 \otimes \cdots \otimes x_k \in B_k(C[1])$ in $B_k^{\rm cyc}(C[1])$. Let $(S(X;\Lambda_0), \delta)$ be an appropriate cochain complex whose cohomology group is the cohomology group of X. (In [FOOO3] we used a countably generated subcomplex of singular chain complex. In the next section we will use de Rham complex.)

Theorem 17.6. (Theorem 3.8.9 [FOOO3]) *There exists a family of the operators*

$$\mathfrak{p}_{\ell;k} : E_\ell(H(X;\Lambda_{0,\rm nov})[2]) \otimes B_k^{\rm cyc}(H(L;\Lambda_{0,\rm nov})[1]) \to S(X;\Lambda_{0,\rm nov})$$

for $k, \ell = 0, 1, \dots$ with the following properties.

(1) *Let $k > 0$ and $\mathbf{y} \in \widehat{E}(H(X;\Lambda_{0,\rm nov})[2])$ and $\mathbf{x} \in \widehat{B}_k^{\rm cyc}(H(L;\Lambda_{0,\rm nov})[1])$. Then, we have*

$$\delta(\mathfrak{p}(\mathbf{y};\mathbf{x})) + \sum_c \mathfrak{p}(\mathbf{y}_c^{(2;1)}; (\widehat{\mathfrak{q}}^{\rm cyc}(\mathbf{y}_c^{(2;2)};\mathbf{x}))) = 0.$$

(2) *Let $1 \in \widehat{B}_0^{\rm cyc}(H(L;\Lambda_{0,\rm nov})[1])$ and $\mathbf{y} \in \widehat{E}(H(X;\Lambda_{0,\rm nov})[2])$. Then we have*

$$\begin{aligned}\delta(\mathfrak{p}(\mathbf{y};1)) + \sum_c (-1)^{\deg \mathbf{y}_c^{(2;1)}} \mathfrak{p}(\mathbf{y}_c^{(2;1)}; (\mathfrak{q}(\mathbf{y}_c^{(2;2)};1))) \\ + \widetilde{GW}_{0,\ell+1}(X)(\mathbf{y}; PD[L]) = 0.\end{aligned}$$

Here $\widetilde{GW}_{0,\ell+1}$ is a homomorphism to $S(X;\Lambda_{0,\rm nov})$ which realize $GW_{0,\ell+1}$ in the cohomology level.

(3) *Let $\mathbf{e}_L = PD[L]$ be the unit, that is the Poincaré dual to the fundamental class of L. Then we have*

$$\mathfrak{p}_{\ell;k}(\mathbf{y}; [\mathbf{e}_L \otimes x_1 \otimes \cdots \otimes x_{k-1}]) = 0.$$

(4) *Let $\mathbf{e}_X = PD[X]$ be the Poincaré dual to the fundamental class of X. Then we have the following:*
   - *If $k > 2$, or $\ell \geq 1$, then*

$$\mathfrak{p}_{\ell+1;k}([\mathbf{e}_X \otimes y_1 \otimes \cdots \otimes y_\ell]; \mathbf{x}) = 0.$$

   - *For $\ell = 1$, $k = 0$,*

$$\mathfrak{p}_{1;0}(\mathbf{e}_X; 1) = i_!(\mathbf{e}_L).$$

   - *For $\ell = 1$, $k = 1$,*

$$\mathfrak{p}_{1;1}(\mathbf{e}_X; x) = i_!(x).$$

(5) *For $\ell \geq 1$ we have $\mathfrak{p}_{k;\ell} \equiv 0 \mod \Lambda_{0,nov}^+$. (Here $\Lambda_{0,nov}^+$ is the set of all elements of $\Lambda_{0,nov}$ such that all the exponents of T is positive.)*

Remark 17.7. (1) Theorems 3.8.9 and Proposition 3.8.78 of [FOOO3] are the chain level versions of Theorem 17.6. It is reduced to the cohomology version in Theorem 7.4.192 [FOOO3], from which Theorem 17.6 immediately follows.

(2) In Proposition 3.8.78 [FOOO3] it is claimed that, in case $\ell = 0$ and $k = 2$,

$$\mathfrak{p}_{2;0}(1; [\mathbf{e}_L \otimes x]) = \mathfrak{m}_2([\mathbf{e}_L \otimes x]). \tag{17.12}$$

This may look inconsistent with (3) above. In fact, since we take the quotient to define $B_k^{\rm cyc}(C[1])$ in this paper (in [FOOO3] the $\mathbb{Z}_k$ invariant part is taken), so (17.12) should be changed. Namely the right hand side becomes

$$\delta^{\rm cyc}([\mathbf{e}_L \otimes x]) = \mathfrak{m}_2(\mathbf{e}_L \otimes x) + (-1)^{\deg' x}\mathfrak{m}_2(x \otimes \mathbf{e}_L) = 0.$$

Here we use $\deg' \mathbf{e}_L = 1$ for the first equality, and the defining relation of a unit $\mathbf{e}_L$

$$\mathfrak{m}_2(\mathbf{e}_L \otimes x) = (-1)^{\deg x}\mathfrak{m}_2(x \otimes \mathbf{e}_L) = x$$

for the second equality (see Condition 3.2.18 [FOOO3]).

(3) In [FOOO3] Theorem 3.8.9, there is a factor $1/\ell!$ in front of the term $\widetilde{GW}_{0,\ell+1}(X)(\mathbf{y}; PD[L])$. We do not have one here since we take the quotient of BC by the action of symmetric group for the definition of EC, instead of taking the invariant subset of BC. See Remark 6.8.

(4) We mention some of the related works in Section 31.

Let $x \in C[1]$ be an element of an even (shifted) degree. We consider the element $e^x = \sum_{k=0}^{\infty} x^{\otimes k} \in B(C[1])$. For $w \in C$, we consider

$$e^x w e^x \in B(C[1]), \quad [we^x] \in B^{\rm cyc}C.$$

Let $\mathfrak{b} \in H^2(X; \Lambda_{0,\rm nov}^+)$ and $b \in H^1(L; \Lambda_{0,\rm nov}^+)$. We recall $e^{\mathfrak{b}} = \sum \mathfrak{b}^{\otimes \ell}/\ell!$. We assume that $(\mathfrak{b}, b)$ is a weak bounding cochain. Namely we assume $\mathfrak{q}(e^{\mathfrak{b}}; e^b) = \mathfrak{PO}_{\mathfrak{b}}(b)\, \mathbf{e}_L$, where $\mathfrak{PO}_{\mathfrak{b}}(b) \in \Lambda_{0,\rm nov}^+$. For $x \in H(L; \Lambda_{0,\rm nov})$ we define an operator $\delta_{\rm can}^{\mathfrak{b},b} : H(L; \Lambda_{0,\rm nov}) \to H(L; \Lambda_{0,\rm nov})$ of degree 1 by $\delta_{\rm can}^{\mathfrak{b},b}(w) = \mathfrak{q}(e^{\mathfrak{b}}; [we^b])$. The basic properties of $\mathfrak{q}$ (see Theorem 3.8.32 [FOOO3] or Theorem 2.1 [FOOO5]) imply $\delta_{\rm can}^{\mathfrak{b},b} \circ \delta_{\rm can}^{\mathfrak{b},b} = 0$.

Definition 17.8. We define a homomorphism

$$i_{*,{\rm qm},(\mathfrak{b},b)} : H(L; \Lambda_{0,\rm nov}) \to S(X; \Lambda_{0,\rm nov})$$

of degree 0 by

$$i_{*,{\rm qm},(\mathfrak{b},b)}(x) = \mathfrak{p}(e^{\mathfrak{b}}; [xe^b]). \tag{17.13}$$

Lemma 17.9. *$i_{*,{\rm qm},(\mathfrak{b},b)}$ is a chain map. Namely we have*

$$\delta \circ i_{*,{\rm qm},(\mathfrak{b},b)} + i_{*,{\rm qm},(\mathfrak{b},b)} \circ \delta_{\rm can}^{\mathfrak{b},b} = 0.$$

Note the boundary operator on $H(X; \Lambda_{0,\rm nov})$ is 0.

Proof. We remark that

$$\Delta(e^{\mathfrak{b}}) = e^{\mathfrak{b}} \otimes e^{\mathfrak{b}}.$$

It follows that

$$\begin{aligned}
&\widehat{\mathfrak{q}}^{\rm cyc}(e^{\mathfrak{b}}; [xe^b]) \\
&= [\mathfrak{q}(e^{\mathfrak{b}}; e^b x e^b) e^b] + (-1)^{\deg' x}[xe^b \mathfrak{q}^{\rm cyc}(e^{\mathfrak{b}}; e^b) e^b] \\
&= [\delta_{\rm can}^{\mathfrak{b},b}(x)\, e^b] + (-1)^{\deg' x}\mathfrak{PO}(\mathfrak{b}, b)\, [xe^b \mathbf{e}_L e^b].
\end{aligned}$$

Therefore Theorem 17.6 (1) and (3) imply

$$-(\delta \circ \mathfrak{p})(e^{\mathfrak{b}}; [xe^b]) = \mathfrak{p}(e^{\mathfrak{b}}; [\delta_{\rm can}^{\mathfrak{b},b}(x) e^b]) = (i_{*,{\rm qm},(\mathfrak{b},b)} \circ \delta_{\rm can}^{\mathfrak{b},b})(x),$$

as required. $\square$

We recall

$$HF((L(u),\mathfrak{b},b);(L(u),\mathfrak{b},b);\Lambda_0)=\frac{\mathrm{Ker}\delta^{\mathfrak{b},b}_{\mathrm{can}}}{\mathrm{Im}\delta^{\mathfrak{b},b}_{\mathrm{can}}}.$$

Therefore $i_{*,\mathrm{qm},(\mathfrak{b},b)}$ induces a map (17.1). We will discuss (17.5) in Section 19.

Remark 17.10. In this section, where we review general story from [FOOO3], we use the universal Novikov ring $\Lambda_{0,\mathrm{nov}}$ which contains two formal parameters T and e. In other sections of this current paper we use Λ_0 which contains only T. The parameter e is used so that all the operators have the well defined degree (for example $\mathfrak{m}_k$ has degree 1). Stories over $\Lambda_{0,\mathrm{nov}}$ coefficients and over Λ_0 are basically the same if we take enough care of the degree. Since the ring Λ_0 behaves better than $\Lambda_{0,\mathrm{nov}}$ from the point of view of commutative algebra, we use Λ_0 in our toric case.

18. CYCLIC SYMMETRY IN THE TORIC CASE

The discussion on the operator $\mathfrak{p}$ in Section 17 shows that it is essential to use a system of perturbations of the moduli spaces $\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta)$ that is invariant under the cyclic permutation of the boundary marked points. The cyclic symmetry is also important when we study the moduli spaces of holomorphic annuli in Section 20. (In doing so we need to consider the fiber product of moduli spaces of holomorphic discs, which appear at the boundary of the moduli space of holomorphic annuli, and to perturb it in a way invariant under the cyclic permutation of the boundary marked points. See the proof of Lemma 20.20.)

Such a perturbation is constructed in [Fu2]. In this section we adapt this construction to the present situation and explain how we can apply a similar scheme to construct the cyclically symmetric analog $\mathfrak{q}^{\mathfrak{c}}$ to the operator $\mathfrak{q}$. We also study the relationship between the operators $\mathfrak{q}$ and $\mathfrak{q}^{\mathfrak{c}}$.

The review of the perturbation to construct the operator $\mathfrak{q}$ is in Section 6. We next discuss the cyclically symmetric perturbation of $\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})$. Let

$$\mathrm{cyc}:\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})\to\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p}) \tag{18.1}$$

be the map induced by the cyclic permutation of marked points that sends $(0,1,\dots,k)$ to $(1,\dots,k,0)$. It generates the action of $\mathbb{Z}_{k+1}$. We can construct the Kuranishi structure of $\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})$ to which the $\mathbb{Z}_{k+1}$ action extends.

Proposition 18.1. *There exists a system of continuous family of multi-sections satisfying the properties (1)-(5) stated in Condition 6.3. In addition, it also satisfies:*

(6) *It is invariant under the above $\mathbb{Z}_{k+1}$-action.*

Remark 18.2. The notion of continuous family of multisections and its application to the smooth correspondence in de Rham theory is discussed in [FOOO5] Section 12. In our case where there is a T^n action, we can perform the construction there so that the family of multisections is T^n-invariant. We need to include the case where the parameter space of the family of multisections admits a T^n action. We will describe it in Sections 29-30 for completeness' sake.

Actually we can work out the whole argument without requiring T^n invariance of the family of multisections in Proposition 18.1. If we perform the construction in a T^n-invariant way, construction of the canonical model becomes much simpler. This is the reason why we use the T^n invariance in Proposition 18.1. On the contrary, in the construction of the operator $\mathfrak q$ where we use a single multisection, T^n invariance is used to assure surjectivity of the evaluation map. This surjectivity is used in [FOOO5] Proposition 6.6 and to construct the embedding $H^1(L(u);\Lambda_0) \subset \mathcal M_{\rm weak}(L(u);\Lambda_0)$. In this regard, the latter usage is more serious than that of the case of Proposition 18.1.

Proof of Proposition 18.1. Except the statement on T^n invariance, Proposition 18.1 follows from [Fu2] Corollary 3.1. We can perform the construction in a T^n-equivariant way. See Sections 29 - 30 for detail. □

Remark 18.3. Here is an important remark to be made. In [Fu2] one of the key idea to construct a cyclically symmetric (family of) perturbations is to include $\mathcal M_{0;\ell}(\beta;\mathbf p)$ also. In [FOOO4] the construction of the $\mathfrak q$-perturbation on $\mathcal M_{k+1;\ell}(\beta;\mathbf p)$ uses the fact that the T^n action there is free. This does not hold for the case $k+1=0$. Therefore the argument of [FOOO4] is not enough to construct cyclically symmetric perturbation. It seems that it is necessary to use a *continuous family* of multisections for the proof of Proposition 18.1.

Definition 18.4. We call the system of continuous family of perturbations in Proposition 18.1 the $\mathfrak c$ *perturbation* and write the corresponding perturbed moduli space as $\mathcal M_{0;\ell}(\beta;\mathbf p)^{\mathfrak c}$.

As we show later in this section, we can use a $\mathfrak c$-perturbation in place of the $\mathfrak q$-perturbation to define an operator that is not only cyclically symmetric but shares most of the properties of $\mathfrak q$.

However there is one property of the $\mathfrak q$-perturbation which is not shared by the $\mathfrak c$-perturbation. The next lemma is related to this point. Let $\mathbf p = (\mathbf p(1),\dots,\mathbf p(\ell))$ be T^n-invariant cycles as in Section 6.

Condition 18.5. X is nef and $\deg \mathbf p(i) = 2$ for all i. (Note $\dim \mathbf p(i) = 2n - \deg \mathbf p(i)$.)

Lemma 18.6. *Assume Condition 18.5. We may take a $\mathfrak c$-perturbation so that the following holds. $\mathcal M_{k+1;\ell}(\beta;\mathbf p)^{\mathfrak c}$ is empty if one of the following is satisfied for $k \ge 0$:*

(1) $\mu(\beta) - \sum_i (2n - \dim \mathbf p(i) - 2) < 0$.
(2) $\mu(\beta) - \sum_i (2n - \dim \mathbf p(i) - 2) = 0$ *and* $\beta \ne 0$.

Proof. Suppose the case (2) holds. Since $\deg \mathbf p(i) \ge 2$, we have

$$\dim \mathcal M_{1;\ell}(\beta;\mathbf p) \le \dim \mathcal M_{1;0}(\beta) = \mu(\beta) + n - 2 \le n-2. \tag{18.2}$$

Here dim is the (virtual) dimension as a space with Kuranishi structure.

We assume that $\mathcal M_{1;0}(\beta)$ is nonempty. By [FOOO4] Theorem 11.1 we have

$$\beta = \beta_{i_1} + \dots + \beta_{i_l} + \alpha$$

where $i_1,\dots,i_l \in \{1,\dots,m\}$, $\alpha \in H_2(X;\mathbb Z)$ and α is realized by a sum of holomorphic spheres. This is impossible since $c_1(\alpha) \ge 0$ by Condition 18.5 and $\mu(\beta) < 0$ (by (18.2) and $\beta \ne 0$) for the case (2). Thus $\mathcal M_{1;\ell}(\beta;\mathbf p) = \emptyset$.

We can prove it also in case (1) similarly.

Thus we do not need to perturb it. Hence it is empty set after perturbation also. □

Remark 18.7. Lemma 18.6 is similar to [FOOO5] Corollary 6.6. However in [FOOO5] Corollary 6.6 we did not need Condition 18.5. In case we can use (a single) multisection, the moduli space with negative (virtual) dimension can be perturbed and becomes an empty set. This is not necessary the case when we use a continuous family of multisections.

Definition 18.8. We use the perturbed moduli space $\mathcal{M}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{c}}$ to define

$$\mathfrak{q}^{\mathfrak{c}}_{\ell;k+1;\beta} : E_\ell(\mathcal{A}[2]) \otimes B_k(H(L(u);\mathbb{R})[1]) \to H(L(u);\mathbb{R})[1]$$

by

$$\mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k) = \mathrm{ev}_{0*}(\mathrm{ev}^*(h_1 \times \cdots \times h_k); \mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{c}}) \tag{18.3}$$

where

$$(\mathrm{ev}_0, \mathrm{ev}) : \mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{c}} \to L(u) \times L(u)^k$$

and h_i are T^n-equivariant forms. (Here we use Convention 6.12.) The right hand side of (18.3) is independent of the choice of the representative $\mathbf{p}$ but depends only on the equivalence class $[\mathbf{p}]$. This is a consequence of Condition 6.3 (5).

Next we define

$$\begin{aligned}&\mathfrak{q}^{\rho,\mathfrak{c}}_{\ell;k}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k)\\ &= \sum_{\ell=0}^{\infty} \sum_{\beta \in H_2(X;L(u);\mathbb{Z})} T^{\beta\cap\omega/2\pi} \rho(\partial\beta) \mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}([\mathbf{p}]; h_1 \otimes \cdots \otimes h_k).\end{aligned}$$

Theorem 18.9. *The operators* $\mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}$ *have the following properties:*

1. (6.11a) *holds. (We replace* $\mathfrak{q}$ *by* $\mathfrak{q}^{\mathfrak{c}}$*.)*
2. (6.11b) *and* (6.11c) *hold. (We replace* $\mathfrak{q}$ *by* $\mathfrak{q}^{\mathfrak{c}}$*.)*
3. *The operator* $\mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}$ *is cyclic, i.e., satisfies*

$$\begin{aligned}&\langle \mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}(\mathbf{y}; h_1, \ldots, h_k), h_0 \rangle_{\mathrm{cyc}}\\ &= (-1)^{\deg' h_0(\deg' h_1 + \cdots + \deg' h_k)} \langle \mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}(\mathbf{y}; h_0, h_1, \ldots, h_{k-1}), h_k \rangle_{\mathrm{cyc}}.\end{aligned} \tag{18.4}$$

Here $\langle \cdot, \cdot \rangle_{\mathrm{cyc}}$ *is defined by* (2.22)*, see also Subsection 26.1.*

Proof. (1) and (2) follows from (1) – (5) Condition 6.3 in the same way as the case of $\mathfrak{q}$. (3) is a consequence of (6) Proposition 18.1. □

Remark 18.10. Actually we need to fix an energy level E_0 and restrict the construction of the family of multisections for the moduli space of maps of homology class β with $\beta \cap \omega \le E_0$, by the reason explained in [FOOO3] Subsection 7.2.3. (After then we can use homological algebra to extend construction of the operators to all β's.)

This process is discussed in [Fu2] Section 14, following [FOOO3] Section 7.2. (See also [Fu2] Section 14.) In the current T^n-equivariant case, we can perform the same construction in a T^n-equivariant way. We will not repeat this kind of remarks in other similar situations in the rest of this paper.

The sign of (18.4) looks different from the one in [FOOO3] at glance but is mathematically consistent. We here use the sign convention of [Fu2]. See [Fu2] Lemma 6.1 about the relationship between two conventions.

Definition 18.11. Let $\mathfrak{b} \in H^{even}(X;\Lambda_0)$ and $\rho : H_1(L(u);\mathbb{Z}) \to \mathbb{C}^*$ a local system. We write

$$\mathfrak{b} = \mathfrak{b}_0 + \mathfrak{b}_2 + \mathfrak{b}_{\text{high}}$$

where $\mathfrak{b}_0 \in H^0(X;\Lambda_0)$, $\mathfrak{b}_2 \in H^2(X;\Lambda_0)$, $\mathfrak{b}_{\text{high}} \in H^{2m}(X;\Lambda_0)$ $(m > 1)$. We define $\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k;\beta} : B_k(H(L;\mathbb{R})[1]) \to H(L;\mathbb{R})[1]$ by

$$\begin{aligned}&\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k}(h_1,\dots,h_k)\\ &= \sum_{\beta}\sum_{\ell=0}^{\infty} T^{\beta\cap\omega/2\pi}\rho(\partial\beta)\exp(\mathfrak{b}_2\cap\beta)\mathfrak{q}^{\mathfrak{c}}_{\ell;k;\beta}(\mathfrak{b}_{\text{high}}^{\ell};h_1,\dots,h_k).\end{aligned} \tag{18.5}$$

Let $b \in H^{odd}(L(u);\Lambda_0)$. We define

$$b = b_0 + b_+$$

where $b_0 \in H^1(L(u);\mathbb{C})$ and $b_+ \in H^1(L(u);\Lambda_+) \oplus \bigoplus_{m>0} H^{2m+1}(L(u);\Lambda_0)$. We define $\rho : H_1(L(u);\mathbb{Z}) \to \mathbb{C}^*$ by $\gamma \mapsto e^{\gamma\cap b_0}$. We define

$$\mathfrak{m}^{\mathfrak{c},\mathfrak{b},b}_{k}(h_1,\dots,h_k) = \sum_{l_0,\dots,l_k} \mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k+\sum l_i;\beta}(b_+^{l_0},h_1,b_+^{1},\dots,b_+^{l_{k-1}},h_k,b_+^{l_k}). \tag{18.6}$$

In the rest of this paper we only consider T^n-invariant differential forms and hence we always work with the canonical model.

Remark 18.12. Convergence of the series in the right hand side of (18.6) is proved in the same way as in the proof of Lemma 7.11.

We handle degree 2 classes $\mathfrak{b}_2$ in a different way from higher degree classes $\mathfrak{b}_{\text{high}}$. We do so for the proof of convergence of (18.6) and similar other series.

Proposition 18.13. *$(H(L(u);\Lambda_0),\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_k\}_{k=0}^{\infty})$ and $(H(L(u);\Lambda_0),\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},b}_k\}_{k=0}^{\infty})$ are unital and cyclically symmetric filtered A_∞ algebras.*

Proof. This follows from Proposition 18.9. □

Lemma 18.14. *If X is nef and $\mathfrak{b} \in H^2(X;\Lambda_0)$, then*

$$H^1(L(u);\Lambda_+) \subset \widehat{\mathcal{M}}_{\text{weak}}((H(L(u);\Lambda_0),\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_k\}_{k=0}^{\infty});\Lambda_+).$$

Proof. This follows from Lemma 18.6 in the same way as Proposition 3.1 [FOOO5] follows from Corollary 6.6 [FOOO5]. □

We remark that without the assumption that X is nef and $\mathfrak{b} \in H^2(X;\Lambda_0)$ it is not yet clear at this stage whether $(H(L(u);\Lambda_0),\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_k\}_{k=0}^{\infty})$ is weakly unobstructed. We will prove this below by comparing $\mathfrak{q}^{\mathfrak{c}}$ with $\mathfrak{q}$. We use the bifurcation method for this purpose. (See [FOOO3] Subsection 7.2.14 about the cobordism method and the bifurcation method.)

Let $H = H(L(u);\Lambda_0)$ and $\overline{H} = H(L(u);\mathbb{R})$. We consider the set of formal sums

$$a(t) + dt \wedge b(t) \tag{18.7}$$

where $a(t) \in C^\infty([0,1],\overline{H}^k)$, $b(t) \in C^\infty([0,1],\overline{H}^{k-1})$. We denote the set of such expressions (18.7) by $C^\infty([0,1]\times\overline{H})^k$. We will consider filtered A_∞ structure on it. More precisely, we proceed as follows.

We assume that, for each $t \in [0,1]$, we have operations:

$$\mathfrak{q}^t_{\ell;k,\beta} : E_\ell(\mathcal{A}[2]) \otimes B_k(\overline{H}[1]) \to \overline{H}[1] \tag{18.8}$$

of degree $-\mu(\beta)+1$ and

$$\mathfrak{Q}^t_{\ell;k,\beta} : E_\ell(\mathcal{A}[2]) \otimes B_k(\overline{H}[1]) \to \overline{H}[1] \tag{18.9}$$

of degree $-\mu(\beta)$.

Definition 18.15. We say $\mathfrak{q}^t_{\ell;k,\beta}$ is *smooth* if for each $y_1,\dots,y_\ell,x_1,\dots,x_k$

$$t \mapsto \mathfrak{q}^t_{\ell;k,\beta}(y_1,\dots,y_\ell;x_1,\dots,x_k)$$

is an element of $C^\infty([0,1],\overline{H})$.

The smoothness of $\mathfrak{Q}^t_{\ell;k,\beta}$ is defined in the same way.

Definition 18.16. We say $(H,\{\mathfrak{q}^t_{\ell;k,\beta}\},\{\mathfrak{Q}^t_{\ell;k,\beta}\})$ is a *pseudo-isotopy* of unital G-gapped filtered A_∞ algebras with bulk if the following holds:

1. $\mathfrak{q}^t_{\ell;k,\beta}$ and $\mathfrak{Q}^t_{\ell;k,\beta}$ are smooth.
2. For each (but fixed) t, the triple $(H,\{\mathfrak{q}^t_{k,\beta}\})$ satisfies the conclusion of Proposition 18.9.
3. Let $\mathbf{x} \in B_k(\overline{H}[1])$, $\mathbf{y} \in E_\ell(\mathcal{A}[2])$. We put

$$\Delta^2\mathbf{x} = \sum_{c_1\in C_1} \mathbf{x}_{c_1;1}\otimes\mathbf{x}_{c_1;2}\otimes\mathbf{x}_{c_1;3}, \quad \Delta\mathbf{y} = \sum_{c_2\in C_2} \mathbf{y}_{c_2;1}\otimes\mathbf{y}_{c_2;2}.$$

   Then we have

$$\begin{aligned}
&\frac{d}{dt}\mathfrak{q}^t_{\ell;k,\beta}(\mathbf{y};\mathbf{x})\\
&+\sum_{c_1\in C_1}\sum_{c_2\in C_2}\sum_{\beta_1+\beta_2=\beta}(-1)^*\mathfrak{Q}^t_{\beta_1}(\mathbf{y}_{c_2;1};\mathbf{x}_{c_1;1},\mathfrak{q}^t_{\beta_2}(\mathbf{y}_{c_2;2};\mathbf{x}_{c_1;2}),\mathbf{x}_{c_1;3})\\
&-\sum_{c_1\in C_1}\sum_{c_2\in C_2}\sum_{\beta_1+\beta_2=\beta}(-1)^{**}\mathfrak{q}^t_{\beta_1}(\mathbf{y}_{c_2;1};\mathbf{x}_{c_1;1},\mathfrak{Q}^t_{\beta_2}(\mathbf{y}_{c_2;2};\mathbf{x}_{c_1;2}),\mathbf{x}_{c_1;3})\\
&=0.
\end{aligned} \tag{18.10}$$

   Here the sign is given by $* = \deg'\mathbf{y}_{c_2;1} + \deg'\mathbf{x}_{c_1;1} + \deg'\mathbf{x}_{c_1;1}\deg'\mathbf{y}_{c_2;2}$ and $** = \deg'\mathbf{x}_{c_1;1}\deg'\mathbf{y}_{c_2;2}$.
4. $\mathfrak{q}^t_{\ell;k,\beta_0}$ is independent of t. $\mathfrak{Q}^t_{k,\beta_0}=0$. (Here $\beta_0 = 0$ in $H_2(X;L(u))$.)
5. $\mathfrak{q}^0_{\ell;k,\beta}$ satisfies (6.11a).

Proposition 18.17. *There exists a pseudo-isotopy* $(H,\{\mathfrak{q}^t_{k,\beta}\},\{\mathfrak{Q}^t_{k,\beta}\})$ *of unital G-gapped filtered A_∞ algebras with bulk such that:*

1. $\mathfrak{q}^0_{\ell;k,\beta} = \mathfrak{q}^\rho_{\ell;k,\beta}$.
2. $\mathfrak{q}^1_{\ell;k,\beta} = \mathfrak{q}^{\rho,\mathfrak{c}}_{\ell;k,\beta}$.

Proof. The proof is based on the following lemma.

Lemma 18.18. *There exists a system of continuous family of multisections* $\mathfrak{par}$ *on* $\mathcal{M}^{\text{main}}_{\ell;k+1;\beta}(\beta;\mathbf{p})\times[0,1]$ *with the following properties. We use s to parametrize* $[0,1]$.

1. *The* $[0,1]$ *parametrized version of Condition 6.3 holds.*
2. *At $s=0$ our family of multisections* $\mathfrak{par}$ *coincides with* $\mathfrak{q}$ *perturbation.*
3. *At $s=1$ our family of multisections* $\mathfrak{par}$ *coincides with* $\mathfrak{c}$ *perturbation.*

Proof. This is a straightforward analogue of the construction of $\mathfrak{q}$ perturbation. □

We use the perturbed moduli space $(\mathcal{M}_{k+1;\ell}(\beta;\mathbf{p})\times[0,1])^{\mathfrak{par}}$ to define

$$\mathfrak{q}\mathfrak{Q}_{\ell;k;\beta} : E_\ell(\mathcal{A}[2])\otimes B_k(H(L(u);\mathbb{R})[1]) \to C^\infty([0,1],\Lambda^*([0,1])\otimes H(L(u);\mathbb{R}))$$

(where $\Lambda^*([0,1])$ is the vector bundle of differential forms on $[0,1]$) by

$$\begin{aligned}&\mathfrak{q}\mathfrak{Q}_{\ell;k;\beta}([\mathbf{p}];h_1\otimes\cdots\otimes h_k)\\&=(\mathrm{ev}_0,\mathrm{ev}_s)_*(\mathrm{ev}^*(h_1\times\cdots\times h_k);\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{par}}).\end{aligned}\tag{18.11}$$

Here

$$(\mathrm{ev}_0,\mathrm{ev}) : (\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})\times[0,1])^{\mathfrak{par}} \to L(u)\times L(u)^k$$

are the evaluation maps at boundary marked points and

$$\mathrm{ev}_s : (\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})\times[0,1])^{\mathfrak{par}} \to [0,1]$$

is the projection.

We next define

$$\begin{aligned}&\mathfrak{q}\mathfrak{Q}^{\rho}_{\ell;k}([\mathbf{p}];h_1\otimes\cdots\otimes h_k)\\&=\sum_{\ell=0}^{\infty}\sum_{\beta\in H_2(X;L(u);\mathbb{Z})}T^{\beta\cap\omega/2\pi}\rho(\partial\beta)\mathfrak{q}\mathfrak{Q}_{\ell;k;\beta}([\mathbf{p}];h_1\otimes\cdots\otimes h_k).\end{aligned}$$

We restrict it to T^n-equivariant forms and obtain $\mathfrak{q}\mathfrak{Q}^{\rho,\mathrm{can}}_{\ell;k}$. By decomposing it to the part which contains ds and to the part which does not contain ds, we obtain $\mathfrak{Q}^t$ and $\mathfrak{q}^t$ which has the required properties. ☐

Lemma 18.19. *Suppose Condition 18.5 holds. Then*

$$\mathfrak{q}^t_{\ell;k;\beta}([\mathbf{p}];h_1,\ldots,h_k)=0,\quad \mathfrak{Q}^t_{\ell;k;\beta}([\mathbf{p}];h_1,\ldots,h_k)=0$$

if

(1) $\mu(\beta)-\sum_i(2n-\dim\mathbf{p}(i)-2)<0$,
(2) $\mu(\beta)-\sum_i(2n-\dim\mathbf{p}(i)-2)=0$ *and* $\beta\neq 0$.

The proof is similar to Lemma 18.6. (The dimension counting argument there works up to *one* parameter family since all $\mu(\beta)$ are even integers.)

Definition 18.20. Let $\mathfrak{b}\in H^{even}(X;\Lambda_0)$ and $\rho : H_1(L(u);\mathbb{Z})\to\mathbb{C}^*$. We write $\mathfrak{b}=\mathfrak{b}_0+\mathfrak{b}_2+\mathfrak{b}_{\mathrm{high}}$ where $\mathfrak{b}_0\in H^0(X;\Lambda_0)$, $\mathfrak{b}_2\in H^2(X;\Lambda_0)$, $\mathfrak{b}_{\mathrm{high}}\in H^{2m}(X;\Lambda_0)$ $(m>1)$. We define

$$\begin{aligned}&\mathfrak{m}^{t,\mathfrak{b},\rho}_k(h_1,\ldots,h_k)\\&=\sum_\beta\sum_{\ell=0}^{\infty}\frac{T^{\beta\cap\omega/2\pi}}{\ell!}\rho(\partial\beta)\exp(\mathfrak{b}_2\cap\beta)\mathfrak{q}^t_{\ell;k;\beta}(\mathfrak{b}^{\ell}_{\mathrm{high}};h_1,\ldots,h_k),\\&\mathfrak{c}^{t,\mathfrak{b},\rho}_{k,\mathfrak{b}}(h_1,\ldots,h_k)\\&=\sum_\beta\sum_{\ell=0}^{\infty}\frac{T^{\beta\cap\omega/2\pi}}{\ell!}\rho(\partial\beta)\exp(\mathfrak{b}_2\cap\beta)\mathfrak{Q}^t_{\ell;k;\beta}(\mathfrak{b}^{\ell}_{\mathrm{high}};h_1,\ldots,h_k).\end{aligned}\tag{18.12}$$

Let $b\in H^{odd}(L(u);\Lambda_0)$. We define $b=b_0+b_+$ where $b_0\in H^1(L(u);\mathbb{C})$ and $b_+\in H^1(L(u);\Lambda_+)\oplus\bigoplus_{m>0}H^{2m+1}(L(u);\Lambda_0)$. We define a local system $\rho : H_1(L(u);\mathbb{Z})\to\mathbb{C}^*$ by

$$\rho\ :\ \gamma\mapsto e^{\gamma\cap b_0}$$

and define

$$\begin{aligned}\mathfrak{m}_k^{t,b}(h_1,\dots,h_k) &= \sum_{l_0,\dots,l_k} \mathfrak{m}^{t,\mathfrak{b},\rho}_{k+\sum l_i;\beta}(b_+^{l_0},h_1,b_+^1,\dots,b_+^{l_{k-1}},h_k,b_+^{l_k}),\\ \mathfrak{c}_k^{t,b}(h_1,\dots,h_k) &= \sum_{l_0,\dots,l_k} \mathfrak{Q}^{t,\mathfrak{b},\rho}_{k+\sum l_i;\beta}(b_+^{l_0},h_1,b_+^1,\dots,b_+^{l_{k-1}},h_k,b_+^{l_k}).\end{aligned} \tag{18.13}$$

We consider $x_i(t) + dt \wedge y_i(t) = \mathbf{x}_i \in C^\infty([0,1],\overline{C})$. We define

$$\widehat{\mathfrak{m}}^{t,\mathfrak{b},\rho}_{k,\beta}(\mathbf{x}_1,\dots,\mathbf{x}_k) = x(t) + dt \wedge y(t),$$

where

$$x(t) = \mathfrak{m}^{t,\mathfrak{b},\rho}_{k,\beta}(x_1(t),\dots,x_k(t)) \tag{18.14a}$$

$$\begin{aligned}y(t) =&\mathfrak{c}^{t,\mathfrak{b},\rho}_{k,\beta}(x_1(t),\dots,x_k(t))\\ &-\sum_{i=1}^{k}(-1)^{*_i}\mathfrak{m}^{t,\rho}_{k,\beta}(x_1(t),\dots,x_{i-1}(t),y_i(t),x_{i+1}(t),\dots,x_k(t))\end{aligned} \tag{18.14b}$$

if $(k,\beta) \ne (1,\beta_0)$ $(\beta_0 = 0 \in H_2(X;L(u)))$ and

$$y(t) = \frac{d}{dt}x_1(t) + \mathfrak{m}^{t,\mathfrak{b},\rho}_{1,\beta_0}(y_1(t)) \tag{18.14c}$$

if $(k,\beta) = (1,\beta_0)$. Here $*_i$ in (18.14b) is $*_i = \deg' x_1 + \cdots + \deg' x_{i-1}$. We define $\widehat{\mathfrak{m}}^{t,\mathfrak{b},b}_{k,\beta}$ in a similar way.

We put

$$\begin{aligned}&(\mathrm{Eval}_{t=0})_1(x(t)+dt\wedge y(t)) = x(0), \quad (\mathrm{Eval}_{t=1})_1(x(t)+dt\wedge y(t)) = x(1),\\ &(\mathrm{Eval}_{t=0})_k = (\mathrm{Eval}_{t=1})_k = 0, \quad \text{for } k \ne 1.\end{aligned}$$

Lemma 18.21. *$(C^\infty([0,1]\times H),\{\widehat{\mathfrak{m}}^{t,\mathfrak{b},\rho}_{k,\beta}\})$ is a G gapped unital filtered A_∞ algebra.*

$\mathrm{Eval}_{t=0}$ is a strict unital filtered A_∞ homomorphism from $(C^\infty([0,1]\times H),\{\widehat{\mathfrak{m}}^{t,\mathfrak{b},\rho}_{k,\beta}\})$ to $(H,\{\mathfrak{m}^{\mathfrak{b},\rho}_{k,\beta}\})$.

$\mathrm{Eval}_{t=1}$ is a strict unital filtered A_∞ homomorphism from $(C^\infty([0,1]\times H),\{\widehat{\mathfrak{m}}^{t,\mathfrak{b},\rho}_{k,\beta}\})$ to $(H,\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k,\beta}\})$.

The proof is a straightforward calculation from Proposition 18.17.

Corollary 18.22. *$(H,\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k,\beta}\})$ is unital isomorphic to $(H,\{\mathfrak{m}^{\mathfrak{b},\rho}_{k,\beta}\})$.*

Since Eval induces an isomorphism on $\mathfrak{m}_{1,0}$-cohomology, this follows from Lemma 18.21 and [FOOO3] Theorem 4.2.45. (Note in the our case $\mathfrak{m}_{1,0} = 0$ so homotopy equivalence has an inverse. See [FOOO3] Proposition 5.4.5.) See also [Fu2] Proposition 4.1. The isomorphism constructed there from the pseudo-isotopy in Lemma 18.21 turns out to be the isomorphism obtained by the cobordism method as in [FOOO3] Subsection 4.6.1 using the time-ordered-product moduli space.

Hereafter we fix an isomorphism

$$\mathfrak{F} : (H,\{\mathfrak{m}^{\mathfrak{b},\rho}_{k,\beta}\}) \to (H,\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k,\beta}\}). \tag{18.15}$$

We note that

$$\mathfrak{F} \equiv \text{identity} \mod \Lambda_+. \tag{18.16}$$

Lemma 18.14 implies

$$\mathfrak{F}_*(H^1(L(u);\Lambda_+)) \subset \mathcal{M}_{\mathrm{weak}}((H,\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k,\beta}\});\Lambda_+) \tag{18.17}$$

and twisting the boundary map by non-unitary local systems ρ enables us to extend the domain of $\mathfrak{F}$ to $H^1(L(u);\Lambda_0)$.

Remark 18.23. Since the isomorphism $\mathfrak{F}$ may increase the degree, the image $\mathfrak{F}_*(H^1(L(u);\Lambda_+))$ may not necessarily be contained in $H^1(L(u);\Lambda_+)$ again. The authors do not know whether we can take $\mathfrak{F}_*$ so that its image be contained in $H^1(L(u);\Lambda_+)$. As a matter of fact, it is in general very hard to calculate the map $\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_{k,\beta}$.

However, there are two particular cases for which $\mathfrak{F}_*(H^1(L(u);\Lambda_+))$ is contained in the degree one cohomology $H^1(L(u);\Lambda_0)$. We discuss these two cases in detail now.

Lemma 18.24. *If dimension of* $L(u)$ *is* 2*, then*

$$H^1(L(u);\Lambda_+) \subset \mathcal{M}_{\rm weak}((H,\{\mathfrak{m}^{\mathfrak{c},\mathfrak{b},\rho}_k\});\Lambda_+).$$

Proof. We remark that $\mathfrak{F}_*$ preserves the parity of the degree of the elements. For the case of dimension 2, the elements of odd degree are necessarily of degree 1. The lemma then follows from (18.17). ☐

Lemma 18.25. *If* X *is nef and* $\mathfrak{b} \in H^2(X;\Lambda_0)$*, then*

$$C^\infty([0,1]\times H^1(L(u);\Lambda_+)) \subset \mathcal{M}_{\rm weak}((C^\infty([0,1]\times H),\{\widehat{\mathfrak{m}}^{t,\mathfrak{b},\rho}_{k,\beta}\});\Lambda_+).$$

Proof. This follows from Lemma 18.19 in the same way as Proposition 3.1 [FOOO5] follows from Corollary 6.6 [FOOO5]. ☐

Corollary 18.26. *If* X *is nef and* $\mathfrak{b} \in H^2(X;\Lambda_0)$*, then* $\mathfrak{F}_*$ *is an identity map on* $H^1(L(u);\Lambda_+)$*. In particular,*

$$H^1(L(u);\Lambda_+) \subset \mathcal{M}_{\rm weak}((H,\{\mathfrak{m}^{\mathfrak{b},\rho}_{k,\beta}\});\Lambda_+).$$

This is immediate from Lemma 18.25.

Definition 18.27. Let $b = b_0 + b_+ \in H^1(L(u);\Lambda_0)$, where $b_0 \in H^1(L(u);\mathbb{C})$ and $b_+ \in H^1(L(u);\Lambda_+)$. The term b_0 induces the representation $\rho = \rho_{b_0}$ as before. We put $b_+^{\mathfrak{c}} = \mathfrak{F}_*(b_+)$ and $b^{\mathfrak{c}} = b_0 + b_+^{\mathfrak{c}}$.

We next consider the homomorphism

$$i^{\#}_{{\rm qm},(\mathfrak{b},b,u)} : H(X;\Lambda_0) \to H(L(u);\Lambda_0)$$

in (17.1). It is defined as:

$$\begin{aligned} &i^{\#}_{{\rm qm},(\mathfrak{b},b,u)}(Q) \\ &= \sum_\beta\sum_\ell\sum_k \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\rho(\partial\beta)e^{\mathfrak{b}_2\cap\beta}\mathfrak{q}_{\ell+1;k;\beta}(Q\otimes\mathfrak{b}^{\ell}_{\rm high};b_+,\dots,b_+). \end{aligned} \tag{18.18}$$

We replace $\mathfrak{q}$ by $\mathfrak{q}^{\mathfrak{c}}$ above and b_+ by $b_+^{\mathfrak{c}}$ and obtain:

$$i^{\mathfrak{c}\#}_{{\rm qm},(\mathfrak{b},b^{\mathfrak{c}},u)} : H(X;\Lambda_0) \to H(L(u);\Lambda_0).$$

Remark 18.28. In case $\deg b_+ = 1$, we can rewrite (18.18) to

$$\sum_\beta\sum_\ell \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\rho^b(\partial\beta)e^{\mathfrak{b}_2\cap\beta}\mathfrak{q}_{\ell+1;0;\beta}(Q\otimes\mathfrak{b}^{\ell}_{\rm high};1). \tag{18.19}$$

See Section 27. In the current circumstance, $b_+^{\mathfrak{c}}$ may not have degree one. Because of this we can not replace (18.18) by (18.19) in our definition of $i^{\mathfrak{c}\#}_{\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}},u)}$.

Lemma 18.29. *If $\mathfrak{F}_*(b_+) = b_+^{\mathfrak{c}}$, then*

$$\mathfrak{F}_* \circ i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)} = i^{\mathfrak{c}\#}_{\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}},u)}. \tag{18.20}$$

Proof. We take $\tilde{b}_+$ which goes to b_+ and $b_+^{\mathfrak{c}}$ by $\mathrm{Eval}_{t=0}$ and $\mathrm{Eval}_{t=1}$ respectively. We use $\mathfrak{q}\mathfrak{Q}_{\ell;k;\beta} : E_\ell(\mathcal{A}[2]) \otimes B_k(H(L(u);\mathbb{R})[1]) \to C^\infty([0,1], H(L(u);\Lambda_0))[1]$ to define $i^{\#\mathfrak{para}}_{\mathrm{qm},(\mathfrak{b},b,u)} : H(X;\Lambda_0) \to C^\infty([0,1] \times H(L(u);\Lambda_0))$ in the same way. Then we have a commutative diagram:

$$\begin{tikzcd}
H(X;\Lambda_0) \arrow[r, "i^{\mathfrak{c}\#}_{\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}},u)}"] \arrow[d, equal] & H(L(u);\Lambda_0) \\
H(X;\Lambda_0) \arrow[r, "i^{\#\mathfrak{para}}_{\mathrm{qm},(\mathfrak{b},\tilde{b},u)}"] \arrow[d, equal] & C^\infty([0,1] \times H(L(u);\Lambda_0)) \arrow[u, "\mathrm{Eval}_{t=1}"'] \arrow[d, "\mathrm{Eval}_{t=0}"] \\
H(X;\Lambda_0) \arrow[r, "i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}"] & H(L(u);\Lambda_0)
\end{tikzcd} \tag{18.21}$$

By taking the cohomology of $C^\infty([0,1] \times H(L(u);\Lambda_0))$, the right hand side becomes an isomorphism which is nothing but $\mathfrak{F}_*$. The lemma follows. □

In the rest of this section, we describe the morphism $\mathfrak{F}$ for the case in which $n = 2$ but X is not nef, and explain how $\mathfrak{q}^{\mathfrak{c}}$ could be different from $\mathfrak{q}$. Namely we compare $\mathfrak{PO}$ and $\mathfrak{PO}^{\mathfrak{c}} = \mathfrak{PO} \circ \mathfrak{F}_*^{-1}$.

Note

$$\mathfrak{PO}^{\mathfrak{c}}(0;b) = \sum_\beta \sum_{k \ge 0} \rho(\partial\beta) e^{\mathfrak{b}_2 \cap \beta} \mathfrak{q}^{\mathfrak{c}}_{0;k+1;\beta}(1; b_+, \dots, b_+). \tag{18.22}$$

Here $b = b_0 + b_+ \in H^1(L(u);\Lambda_0)$, $b_0 \in H^1(L(u);\mathbb{C}), b_+ \in H^1(L(u);\Lambda_+)$, and ρ is induced by b_0 as before.

We consider the moduli space $[0,1] \times \mathcal{M}^{\mathrm{main}}_{0;k+1}(\beta)$ and its family of multisections $\mathfrak{par}$ as in Lemma 18.18.

We remark that in (18.22) we take the sum over β with $\mu(\beta) = 2$. On the other hand,

$$\dim([0,1] \times \mathcal{M}^{\mathrm{main}}_{0;1}(\beta)) = n + \mu(\beta) + 1 - 3 + 1 = \mu(\beta) + 1. \tag{18.23}$$

(Here we use $n = 2$.) Therefore, in case $\mu(\beta) = 0, 2$, this moduli space can be nonempty and

$$\begin{aligned}
&\mathrm{ev}_{0*}\left(([0,1] \times \mathcal{M}^{\mathrm{main}}_{0;1}(\beta))^{\mathfrak{par}}\right) \\
&\in C^\infty([0,1], H^{2-\mu(\beta)}(L(u);\mathbb{R})) \oplus dt \wedge C^\infty([0,1], H^{1-\mu(\beta)}(L(u);\mathbb{R})).
\end{aligned}$$

We put the left hand side as

$$c(\beta)(t) + dt \wedge e(\beta)(t).$$

Now let $b_+ \in H^1(L(u);\Lambda_+)$ and put

$$b_+ = \sum_{\beta:\mu(\beta)=2} T^{\beta\cap\omega/2\pi} b_\beta.$$

We solve the ordinary differential equation

$$\frac{db_\beta(t)}{dt} + \sum_{\substack{\beta_1,\beta_2\\ \beta_1+\beta_2=\beta,\mu(\beta_1)=0}} e(\beta_1)(t) \wedge b_{\beta_2}(t) = 0$$

with initial value $b_\beta(0) = b_\beta$.

We can prove that

$$\mathfrak{F}_*(b_+) = \sum_{\beta:\mu(\beta)=2} T^{\beta\cap\omega/2\pi} b_\beta(1).$$

(There is a related argument in Auroux [Aur1].)

19. Operator $\mathfrak{p}$ in the toric case

We now start the definition of $\mathfrak{p}$. Consider the moduli space $\mathcal{M}^{\text{main}}_{k;\ell+1}(\beta)$ and the evaluation map

$$\text{ev} = (\text{ev}^{\text{int}}, \text{ev}, \text{ev}^+_0) : \mathcal{M}^{\text{main}}_{k;\ell+1}(\beta) \to X^\ell \times L(u)^k \times X.$$

We remark that the 0-th marked point is an *interior* marked point in this case. So the target of ev^+_0 is X. Let $\mathbf{p} = (\mathbf{p}(1), \dots, \mathbf{p}(\ell))$ be T^n invariant cycles as in Section 6. We put

$$\mathcal{M}^{\text{main}}_{k;\ell+1}(\beta;\mathbf{p}) = \mathcal{M}^{\text{main}}_{k;\ell+1}(\beta)\,{}_{\text{ev}^{\text{int}}} \times (\mathbf{p}(1) \times \cdots \times \mathbf{p}(\ell)).$$

Lemma 19.1. *The moduli spaces $\mathcal{M}^{\text{main}}_{k;\ell+1}(\beta;\mathbf{p})$ have T^n-equivariant Kuranishi structures with boundaries and corners. There exists a system of families of multisections $\mathfrak{p}$ on them such that:*

1. *It is T^n-equivariant.*
2. *It is invariant under the cyclic permutation of boundary marked points.*
3. *It is invariant under the forgetful map of boundary marked points.*
4. *It is transversal to zero.*
5. $\text{ev}^+_0 : \mathcal{M}^{\text{main}}_{k;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}} \to X$ *is a submersion.*
6. *If $k+1>1$, then its boundary is described by the following product:*
$$\mathcal{M}^{\text{main}}_{k_1+1;\ell_1}(\beta_1;\mathbf{p}_1)^{\mathfrak{c}}\,{}_{\text{ev}_0}\times_{\text{ev}_i} \mathcal{M}^{\text{main}}_{k_2;\ell_2+1}(\beta_2;\mathbf{p}_2)^{\mathfrak{p}}$$
Here $\beta_1+\beta_2=\beta$, $k_1+k_2=k$, $(\mathbf{p}_1,\mathbf{p}_2) = \text{Split}(\mathbf{p},(\mathbb{L}_1,\mathbb{L}_2))$, $|\mathbf{p}_i| = \ell_i$, $i = 1,\dots,k_2$.
7. *If $k+1=1$, there is another type of boundary component*
$$\mathcal{M}^{\text{main}}_{0;\ell+2}(\beta;\mathbf{p})^{\mathfrak{p}}\,{}_{\text{ev}_{\ell+1}}\times_X L(u).$$

The lemma can be proved in the same way as [Fu2] Theorem 3.1 and Corollary 3.1. (The statement on T^n invariance is proved in the same way as in Sections 28 - 30 of this paper.) We remark that on the 'bubble' components in (6) we always use the $\mathfrak{c}$-perturbation.

To eliminate the boundary of type (7) we proceed as follows. (We remark that in the present toric case the homology class of $[L(u)]$ is zero so the Gromov-Witten term in Theorem 17.6 (2) drops.) We take a T^n-invariant submanifold $\widehat{L}$ that bounds $L(u)$ as follows. Take a path C joining u to a point u' lying in the interior of a codimension one boundary of P. We put $\widehat{L} = \pi^{-1}(C)$. We now consider

$$\mathcal{M}^{\text{main}}_{0;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}} \cup \mathcal{M}^{\text{main}}_{0;\ell+2}(\beta;\mathbf{p}\otimes\widehat{L})^{\mathfrak{p}}. \tag{19.1}$$

The second term cancels the boundary of type (7) in Lemma 19.1. Moreover the T^n-symmetry is not broken. From now on we change the notation and denote the space (19.1) by $\mathcal{M}^{\rm main}_{0;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}}$. (Note the case $k+1=0$ never appears in the boundary of the other moduli spaces. So changing the notation does not affect the other moduli spaces.) Thus we can forget the boundary of type (7) from now on.

To study Poincaré duality (17.5), it is convenient to use de Rham complex of X as the target of the operator $\mathfrak{p}$. This leads us to use a continuous family of multisections. We remark that we have already chosen multisections on $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})^{\mathfrak{c}}$.

Definition 19.2. We define $\mathfrak{p}_{\ell;k;\beta} : E_\ell(\mathcal{A}(\Lambda_0)[2]) \otimes B_k^{\rm cyc}(H(L(u);\Lambda_0)[1]) \to \Omega(X)$ by

$$\mathfrak{p}_{\ell;k;\beta}([\mathbf{p}];[h_1\otimes\cdots\otimes h_k]) = {\rm ev}^+_{0*}({\rm ev}^*(h_1\times\cdots\times h_k);\mathcal{M}^{\rm main}_{k;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}}). \tag{19.2}$$

(We use Convention 6.12.) The right hand side is invariant under the choice of representative $h_1\otimes\cdots\otimes h_k$ but depends only on its equivalence class in $B_k^{\rm cyc}(H(L(u);\Lambda_0)[1])$ by Lemma 19.1 (2).

We put

$$\mathfrak{p}^{(u,b)}_{\ell;k} = \sum_{\ell=0}^{\infty}\sum_{\beta\in H_2(X;L(u);\mathbb{Z})} T^{\beta\cap\omega/2\pi} y(u)_1^{\partial\beta\cap\mathbf{e}_1}\cdots y(u)_n^{\partial\beta\cap\mathbf{e}_n}\mathfrak{p}_{\ell;k;\beta}.$$

This defines a map $E_\ell\mathcal{A}(\Lambda_0)[2]\otimes B_k^{\rm cyc}(H(L(u);\Lambda_0)[1]) \to \Omega(X)\,\widehat{\otimes}\,\Lambda\langle\langle y(u), y(u)^{-1}\rangle\rangle$.

When $\mathbf{p}$ contains $\mathbf{f}_0 = PD[X] = \mathbf{e}_X$, we use (4) of Theorem 17.6 as the definition of $\mathfrak{p}$.

Here we fix u. (In Sections 19-21 it is unnecessary to move u and regard the operations as the ones over $\Lambda_0^{\overset{\circ}{P}}\langle\langle y, y^{-1}\rangle\rangle$.)

Theorem 19.3. *The map* $\mathfrak{p}$ *defined above satisfies Theorem 17.6* (1) – (5)*, where we take* $\mathfrak{q}^{\mathfrak{c}}$ *instead of* $\mathfrak{q}$*. Moreover in* (2) *we may take* $\widetilde{GW}_{0,\ell+1}(X) = 0$.

Proof. (1) & (2): These follow from Lemma 19.1 and the definition. (We use Lemmata 12.18 and 12.20 [FOOO4] also.)

(3) Note $\mathbf{e}_L = 1 \in H^0(L(u);\Lambda_0)$. On the other hand, (3) of Lemma 19.1 implies that the following diagram commutes:

$$\begin{tikzcd}
\mathcal{M}^{\rm main}_{k;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}} \arrow[r, "{\rm ev}_0^+"] \arrow[d, "\mathfrak{forget}"] & X \arrow[d, equal] \\
\mathcal{M}^{\rm main}_{k-1;\ell+1}(\beta;\mathbf{p})^{\mathfrak{p}} \arrow[r, "{\rm ev}_0^+"] & X
\end{tikzcd} \tag{19.3}$$

(3) follows easily.

(4) is a consequence of the definition.

(5) follows from the fact that the moduli space for $\beta = 0$ contributes only to $\mathfrak{p}_{2,0}$ which gives the usual wedge product. □

Now we use $\mathfrak{p}^{(u,b)}$ to define

$$i_{*,{\rm qm},(\mathfrak{b},b,u)} : H(L(u);\Lambda_0) \to \Omega(X).$$

Write $\mathfrak{b} = \mathfrak{b}_2 + \mathfrak{b}_{\rm high}$ where $\mathfrak{b}_2$ has degree 2 and $\mathfrak{b}_{\rm high}$ has degree higher than 2.

Definition 19.4. We define

$$i_{*,\mathrm{qm},(\mathfrak{b},b,u)}(h) = \sum_{\beta}\sum_{\ell} \frac{T^{\beta\cap\omega/2\pi}}{\ell!} \exp{(\beta\cap\mathfrak{b}_2)}\rho(\partial\beta)\mathfrak{p}_{\ell;1;\beta}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell};h).$$

Here ρ is the representation of $\pi_1(L(u))$ obtained from b_0 as before.

Remark 19.5. If $b = b_0 + b_+$ where b_0 is of degree 1 and the degree of b_+ is > 1, then we have

$$i_{*,\mathrm{qm},(\mathfrak{b},b,u)}(h) = \sum_{\beta}\sum_{\ell,k} \frac{T^{\beta\cap\omega/2\pi}}{\ell!} \exp{(\beta\cap\mathfrak{b}_2)}\exp(b_0\cap\partial\beta)\mathfrak{p}_{\ell;k+1;\beta}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell};[h\otimes b_+^{\otimes k}]).$$

This expression is closer to (17.13) than the above definition. See Section 27.

Lemma 19.6. *$i_{*,\mathrm{qm},(\mathfrak{b},b,u)}$ is a chain map.*

The proof is the same as the proof of Lemma 17.9. Thus we have defined

$$i_{\#,\mathrm{qm},(\mathfrak{b},b,u)} : HF((L(u);(\mathfrak{b},b)),(L(u);(\mathfrak{b},b));\Lambda_0) \to H(X;\Lambda_0). \tag{19.4}$$

We now consider the pairing $\langle\cdot,\cdot\rangle_{\mathrm{PD}_{L(u)}}$ by

$$\langle h',h\rangle_{\mathrm{PD}_{L(u)}} = \int_{L(u)} h'\wedge h \tag{19.5}$$

and $\langle\cdot,\cdot\rangle_{\mathrm{PD}_X}$ by

$$\langle Q,v\rangle_{\mathrm{PD}_X} = \int_Q v = \int_X PD_X(Q)\wedge v. \tag{19.6}$$

We also recall the map

$$i^{\#\mathfrak{c}}_{\mathrm{qm},(\mathfrak{b},b,u)} : H(X;\Lambda_0) \to H(L(u);\Lambda_0)$$

introduced in Section 18.

Lemma 19.7. *For any $Q\in\mathcal{A}(X)$, the element $i^{\#\mathfrak{c}}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q)$ is contained in $\Lambda_0\cdot\{e_L\}$.*

Proof. By definition, $i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q)$ is proportional to the unit which is a degree-zero form. (See Lemma 17.1.) Since $\mathfrak{F}$ is unital filtered A_∞ homomorphism $\mathfrak{F}_*$ sends the unit to the unit. Lemma 19.7 then follows from Lemma 18.29. □

The following theorem describes the relationship between the maps $i^{\mathfrak{c}\#}_{\mathrm{qm},(\mathfrak{b},b,u)}$ and $i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}$.

Theorem 19.8. *Let $Q\in\mathcal{A}(X)$ and $h\in H^*(L(u);\Lambda_0)$. Then we have*

$$\langle i^{\mathfrak{c}\#}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q),h\rangle_{\mathrm{PD}_{L(u)}} = \langle Q, i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(h)\rangle_{\mathrm{PD}_X}. \tag{19.7}$$

Corollary 19.9. *If the degree n component of h is zero, then the right hand side of (19.7) is zero.*

Proof. This follows from Theorem 19.8 and Lemma 19.7. □

Proof of Theorem 19.8. Let $\mathbf{p}\in E_\ell(\mathcal{A}[2])$. We put

$$\Delta\mathbf{p} = \sum_c \mathbf{p}_{1;c}\otimes\mathbf{p}_{2;c},$$

where $\mathbf{p}_{i;c}\in E_{\ell_{c;i}}(\mathcal{A}[2])$. The following is the key geometric result on which the proof of (19.7) is based.

Lemma 19.10. $[0,1]\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(\beta;\mathbf{p})$ *has a Kuranishi structure and its boundary is the union of following* 4 *types of fiber products:*

1. $\{0\}\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(\beta;\mathbf{p})$.
2. $\{1\}\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(\beta;\mathbf{p})$.
3. $\mathcal{M}_{k_1+1;\ell_{c;1}}(\beta_1;\mathbf{p}_{1;c})\,{}_{\mathrm{ev}_0}\times_{\mathrm{ev}_j}\Big([0,1]\times\mathcal{M}^{\mathrm{main}}_{k_2+2;\ell_{c;2}+1}(\beta_2;\mathbf{p}_{2;c})\Big)$. *Here* $k_1+k_2=k$, $\beta_1+\beta_2=\beta$, $j=1,\dots,k_2+1$.
4. $\big([0,1]\times\mathcal{M}_{k_1+1;\ell_{c;1}+1}(\beta_1;\mathbf{p}_{1;c})\big)\,{}_{\mathrm{ev}_0}\times_{\mathrm{ev}_j}\mathcal{M}^{\mathrm{main}}_{k_2+2;\ell_{c;1}}(\beta_2;\mathbf{p}_{2;c})$. *Here* $k_1+k_2=k$, $\beta_1+\beta_2=\beta$, $j=1,\dots,k_2$.

And there exists a system $\mathfrak{parap}$ *of a continuous family of multisections on* $[0,1]\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell}(\beta;\mathbf{p})$ *with the following properties:*

(a) *It is transversal to zero and* T^n*-invariant.*
(b) *It is invariant under the cyclic permutation of the boundary marked points and under the permutations of interior marked points.*
(c) *It is compatible with the forgetful map of the boundary marked points.*
(d) *On the boundary component of type* (1) *above it coincides with* $\mathfrak{c}$ *perturbation and one of type* (2) *above it coincides with* $\mathfrak{p}$ *perturbation.*
(e) *On the boundary component of type* (3) *above it is compatible. Here we put the* $\mathfrak{c}$ *perturbation for the first factor.*
(f) *On the boundary component of type* (4) *above it is compatible. Here we put the* $\mathfrak{c}$ *perturbation for the second factor.*

Proof. We remark that the 0-th marked point is the interior marked point when we define $\mathfrak{p}$ and $\mathfrak{parap}$. So when we consider the codimension one boundary of $\mathcal{M}^{\mathrm{main}}_{k,\ell+1}(\beta;\mathbf{p})$ the $\mathfrak{p}$ or $\mathfrak{parap}$-perturbation is applied to the factor containing the 0-th marked point. Once this point is understood, the proof is the same as other similar statements we discussed many times before. ☐

Going back to the proof of the theorem, we apply Stokes' formula and obtain

$$\begin{aligned}&\int_{\partial([0,1]\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(\beta;\mathbf{p}))^{\mathfrak{parap}}}(\mathrm{ev},\mathrm{ev}_0)^*((b_+^{\mathfrak{c}})^{\otimes k}\times h)\\ &=\int_{([0,1]\times\mathcal{M}^{\mathrm{main}}_{k+1;\ell+1}(\beta;\mathbf{p}))^{\mathfrak{parap}}}(\mathrm{ev},\mathrm{ev}_0)^*(d((b_+^{\mathfrak{c}})^{\otimes k}\times h))=0.\end{aligned}\tag{19.8}$$

Remark 19.11. In the proofs of the identity (19.8) and of similar others, we use the integration of a differential form (or integration along the fiber of a map) on the zero set of a *continuous family of* multisections. The dimension of this zero set is given by the sum of the (virtual) dimension of the space with Kuranishi structure and that of the parameter space W. (Note W is defined only locally.) We choose a (T^n-invariant) smooth form ω with compact support on W that has degree $\dim W$ and total mass 1, and do integration of a differential form after we take a wedge product with this form. So strictly speaking, the integrand of the left hand side of (19.8) should actually be $(\mathrm{ev},\mathrm{ev}_0)^*((b_+^{\mathfrak{c}})^{\otimes k}\times h)\wedge\omega)$, for example. By an abuse of notations, however, we write this integration just as (19.8) here and henceforth. See [FOOO5] Section 12 for the precise definition.

We decompose the left hand side of (19.8) according to the decomposition of the boundary to (1) - (4) of Lemma 19.10. Let us write the term corresponding to (m) as

$$C(\beta;\mathbf{p};k;\ell;m),\quad m=1,2,3,4.$$

We put

$$\mathfrak{b}_{\rm high}^{\otimes\ell} = \sum_{I\in\mathfrak{I}_\ell} c_I \mathbf{f}_{i_1}\otimes\cdots\otimes\mathbf{f}_{i_\ell} = \sum_{I\in\mathfrak{I}_\ell} c_I\mathbf{f}_I. \tag{19.9}$$

Here $\mathfrak{I}_\ell$ is the set of multi-indices $I = (i_1,\dots,i_\ell)$ and $c_I\in\Lambda_0$.

We now calculate

$$\sum_{\ell,k,\beta}\sum_{I\in\mathfrak{I}_\ell}\frac{c_I}{\ell!}\rho(\partial\beta)e^{\mathfrak{b}_2\cap\omega}C(\beta;\mathbf{f}_I\otimes Q;k;\ell;m) \tag{19.10}$$

for $m = 1,2,3,4$.

The cases $m=1$ and $m=2$ give rise to the left and the right hand sides of Theorem 19.8 by Lemma 19.10 (d) respectively.

Let us consider the case $m=3$. Decompose $I = I_1\sqcup I_2$ for $I_j\in\mathfrak{I}_{\ell_j}$, $j=1,2$ with $\ell_1+\ell_2=\ell$ and

$$k = k_1+k_2,\ \beta=\beta_1+\beta_2,\ \mathbf{p}=\mathbf{p}_1\otimes\mathbf{p}_2.$$

We put

$$E = \sum_{\beta_1,\ell_1,k_1}\sum_{I_1\in\mathfrak{I}_{\ell_1}}\frac{c_{I_1}}{\ell!}{\rm ev}_{0*}\left(({\rm ev})^*((b_+^{\mathfrak{c}})^{\otimes k_1});\mathcal{M}_{k_1+1;\ell_1+1}^{\rm main}(\beta;\mathbf{f}_I\times Q)^{\mathfrak{c}}\right). \tag{19.11}$$

This is an element of $H(L(u);\Lambda_0)$. Then thanks to $\Delta(e^{\mathfrak{b}}) = e^{\mathfrak{b}}\otimes e^{\mathfrak{b}}$, the case $m=3$ of (19.10) becomes:

$$\begin{aligned}&\sum_{\beta_2,\ell_2,k_2}\sum_{j=1}^{k_2}\sum_{I_2\in\mathfrak{I}_{\ell_2}}\frac{c_{I_2}}{\ell!}\rho(\partial\beta)e^{\mathfrak{b}_2\cap\omega}\\ &\int_{([0,1]\times\mathcal{M}_{k_2+1;\ell_2+1}^{\rm main}(\beta_2;\mathbf{p}_2))^{\mathfrak{parap}}}({\rm ev},{\rm ev}_0)^*((b_+^{\mathfrak{c}})^j\times E\times(b_+^{\mathfrak{c}})^{k_2-j}\times h).\end{aligned} \tag{19.12}$$

In the same way as the proof of Lemma 19.7, we can prove that E is proportional to the unit. Therefore Lemma 19.10 (c) implies that (19.12) is 0. Namely (19.10) is zero for the case $m=3$.

We finally consider the case $m=4$. We define

$$\begin{aligned}F = &\sum_{\beta_1,\ell_1,k_1}\sum_{I_1\in\mathfrak{I}_{\ell_1}}\frac{c_{I_1}}{\ell_1!}\rho(\partial\beta)e^{\mathfrak{b}_2\cap\omega}\\ &{\rm ev}_{0*}\left({\rm ev}^*(b_+^{\mathfrak{c}})^{k_1};([0,1]\times\mathcal{M}_{k_1+1;\ell_1+1}^{\rm main}(\beta;\mathbf{f}_{I_1}\times Q)^{\mathfrak{parap}}\right).\end{aligned} \tag{19.13}$$

Then, by using $\Delta(e^{\mathfrak{b}}) = e^{\mathfrak{b}}\otimes e^{\mathfrak{b}}$, the case $m=4$ of (19.10) is

$$\begin{aligned}&\sum_{\beta_2,\ell_2,k_2}\sum_{I_2\in\mathfrak{I}_{\ell_2}}\sum_{j=1}^{k_2}\frac{c_{I_2}}{\ell_2!}\rho(\partial\beta)e^{\mathfrak{b}_2\cap\omega}\\ &{\rm ev}_{0*}\left(({\rm ev})^*((b_+^{\mathfrak{c}})^j\times F\times(b_+^{\mathfrak{c}})^{(k_2-j)}\times h);\mathcal{M}_{k_2+1;\ell_2+1}^{\rm main}(\beta;\mathbf{f}_{I_2})^{\mathfrak{c}}\right).\end{aligned} \tag{19.14}$$

By definition this is

$$\langle\delta_{\mathfrak{c}}^{\mathfrak{b},b^{\mathfrak{c}}}(F),h\rangle_{{\rm PD}_{L(u)}}$$

and is zero. This is because $(u,b^{\mathfrak{c}})$ is a critical point of $\mathfrak{PO}_{\mathfrak{b}}^{\mathfrak{c}}$ and so $\delta_{\mathfrak{c}}^{\mathfrak{b},b^{\mathfrak{c}}}=0$ and Corollary 18.22. The proof of Theorem 19.8 is complete. □

Remark 19.12. We used cyclic symmetry in the proof of Theorem 19.8 above. In other words the above proof does not work to prove the same conclusion (19.7) when we replace the left hand side by $\langle i^{\#}_{\mathrm{qm},(\mathfrak{b},b,u)}(Q), h\rangle_{\mathrm{PD}_{L(u)}}$.

The reason is as follows.

To construct the $\mathfrak{p}$ perturbation which is compatible to $\mathfrak{q}$ perturbation, we need to take the $\mathfrak{q}$ perturbation for the first factor in Lemma 19.1 (6). Moreover we have to take the 0-th marked point of this factor to the point which intersects with the second factor. (Namely the point which becomes the sigular point after gluing.)

On the other hand, for the Formula (19.7) to hold, we need to take the 0-th marked point so that we pull back the form h by ev at the 0-th marked point.

We cannot realize both of them at the same time in the situation of Lemma 19.10 (4).

20. Moduli space of holomorphic annuli

Let $(u,b) \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$. We take a basis $\mathbf{e}_I$ with $I \in 2^{\{1,\dots,n\}}$ of $H(L(u);\mathbb{Q})$. Let

$$g_{IJ} = \langle \mathbf{e}_I, \mathbf{e}_J\rangle_{\mathrm{PD}_{L(u)}}, \qquad [g^{IJ}] = [g_{IJ}]^{-1}.$$

Here $[g^{IJ}]$, $[g_{IJ}]$ are $2^n \times 2^n$ matrices. We recall that the operator

$$\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b} : HF((L(u),\mathfrak{b},b),(L(u),\mathfrak{b},b);\Lambda_0)^{\otimes 2} \to HF((L(u),\mathfrak{b},b),(L(u),\mathfrak{b},b);\Lambda_0)$$

is defined by

$$\begin{aligned}\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b}(h_1,h_2) = &\sum_{k_1,k_2,k_3}\sum_{\ell,\beta} T^{\beta\cap\omega/2\pi}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta) \\ &\mathfrak{q}^{\mathfrak{c}}_{\ell;2+k_1+k_2+k_3;\beta}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell};(b^{\mathfrak{c}}_+)^{k_1}\times h_1\times(b^{\mathfrak{c}}_+)^{k_2}\times h_2\times(b^{\mathfrak{c}}_+)^{k_3}).\end{aligned} \tag{20.1}$$

(See (18.5), (18.6).) In the next theorem, we involve the harmonic forms on $L(u)$ of all degrees beside those of top degree n, although in the end only the top degree forms turn out to give non-trivial contributions in the formula below.

Theorem 20.1. *Let $\mathfrak{v},\mathfrak{w} \in H(L(u);\Lambda_0)$. Then we have*

$$\begin{aligned}&\langle i_{\#,\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}})}(\mathfrak{v}), i_{\#,\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}})}(\mathfrak{w})\rangle_{\mathrm{PD}_X} \\ &= \sum_{I,J\in 2^{\{1,\dots,n\}}} (-1)^* g^{IJ}\langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}_I,\mathfrak{v}), \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}_J,\mathfrak{w})\rangle_{\mathrm{PD}_{L(u)}}.\end{aligned} \tag{20.2}$$

Here $* = \frac{n(n-1)}{2}$ in case the degree of $\mathfrak{v},\mathfrak{w}$ are n. We prove it in Subsection 26.2. (See (26.21).) For the sign $*$ in the general case, see Proposition 26.17.

Remark 20.2. We do not assume nondegeneracy of the critical point (u,b) of $\mathfrak{PO}_{\mathfrak{b}}$ in Theorem 20.1. Theorem 22.2 implies that, in the case of degenerate (u,b) and (1) or (2) of Condition 2.19 is satisfied, the right hand side of (20.2) is 0 and so the left hand side is also 0.

The right hand side is closely related to the invariant $Z(\mathfrak{b},b)$ in Definition 2.23.

The proof of Theorem 20.1 is based on the study of the moduli space of holomorphic maps from the annuli. We begin with defining the corresponding moduli space.

Consider the bordered semi-stable curve Σ of genus 0 with two boundary components $\partial_1\Sigma$, $\partial_2\Sigma$ and k_1+1, k_2+1 boundary marked points $\vec{z}_1 = (z_{1,0}, z_{1,1},\dots,z_{1,k_1})$,

$\vec{z}_2 = (z_{2,0}, z_{2,1}, \dots, z_{2,k_2})$ on each of the components $\partial_1\Sigma$, $\partial_2\Sigma$ of the boundary, and ℓ interior marked points $\vec{z}^{\,+} = (z_1^+, \dots, z_\ell^+)$. We denote it by $(\Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ or sometimes by Σ for short. We say that $(\Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ is in the *main component* if $\vec{z}_i$ respects the cyclic order of the boundary. (When we regard Σ as an annulus in $\mathbb{C}$, we take the counter clockwise cyclic order for the outer circle and the clockwise cyclic order for the inner circle.) Let $(\Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ be as above and consider a holomorphic map $w : (\Sigma, \partial\Sigma) \to (X, L(u))$. We say $(w; \Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ is a *stable map* if its automorphism is trivial in an obvious sense.

Definition 20.3. We put $\beta \in H_2(X, L(u); \Lambda_0)$. We denote by

$$\mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}(\beta)$$

the set of all isomorphism classes of stable maps $(w; \Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ such that $w_*[\Sigma] = \beta$ and $(\Sigma; \vec{z}_1, \vec{z}_2, \vec{z}^{\,+})$ is in the main component, which we described above.

In the same way as Section 3 [FOOO1] and Section 7.1 [FOOO3] we can prove that $\mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}$ has a Kuranishi structure with boundary and corners. There are obvious evaluation maps

$$\begin{aligned} {\rm ev} =& ({\rm ev}^1, {\rm ev}^2, {\rm ev}^{\rm int}, {\rm ev}^1_0, {\rm ev}^2_0) \\ &: \mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}(\beta) \to L(u)^{k_1} \times L(u)^{k_2} \times X^2 \times L(u) \times L(u). \end{aligned}$$

Here ${\rm ev}^1_0$, ${\rm ev}^2_0$ are the evaluation maps at z^1_0, z^2_0, and ${\rm ev}^1$, ${\rm ev}^2$ are evaluation maps at $(z_{1,1}, \dots, z_{1,k_1})$ and $(z_{2,1}, \dots, z_{2,k_2})$, respectively. We consider the case $k_1 \ge 0$, $k_2 \ge 0$ only, in this paper.

Let $\mathcal{M}^{\rm ann;main}_{(1,1);0}$ be the moduli space $\mathcal{M}^{\rm ann;main}_{(0+1,0+1);0}(0)$ in the case $X = L(u)$ =one point. There exists an obvious forgetful map

$$\mathfrak{forget} : \mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}(\beta) \to \mathcal{M}^{\rm ann;main}_{(1,1);0}. \tag{20.3}$$

(Here we forget $(z_{1,1}, \dots, z_{1,k_1})$ and $(z_{2,1}, \dots, z_{2,k_2})$ among the boundary marked points.)

Convention 20.4. Hereafter we write

$$\mathfrak{forget}^{-1}(X) \cap \mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}(\beta)$$

for the inverse image of $X \subset \mathcal{M}^{\rm ann;main}_{(1,1);0}$ by (20.3).

Lemma 20.5. *The boundary of $\mathcal{M}^{\rm ann;main}_{(k_1+1,k_2+1);\ell}(\beta)$ is the union of the following* 5 *types of fiber products:*

(1)
$$\mathcal{M}_{k'_1+1;\ell'}(\beta')^{\mathfrak{c}}\,{}_{{\rm ev}_0}\times_{{\rm ev}^1_i} \mathcal{M}^{\rm ann;main}_{(k''_1+1,k_2+1);\ell''}(\beta'').$$
Here $k'_1 + k''_1 = k_1 + 1$, $\ell' + \ell'' = \ell$, $\beta' + \beta'' = \beta$, $i = 1, \dots, k''_1$.

(2)
$$\mathcal{M}_{k'_2+1;\ell'}(\beta')^{\mathfrak{c}}\,{}_{{\rm ev}_0}\times_{{\rm ev}^2_i} \mathcal{M}^{\rm ann;main}_{(k_1+1,k''_2+1);\ell''}(\beta'').$$
Here $k'_2 + k''_2 = k_2 + 1$, $\ell' + \ell'' = \ell$, $\beta' + \beta'' = \beta$, $i = 1, \dots, k''_2$.

(3)
$$\mathcal{M}^{\rm ann;main}_{(k'_1+1,k_2+1);\ell'}(\beta')\,{}_{{\rm ev}^1_0}\times_{{\rm ev}_i} \mathcal{M}_{k''_1+1;\ell''}(\beta'')^{\mathfrak{c}}.$$
Here $k'_1 + k''_1 = k_1 + 1$, $\ell' + \ell'' = \ell$, $\beta' + \beta'' = \beta$, $i = 1, \dots, k''_1$.

(4)
$$\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2'+1);\ell'}(\beta')\ {}_{\mathrm{ev}_0^2}\times_{\mathrm{ev}_i} \mathcal{M}_{k_2''+1;\ell''}(\beta'')^{\mathfrak{c}}.$$
Here $k_2'+k_2''=k_1+1$, $\ell'+\ell''=\ell$, $\beta'+\beta''=\beta$, $i=1,\dots,k_2''$.

(5) $\mathfrak{forget}^{-1}(\partial\mathcal{M}^{\text{ann;main}}_{(1,1);0})\cap\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta)$.

This is obvious from the construction of the Kuranishi structure. We remark that we put $\mathfrak{c}$ in the first factors of (1),(2) and third factors of (3),(4). This is to clarify we always use the '$\mathfrak{c}$' perturbation for such factor. This holds for many other similar cases. Namely we use the perturbation '$\mathfrak{c}$' for component of the disk bubble.

For $\mathbf{p}$ as in Section 6, we define

$$\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})=\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta)_{\mathrm{ev}^{\mathrm{int}}}\times_{X^\ell}(\mathbf{p}(1)\times\cdots\times\mathbf{p}(\ell)). \tag{20.4}$$

We can describe its boundary by using Lemma 20.5, in the way similar to the cases we have already discussed in several other occasions.

In a way similar to Lemma 9.3 we can describe $\mathcal{M}^{\text{main}}_{(1,1);0}$ as follows.

Lemma 20.6. $\mathcal{M}^{\text{ann;main}}_{(1,1);0}$ *has a stratification such that the interior of each stratum are as follows.*

(1) $\operatorname{Int}D^2\setminus\{0\}$.
(2) *An arc* $(-1,1)$.
(3) *Two points.* $[\Sigma_1]$, $[\Sigma_2]$.

The point $0\in D^2$ *becomes* $[\Sigma_1]$ *in the closure of the stratum* (1). *The two boundary points* ±1 *in the closure of* $(-1,1)$ *become* $[\Sigma_2]$. *Thus the union of the two strata* $(-1,1)$, $[\Sigma_2]$ *becomes a circle, which turns out to be* ∂D^2 *in the closure of the stratum* (1).

Each of the above strata corresponds to the combinatorial type of the elements of $\mathcal{M}^{\text{main}}_{1;2}$.

Proof. We first define two points $[\Sigma_1]$, $[\Sigma_2]$ of $\mathcal{M}^{\text{ann;main}}_{(1,1);0}$.

Let D_1^2 and D_2^2 be two copies of disk $D^2=\{z\in\mathbb{C}\mid |z|\le 1\}$. We identify $0\in D_1^2$ and $0\in D_2^2$ to obtain Σ_1. We put $\partial_1\Sigma_1=\partial D_1^2$, $\partial_2\Sigma_1=\partial D_2^2$, $z_0^1=1\in\partial D_1^2$, $z_0^2=1\in\partial D_2^2$. Thus we have $(\Sigma_1;z_1^1,z_2^1)\in\mathcal{M}^{\text{ann;main}}_{(1,1);0}$.

We next define Σ_2. We consider two copies D_1^2, D_2^2 of disk. We put $\mathfrak{z}_k^i=e^{2k\pi\sqrt{-1}/3}\in\partial D_i^2$ for $i=1,2$, $k=0,1,2$.

We identify $\mathfrak{z}_1^1$ with $\mathfrak{z}_1^2$ and $\mathfrak{z}_2^1$ with $\mathfrak{z}_2^2$ respectively and obtain Σ_2. We put $\mathfrak{z}_0^i=z_i^1$ and $i=1,2$. We call $\partial_i\Sigma_2$ the component which contains z_i^1. We thus defined $(\Sigma_2;z_1^1,z_2^1)\in\mathcal{M}^{\text{ann;main}}_{(1,1);0}$.

We next describe the stratum (2). We consider the moduli space $\mathcal{M}^{\text{main}}_{4;0}$ of disks with 4 boundary marked points without interior marked points. It is an arc $[0,1]$. Let $(D^2;\mathfrak{z}_1,\mathfrak{z}_2,\mathfrak{z}_3,\mathfrak{z}_4)$ be a point in the interior of $\mathcal{M}^{\text{main}}_{4;0}$. We identify $\mathfrak{z}_2$ with $\mathfrak{z}_4$ to obtain Σ and put $z_1=\mathfrak{z}_1$, $z_2=\mathfrak{z}_4$. Then $(\Sigma;z_1,z_2)\in\mathcal{M}^{\text{ann;main}}_{(1,1);0}$.

It is easy to see that as $(D^2;\mathfrak{z}_1,\mathfrak{z}_2,\mathfrak{z}_3,\mathfrak{z}_4)$ approaches to the boundary of $\mathcal{M}^{\text{main}}_{4;0}$ $(\Sigma;z_1,z_2)$ converges to $[\Sigma_2]$ in $\mathcal{M}^{\text{ann;main}}_{(1,1);0}$. Thus the union of the strata (2), (3) we have described above becomes a circle. It is easy to see that it gives the stratification as in Lemma 20.6. □

The proof of the next lemma is easy.

Lemma 20.7. *The forgetful map* (*20.3*) *is continuous and is a smooth submersion on each stratum.*

The proof of Theorem 20.1 consists of a series of propositions. We recall that, for $h_1, \dots, h_k \in H(L(u);\Lambda_0)$, the symbol $[h_1, \dots, h_k]$ denotes the equivalence class of $h_1 \otimes \cdots \otimes h_k \in B_k(H(L(u);\Lambda_0)[1])$ in $B_k^{\mathrm{cyc}}(H(L(u);\Lambda_0)[1])$. We also write

$$[\mathfrak{v}; k] = [\underbrace{b^{\mathfrak{c}}_{+}, \dots, b^{\mathfrak{c}}_{+}}_{k}, \mathfrak{v}], \tag{20.5}$$

for the simplicity of notations.

Proposition 20.8.

$$\begin{aligned}
&\sum_{\beta}\sum_{\ell}\sum_{k_1,k_2} \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta) \\
&\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]) \\
&= \sum_{\beta_1+\beta_2=\beta}\ \sum_{\substack{\ell_1,\ell_2\\ :\ell_1+\ell_2=\ell}}\ \sum_{k_1,k_2} \frac{T^{(\beta_1+\beta_2)\cap\omega}}{\ell_1!\ell_2!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial(\beta_1+\beta_2)) \\
&\qquad\qquad\langle \mathfrak{p}_{\ell_1,\beta_1}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_1};[\mathfrak{v};k_1]),\mathfrak{p}_{\ell_2,\beta_2}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_2};[\mathfrak{w};k_2])\rangle_{\mathrm{PD}_X}.
\end{aligned}$$

Proposition 20.9.

$$\begin{aligned}
&\sum_{\beta}\sum_{\ell}\sum_{k_1,k_2} \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta) \\
&\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]) \\
&= \sum_{\beta}\sum_{\ell}\sum_{k_1,k_2} \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta) \\
&\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma_2])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]).
\end{aligned}$$

Proposition 20.10.

$$\begin{aligned}
&\sum_{\beta}\sum_{\ell}\sum_{k_1,k_2} \frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta) \\
&\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma_2])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]) \\
&= \sum_{\beta_1+\beta_2=\beta}\ \sum_{\substack{\ell_1,\ell_2\\ :\ell_1+\ell_2=\ell}}\ \sum_{I,J\in 2^{\{1,\dots,n\}}} g^{IJ}\frac{T^{(\beta_1+\beta_2)\cap\omega}}{\ell_1!\ell_2!}\exp((\beta_1+\beta_2)\cap\mathfrak{b}_2)\rho(\partial(\beta_1+\beta_2)) \\
&\qquad\qquad\langle \mathfrak{q}^{\mathfrak{c},\rho}_{\ell_1;2;\beta_1}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_1};\mathfrak{v}\otimes\mathbf{e}_I),\mathfrak{q}^{\mathfrak{c},\rho}_{\ell_2;2;\beta_2}(\mathfrak{b}_{\mathrm{high}}^{\otimes\ell_2};\mathbf{e}_J\otimes\mathfrak{w})\rangle_{\mathrm{PD}_L}.
\end{aligned}$$

Here in the right hand side of Proposition 20.10, the operator $q^{\mathfrak{c},\rho}_{\ell;2;\beta}$ is defined by

$$\begin{aligned}&\mathfrak{q}^{\mathfrak{c},\rho}_{\ell;2;\beta}([\mathbf{p}];x_1,x_2)\\&=\sum_{k_1,k_2,k_3}\mathfrak{q}^{\mathfrak{c},\rho}_{\ell;2+k_1+k_2+k_3;\beta}([\mathbf{p}];(b^{\mathfrak{c}}_+)^{k_1}\otimes x_1\otimes(b^{\mathfrak{c}}_+)^{k_2}\otimes x_2\otimes(b^{\mathfrak{c}}_+)^{k_3}).\end{aligned}\tag{20.6}$$

It is straightforward to see that Propositions 20.8, 20.9 and 20.10 imply Theorem 20.1. The rest of this section is occupied with the proofs of Propositions 20.8, 20.9 and 20.10. Actually the geometric idea behind the validity of these propositions is clear. The main task is to construct appropriate perturbations (a family of multisections) of the moduli spaces involved so that they are consistent with the perturbations we fixed in the earlier stage to define various other operators considered. We explain the details of this construction in this section. In the course of doing this, we also determine our choice of perturbations made for $\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\rm ann;main}_{(1,1);\ell}(\beta;\mathfrak{b}^{\otimes\ell}_{\rm high})$ and for $\mathfrak{forget}^{-1}([\Sigma_2])\cap\mathcal{M}^{\rm ann;main}_{(1,1);\ell}(\beta;\mathfrak{b}^{\otimes\ell}_{\rm high})$.

Proof of Proposition 20.8. The arguments used in this proof is similar to the arguments used around Lemma 9.8. We start with the following:

Definition 20.11.

$$\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)=\mathcal{M}^{\rm main}_{k_1;\ell_1+1}(\beta_1;\mathbf{p}_1)\ {}_{{\rm ev}^{\rm int}_{\ell_1+1}}\times_{{\rm ev}^{\rm int}_{\ell_2+1}}\mathcal{M}^{\rm main}_{k_2;\ell_2+1}(\beta_2;\mathbf{p}_2).$$

Here the fiber product is taken over X. Since evaluation maps are weakly submersive, this space has a fiber product Kuranishi structure. (See [FOOO3] Section A.1.2.)

Let $\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})=(\mathbf{p}_1,\mathbf{p}_2)$. (See Section 6 for the notation.) We can glue elements of $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$ at the ℓ_1+1-th and ℓ_2+1-th marked points. We then get an element of

$$\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\rm ann;main}_{(k_1,k_2);\ell}(\beta_1+\beta_2;\mathbf{p})$$

where $\ell=\ell_1+\ell_2$. We have thus obtained the map

$$\begin{aligned}\mathrm{Glue}\ :\ &\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)\\&\to\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\rm ann;main}_{(k_1,k_2);\ell}(\beta_1+\beta_2;\mathbf{p}).\end{aligned}\tag{20.7}$$

Let β, k_1,k_2, ℓ be given. We consider the disjoint union of the moduli spaces $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$ where $(\mathbf{p}_1,\mathbf{p}_2)=\mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})$ with $(\mathbb{L}_1,\mathbb{L}_2)\in\mathrm{Shuff}(\ell)$ and $\beta_1+\beta_2=\beta$.

Lemma 20.12. *The map (20.7) defines a map from the above union of the moduli spaces* $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$ *to* $\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\rm ann;main}_{(k_1,k_2);\ell}(\beta;\mathbf{p})$. *It is surjective and is an isomorphism outside the codimension* 2 *strata.*

The proof is the same as the proof of Lemma 9.8 and omitted. The reason why the map in Lemma 20.12 is not an isomorphism is the same as in the case of Lemma 9.4. Namely it should be regarded as a 'normalization' of the 'normal crossing divisor' $\mathfrak{forget}^{-1}([\Sigma_1])$.

Remark 20.13. There occurs a similar problem as the one mentioned in Remark 9.14. We will discuss it later in Section 28.

Lemma 20.14. *There exists a system of continuous family of multisections defined on a neighborhood of* $\mathfrak{forget}^{-1}([\Sigma_1])$ *with the following properties:*

(1) *It is transversal to* 0 *and is* T^n*-invariant.*
(2) *It is compatible at the boundaries described in Lemma 20.5.*
(3) *It is compatible with the forgetful map of the boundary marked points.*
(4) *It is invariant under the cyclic permutations of boundary marked points and arbitrary permutations of the interior marked points.*
(5) *It is transversal to the map* (*20.12*) *in the sense we describe below.*

We now make the statements of Condition (5) precise. Let U be a Kuranishi neighborhood of a point $\mathfrak{x}$ in $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$. Denote

$$\mathrm{Glue}_{k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2}(\mathfrak{x}) = \mathfrak{y} \in \mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta_1+\beta_2;\mathbf{p}).$$

Let U' be a Kuranishi neighborhood of $\mathfrak{y}$ and E' an obstruction bundle. Let $\mathfrak{s}_i(x,w)$ be a branch of our continuous family of multisections. Here $w \in W$ is an element of the parameter space of our continuous family of multisections and $x \in U'$ is an element of the Kuranishi neighborhood. We require that $\mathfrak{s}_i^{-1}(0)$ is transversal to the map : $U \times W \to U' \times W$ induced by $\mathrm{Glue}_{k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2}$.

Proof. The proof can be done by an induction of $\beta \cap \omega$ and the number of boundary marked points in the same way as we already did several times before. ☐

We remark that by Lemma 20.12 or by construction, the obstruction bundle E on the above Kuranishi neighborhood U is the pull-back of E'. Thus the choice of a system of continuous families of multisections satisfying Lemma 20.12 induces a system of continuous families of multisections on $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$. We call this a *pull-back multisection* and denote it by $\mathfrak{back}$. We denote its zero set by

$$\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)^{\mathfrak{back}}.$$

Lemma 20.12 immediately implies the equality

$$\begin{aligned}
&\int_{\mathfrak{forget}^{-1}([\Sigma_1])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})} (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]) \\
&= \sum_{\beta_1+\beta_2=\beta}\ \sum_{(\mathbb{L}_1,\mathbb{L}_2)\in\mathrm{Shuff}(\ell)} \\
&\qquad\qquad \int_{\mathcal{M}(k_1+1,k_2+1;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)^{\mathfrak{back}}} (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]),
\end{aligned} \tag{20.8}$$

where $(\mathbf{p}_1,\mathbf{p}_2) = \mathrm{Split}((\mathbb{L}_1,\mathbb{L}_2),\mathbf{p})$.

We next study the right hand side of Proposition 20.8. Let $\mathbf{p}_i \in E_{\ell_i}(\mathcal{A}[2])$. We assume that they do not contain the degree zero element $\mathbf{f}_0$. We fixed a continuous family of multisections on $\mathcal{M}_{k_i;\ell_i+1}(\beta_i;\mathbf{p}_i)$ when we defined an operator $\mathfrak{p}$. It is chosen so that $\mathrm{ev}^{\mathrm{int}}_{\ell_i+1}$ becomes a submersion. Therefore it induces a continuous family of multisections on $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$. We call this multisection the $\mathfrak{p}$*-multisection* and denote its zero set by $\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)^{\mathfrak{p}}$. Then the next formula is immediate from definition.

$$\begin{aligned}
&\int_{\mathcal{M}(k_1+1,k_2+1;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)^{\mathfrak{p}}} (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2]) \\
&\qquad = \langle \mathfrak{p}_{\ell_1,\beta_1}(\mathbf{p}_1\otimes[\mathfrak{v};k_1]), \mathfrak{p}_{\ell_2,\beta_2}(\mathbf{p}_2\otimes[\mathfrak{w};k_2])\rangle_{\mathrm{PD}_X}.
\end{aligned} \tag{20.9}$$

Lemma 20.15. *There exists a system of continuous families of multisections* $\mathfrak{s}$ *on the space* $[0,1]\times\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1,\mathbf{p}_2)$ *with the following properties.*

1. *It is transversal to* 0 *and is* T^n*-invariant.*
2. *It is invariant under the cyclic permutations of* k_1 *(resp.* k_2*) boundary marked points on each of the boundary components. It is invariant under arbitrary permutations of the interior marked points.*
3. *It is compatible with the forgetful map of the boundary marked points.*
4. *At* $\{1\} \times \mathcal{M}(k_1, k_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1, \mathbf{p}_2)$ *it becomes* $\mathfrak{p}$.
5. *At* $\{0\} \times \mathcal{M}(k_1, k_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1, \mathbf{p}_2)$ *it becomes* $\mathfrak{back}$.
6. *It is compatible at the boundary component*
   (a)
   $$\mathcal{M}^{\text{main}}_{k'_1+1,\ell'_1}(\beta'_1; \mathbf{p}'_1)^{\mathfrak{c}}{}_{\text{ev}_0} \times_{\text{ev}^1_i} ([0,1] \times \mathcal{M}(k''_1, k_2; \ell''_1, \ell_2; \beta''_1, \beta_2; \mathbf{p}''_1, \mathbf{p}_2))$$
   where $\ell'_1 + \ell''_1 = \ell_1$, $k'_1 + k''_1 = k_1 + 1$, $\beta'_1 + \beta''_1 = \beta_1$, $(\mathbf{p}'_1, \mathbf{p}''_1) = \operatorname{Split}(\mathbf{p}_1, (\text{\L}_1, \text{\L}_2))$, $i = 1, \dots, k''_1$. *We put* $\mathfrak{c}$ *perturbation on the first factor, and*
   (b)
   $$([0,1] \times \mathcal{M}(k''_1, k_2; \ell''_1, \ell_2; \beta''_1, \beta_2; \mathbf{p}''_1, \mathbf{p}_2))_{\text{ev}_0} \times_{\text{ev}^1_i} \mathcal{M}^{\text{main}}_{k'_1+1,\ell'_1}(\beta'_1; \mathbf{p}'_1)^{\mathfrak{c}}$$
   where $\ell'_1 + \ell''_1 = \ell_1$, $k'_1 + k''_1 = k_1 + 1$, $\beta'_1 + \beta''_1 = \beta_1$, $(\mathbf{p}'_1, \mathbf{p}''_1) = \operatorname{Split}(\mathbf{p}_1, (\text{\L}_1, \text{\L}_2))$, $i = 1, \dots, k''_1$. *We put* $\mathfrak{c}$ *perturbation on the second factor.*
7. *It is compatible at the boundary component*
   (a)
   $$\mathcal{M}^{\text{main}}_{k'_2+1,\ell'_2}(\beta'_2; \mathbf{p}'_2)^{\mathfrak{c}}{}_{\text{ev}_0} \times_{\text{ev}^2_i} ([0,1] \times \mathcal{M}(k_1, k''_2; \ell_1, \ell''_2; \beta_1, \beta''_2; \mathbf{p}_1, \mathbf{p}''_2))$$
   where $\ell'_2 + \ell''_2 = \ell_2$, $k'_2 + k''_2 = k_2 + 1$, $\beta'_2 + \beta''_2 = \beta_2$, $(\mathbf{p}'_2, \mathbf{p}''_2) = \operatorname{Split}(\mathbf{p}_2, (\text{\L}_1, \text{\L}_2))$, $i = 1, \dots, k''_2$. *We put* $\mathfrak{c}$ *perturbation on the first factor, and*
   (b)
   $$([0,1] \times \mathcal{M}(k_1, k''_2; \ell_1, \ell''_2; \beta_1, \beta''_2; \mathbf{p}_1, \mathbf{p}''_2))_{\text{ev}_0} \times_{\text{ev}^2_i} \mathcal{M}^{\text{main}}_{k'_2+1,\ell'_2}(\beta'_2; \mathbf{p}'_2)^{\mathfrak{c}}$$
   where $\ell'_2 + \ell''_2 = \ell_2$, $k'_2 + k''_2 = k_2 + 1$, $\beta'_2 + \beta''_2 = \beta_2$, $(\mathbf{p}'_2, \mathbf{p}''_2) = \operatorname{Split}(\mathbf{p}_2, (\text{\L}_1, \text{\L}_2))$, $i = 1, \dots, k''_2$. *We put* $\mathfrak{c}$ *perturbation on the second factor.*

Proof. The proof is the same as other similar results we already discussed many times. □

We use Lemma 20.15 to prove the next lemma.

Lemma 20.16.

$$\begin{aligned}
&\sum_{\beta_1,\beta_2} \sum_{\ell_1+\ell_2=\ell} \sum_{k_1,k_2} \frac{T^{(\beta_1+\beta_2)\cap\omega/2\pi}}{\ell_1!\ell_2!} \exp((\beta_1+\beta_2)\cap\mathfrak{b}_2)\rho(\partial(\beta_1+\beta_2)) \\
&\qquad\qquad \int_{(\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak{b}^{\otimes\ell_1}_{\text{high}},\mathfrak{b}^{\otimes\ell_2}_{\text{high}})^{\mathfrak{p}}} (\text{ev}^1,\text{ev}^2)^*([\mathfrak{v};k_1] \times [\mathfrak{w};k_2]) \\
= &\sum_{\beta_1,\beta_2} \sum_{\ell_1+\ell_2=\ell} \sum_{k_1,k_2} \frac{T^{(\beta_1+\beta_2)\cap\omega/2\pi}}{\ell_1!\ell_2!} \exp((\beta_1+\beta_2)\cap\mathfrak{b}_2)\rho(\partial(\beta_1+\beta_2)) \\
&\qquad\qquad \int_{(\mathcal{M}(k_1,k_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak{b}^{\otimes\ell_1}_{\text{high}},\mathfrak{b}^{\otimes\ell_2}_{\text{high}})^{\mathfrak{back}}} (\text{ev}^1,\text{ev}^2)^*([\mathfrak{v};k_1] \times [\mathfrak{w};k_2]).
\end{aligned}$$

Proof. We put

$$\mathfrak{b}_{\rm high}^{\otimes \ell} = \sum_{I\in\mathfrak{I}_\ell} c_I \mathbf{f}_{i_1}\otimes\cdots\otimes\mathbf{f}_{i_\ell} = \sum_{I\in\mathfrak{I}_\ell} c_{I,\ell}\mathbf{f}_{I,\ell}. \tag{20.10}$$

We apply Stokes' theorem (Lemma 12.18 [FOOO5]) on

$$([0,1]\times M(k_1+1,k_2+1;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{f}_{I,\ell_1},\mathbf{f}_{I,\ell_2}))^{\mathfrak{s}} \tag{20.11}$$

to the pull back $(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2])$, which is a closed form.

Let us study of the integration of $(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1]\times[\mathfrak{w};k_2])$ on each component of the boundary of (20.11).

The case (4) and (5) of Lemma 20.15 give the left and right hand sides of the formula in Lemma 20.16, respectively after taking weighted sum with weight

$$\frac{T^{\beta\cap\omega/2\pi}}{\ell_1!\ell_2!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta). \tag{20.12}$$

Let us study (6) of Lemma 20.15. We remark that

$$[\mathfrak{v};k_1] = [(b_+^{\mathfrak{c}})^{k_1}\times\mathfrak{v}].$$

We integrate it on the boundary component described by Lemma 20.15 (6). We divide it into two cases.

(Case 1) Lemma 20.15 (6) (a) This is the case when the boundary marked point corresponding to $\mathfrak{v}$ is on the second factor.

(Case 2) Lemma 20.15 (6) (b). This is the case when the boundary marked point corresponding to $\mathfrak{v}$ is on the first factor.

We can show that contribution of both cases vanish in the same way as in the proof of Theorem 19.8. Namely we can study (Case 1) in the same way as (3) Lemma 19.10 and (Case 2) in the same way as (4) Lemma 19.10 . We repeat the detail below for completeness' sake.

We first study Case 1. We define E by (19.11). Then the contribution of Case 1 is

$$\sum_{\substack{k_1,k_2,\\ \ell_1,\ell_2}}\sum_{\substack{I_1\in\mathfrak{I}_{\ell_1}\\ I_2\in\mathfrak{I}_{\ell_2}}}\sum_{i=1}^{k_1+1}\sum_{\beta_1,\beta_2} T^{(\beta_1+\beta_2)\cap\omega/2\pi}\exp((\beta_1+\beta_2)\cap\mathfrak{b}_2)\frac{c_{I_1}c_{I_2}}{\ell_1!\ell_2!}\rho(\partial(\beta_1+\beta_2))$$
$$\int_{([0,1]\times\mathcal{M}(k_1,k_2;\ell_1;\ell_2;\beta_1;\beta_2;\mathbf{p}_1,\mathbf{p}_2))^{\mathfrak{s}}}(\mathrm{ev}_1,\mathrm{ev}_2)^*([(b_+^{\mathfrak{c}})^{i-1}\times E\times(b_+^{\mathfrak{c}})^{k_1+1-i}\times\mathfrak{v}],[\mathfrak{w};k_2]).$$

In the same way as the proof of Lemma 19.7, we can prove that E is proportional to the unit. Therefore this contribution vanishes by Lemma 20.15 (3).

We next study Case 2. We define R by

$$\begin{aligned} R = \sum_{\substack{k_1,k_2,\\ \ell_1,\ell_2}}\sum_{\substack{I_1\in\mathfrak{I}_{\ell_1}\\ I_2\in\mathfrak{I}_{\ell_2}}}\sum_{\beta_1,\beta_2} & T^{\beta\cap\omega/2\pi}\exp(\beta\cap\mathfrak{b}_2)\frac{c_I}{\ell!}\rho(\partial\beta) \\ & (\mathrm{ev}_0)_*\big(((\mathrm{ev}_1,\mathrm{ev}_2,\dots,\mathrm{ev}_{k_1}),\mathrm{ev}_2)^*((b_+^{\mathfrak{c}})^{k_1},[\mathfrak{w};k_2]); \\ & \qquad ([0,1]\times\mathcal{M}(k_1+1,k_2;\ell_1;\ell_2;\beta_1;\beta_2;\mathbf{p}_1,\mathbf{p}_2))^{\mathfrak{s}}\big). \end{aligned} \tag{20.13}$$

By definition the contribution of Case 2 is

$$\langle\delta^{\mathfrak{b},b^{\mathfrak{c}}}(R),\mathfrak{v}\rangle = 0.$$

We can show that the contribution of (7) of Lemma 20.15 vanishes in the same way. The proof of Lemma 20.16 is now complete. □

Proposition 20.8 follows from Formulae (20.8), (20.9) and Lemma 20.16. □

Before proceeding to the proofs of Propositions 20.9 and 20.10, we fix the choice of perturbation on $\mathfrak{forget}^{-1}([\Sigma_2]) \cap \mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta; \mathfrak{b}^{\otimes\ell}_{\text{high}})$.

Let $(\mathbf{p}_1, \mathbf{p}_2) = \text{Split}(\mathbf{p}, (\text{Ł}_1, \text{Ł}_2))$ and $\ell_j = |\mathbf{p}_j|$, $j = 1, 2$.

We consider $\mathcal{M}^{\text{main}}_{k_j+2;\ell_j}(\beta_j; \mathbf{p}_j)$ and take perturbation $\mathfrak{c}$. Let $a_j \in \{1, \dots, k_j + 1\}$. We consider

$$(\text{ev}_0, \text{ev}_{a_1}) : \mathcal{M}^{\text{main}}_{k_1+2;\ell_1}(\beta_1; \mathbf{p}_1)^{\mathfrak{c}} \to L(u) \times L(u) \tag{20.14}$$

and

$$(\text{ev}_{a_2}, \text{ev}_0) : \mathcal{M}^{\text{main}}_{k_2+2;\ell_2}(\beta_2; \mathbf{p}_2)^{\mathfrak{c}} \to L(u) \times L(u). \tag{20.15}$$

Lemma 20.17. *The two maps (20.14) and (20.15) are transversal.*

Proof.

$$(\text{ev}_0)_* \oplus 0 : T\mathcal{M}^{\text{main}}_{k_1+2;\ell_1}(\beta_1; \mathbf{p}_1)^{\mathfrak{c}} \to TL(u) \oplus \{0\}$$

and

$$0 \oplus (\text{ev}_0)_* : T\mathcal{M}^{\text{main}}_{k_2+2;\ell_2}(\beta_2; \mathbf{p}_2)^{\mathfrak{c}} \to \{0\} \oplus TL(u)$$

are both surjective, by T^n equivariance. The lemma follows immediately. □

Remark 20.18. We are studying the moduli space of holomorphic annuli (that is the case of one loop). The proof of the above lemma works only in the one loop case. Namely if we consider the moduli space of pseudo-holomorphic curves from bordered Riemann surface which is more complicated than annuli, the proof of Lemma 20.17 does not work to prove the transversality of the configuration of the discs which appear at the end of its compactification. For example, let us consider the fiber product

$$\mathcal{M}^{\text{main}}_3(\beta_1) \,{}_{(\text{ev}_0,\text{ev}_1,\text{ev}_2)}\times_{(\text{ev}_0,\text{ev}_1,\text{ev}_2)} \mathcal{M}^{\text{main}}_3(\beta_2). \tag{20.16}$$

This fiber product appears at the end of the moduli space of pseudo-holomorphic maps from the Riemann surface which is D^2 minus disjoint union of two D^2's. The perturbation ($\mathfrak{c}$ or $\mathfrak{q}$) is designed so that the evaluation map at one boundary marked point is a submersion. (We may take *any* of the marked points.) This is enough for the transversality appearing in Lemma 20.17 but is *not* enough for the fiber product (20.16) to be transversal automatically after this perturbation is applied to each of the factors.

Definition 20.19. We denote by $\mathcal{N}(k_1, k_2; a_1, a_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1; \mathbf{p}_2)$ the fiber product of the maps (20.14) and (20.15).

There is an obvious map

$$\begin{aligned}\text{Glue} : \mathcal{N}&(k_1, k_2; a_1, a_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1; \mathbf{p}_2) \\ &\to \mathcal{M}^{\text{ann;main}}_{(k_2+a_1-a_2,k_1-a_1+a_2);\ell}(\beta_1 + \beta_2; \mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2]).\end{aligned} \tag{20.17}$$

Lemma 20.20. *The map* Glue *defines an isomorphic to the image. The moduli space* $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta; \mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2])$ *is decomposed to the images of* Glue*'s from difference sources. If* $\mathbf{x} \in \mathcal{M}^{\text{ann;main}}_{(k,k');\ell}(\beta; \mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2])$ *is an image of a codimension* N *corner point of* $\mathcal{N}(k_1, k_2; a_1, a_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1; \mathbf{p}_2)$*, then it is an image of* N *points from the union of the sources of* Glue.

Kuranishi structures of the sources are compatible at the overlapped part of the target.

Furthermore, the multisections $\mathfrak{c}$ are compatible also.

Proof. In fact, the proof is the same as Lemmata 9.20, 9.21. We however emphasis that cyclic symmetry is crucial for this lemma to hold. To elaborate this point, let us consider the subset consisting of elements $((D^2,u_1),(D^2,u_2),(D^2,u_3))$ of $\mathcal{M}_3^{\rm main}(\beta_1)\times\mathcal{M}_2^{\rm main}(\beta_2)\times\mathcal{M}_3^{\rm main}(\beta_3)$ such that

$$\mathrm{ev}_1(u_1)=\mathrm{ev}_1(u_2),\quad \mathrm{ev}_2(u_1)=\mathrm{ev}_2(u_3),\quad \mathrm{ev}_0(u_2)=\mathrm{ev}_1(u_3). \tag{20.18}$$



$\mathrm{ev}_1(u_1)=\mathrm{ev}_1(u_2)$
$u_1(D^2)$
$u_2(D^2)$
ev_0
$\mathrm{ev}_2(u_1)=\mathrm{ev}_2(u_3)$
$u_3(D^2)$
$\mathrm{ev}_2(u_2)=\mathrm{ev}_1(u_3)$
ev_0

Figure 20.1

This set is a subset of $\mathcal{M}_{(1,1);0}^{\rm ann;main}(\beta_1+\beta_2+\beta_3)\cap\mathfrak{forget}^{-1}(\Sigma_2)$ and lies in its codimension 2 strata.

We consider 2 (among 3) ways to resolve singularities. Namely at $\mathrm{ev}_1(u_1)=\mathrm{ev}_1(u_2)$ or at $\mathrm{ev}_0(u_2)=\mathrm{ev}_1(u_3)$.

We remark that for $((D^2,u_1),\ (D^2,u_3))$ the choice of the 0-th marked point (among the 3 boundary marked points) is canonical. Namely there is a marked point which does not meet to other irreducible component. We take it as the 0-th one.

On the other hand, there is no good way to decide which (among 2) boundary marked point is the 0-th one for (D^2,u_2).

In fact, if we insist the consistency with the process to resolve $\mathrm{ev}_1(u_1)=\mathrm{ev}_1(u_2)$, the marked point of (D^2,u_2) which intersects with (D^2,u_1) must be 0-th one. If we insist the consistency with the process to resolve $\mathrm{ev}_0(u_2)=\mathrm{ev}_1(u_3)$, the marked point of (D^2,u_2) which intersects with (D^2,u_3) must be 0-th one.

Thus for the consistency of both, we need to take our perturbation of $\mathcal{M}_2^{\rm main}(\beta_2)$ to be invariant under cyclic symmetry. Actually this is one of the reasons we take the $\mathfrak{c}$ perturbation. (See however [Ab], where M. Abouzaid uses less symmetry than ours. We use cyclic symmetry also in Sections 19 and 22. See Remarks 19.12 and 22.5.)

Once this point is understood, the proof is the same as Lemmata 9.20, 9.21. □

By Lemma 20.20 we can define a system of continuous families of multisections $\mathfrak{s}$ on $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2])$ so that it coincides with the fiber product of $\mathfrak{c}$ multisection on each factor of $\mathcal{N}(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2)$.

We thus fixed our family of multisections for $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2])$. It has the following properties.

Condition 20.21.
1. It is transversal to 0 and is T^n-invariant.
2. It is invariant under the cyclic permutations of boundary marked points and arbitrary permutations of the interior marked points.
3. It is compatible with the forgetful map of the boundary marked points.
4. It is transversal to the map (20.17) in a sense similar to Lemma 9.15 (5).
5. It is compatible at the boundary components.

The boundary components of various irreducible components of $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2])$ are divided in 6 cases as in Figure 20.2 below.



Case 1



Case 2



Case 3



Case 4



Case 5 Case 6

Figure 20.2

Proof of Proposition 20.9. We extend the above multisection to a neighborhood of $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma_2])$ in $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})$. Then for each $[\Sigma'] \in \mathcal{M}^{\text{ann;main}}_{(1,1);0}$ sufficiently close to $[\Sigma_2]$ we obtain a multisection on $\mathcal{M}^{\text{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma'])$ which satisfies the same conditions as Condition 20.21 above. We choose a sequence $[\Sigma_i'] \in \mathcal{M}^{\text{ann;main}}_{(1,1);0}$ which is in stratum (1) in Lemma 20.6 and

$$\lim_{i\to\infty}[\Sigma_i'] = [\Sigma_2].$$

Lemma 20.22.

$$\begin{aligned}&\lim_{i\to\infty}\int_{\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma_i'])}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2])\\ &=\int_{\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma_2])}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2]).\end{aligned}$$

Proof. This lemma is slightly nontrivial since $\mathfrak{forget}$ is not smooth at the fiber of $[\Sigma_2]$. Let us decompose $\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma_2]))$ to the union of $\mathcal{N}(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2)$ by Lemma 20.20. We take an increasing sequence of compact subsets

$$\mathcal{N}_K(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2),\quad K=1,\dots,\to\infty$$

of the union $\mathrm{Int}\,\mathcal{N}(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2)$ of the interior of its irreducible components such that

$$\begin{aligned}&\bigcup_K \mathcal{N}_l(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2)\\ &=\mathrm{Int}\,\mathcal{N}(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2).\end{aligned}$$

We consider the irreducible components of dimension $\deg[\mathfrak{v};k_1]+\deg[w;k_2]$ only. For each K we can construct a map

$$\begin{aligned}\Phi_{K,i}:&\mathcal{N}(k_1+1,k_2+1;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathbf{p}_1;\mathbf{p}_2)\\ &\to\mathcal{M}^{\text{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma_i'])\end{aligned}$$

such that

$$\lim_{i\to\infty} \Phi^*_{K,i}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2]) = (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2]).$$

By transversality and definition, the contribution to the integration of the right hand side of Lemma 20.22 of the codimension one stratum is 0. Therefore the integration of $(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2])$ on the union of the images of $\Phi_{K,i}$ can be chosen arbitrary close to the right hand side if we take K, i large.

On the other hand, the complement of the union of the images of $\Phi_{K,i}$ are obtained by gluing at least three pieces of discs. (Those configurations are described as in Figure 20.3.) So there are at least 3 parameters T_1, T_2, T_3 for gluing. When we restrict these 3 parameters by requiring that the resulting source becomes the given Σ'_i, we still have one more parameter T. We can estimate the differential of $(\mathrm{ev}^1,\mathrm{ev}^2)$ with respect to T in the same way as Lemma A1.58 [FOOO2] and find that it is of order e^{-cT}. Therefore the integration of $(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2])$ on the complement of the union of images of $\Phi_{l,i}$ goes to zero as $l,i\to\infty$. Hence the lemma.



Figure 20.3

□

We next choose a sequence $[\Sigma''_i] \in \mathcal{M}^{\mathrm{ann;main}}_{(1,1);0}$ which is in stratum (1) in Lemma 20.6 and

$$\lim_{i\to\infty}[\Sigma''_i] = [\Sigma_1].$$

We take a perturbation of $\mathcal{M}^{\mathrm{ann;main}}_{(1,1);\ell}(\beta;\mathbf{p}) \cap \mathfrak{forget}^{-1}([\Sigma''_i])$ by restricting the perturbation given in Lemma 20.14.

Lemma 20.23.

$$\begin{aligned}&\lim_{i\to\infty}\int_{\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma''_i])}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2])\\ &=\int_{\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathbf{p})\cap\mathfrak{forget}^{-1}([\Sigma_1])}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2]).\end{aligned}$$

The proof is the same as the proof of Lemma 20.22.

Lemma 20.24. *For any i we have*

$$\begin{aligned}&\sum_{\beta}\sum_{\ell}\sum_{k_1,k_2}\frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta)\\ &\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma'_i])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2])\\ &=\sum_{\beta}\sum_{\ell}\sum_{k_1,k_2}\frac{T^{\beta\cap\omega/2\pi}}{\ell!}\exp(\beta\cap\mathfrak{b}_2)\rho(\partial\beta)\\ &\qquad\qquad\int_{\mathfrak{forget}^{-1}([\Sigma''_i])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k_2+1);\ell}(\beta;\mathfrak{b}_{\mathrm{high}}^{\otimes\ell})}(\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak{v};k_1],[\mathfrak{w};k_2]).\end{aligned}$$

Proof. We take a path $\gamma:[0,1]\to\mathcal{M}^{\mathrm{ann;main}}_{(1,1);0}$ joining $[\Sigma'_i]$ with $[\Sigma''_i]$ in the stratum (1) in Lemma 20.6. We take the fiber product

$$[0,1]_{\gamma}\times_{\mathfrak{forget}}\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})$$

and denote it by $\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p};\gamma)$. This space has a Kuranishi structure and its boundary is described as follows:

(1) $\mathfrak{forget}^{-1}([\Sigma'_i])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})$.

(2) $\mathfrak{forget}^{-1}([\Sigma''_i])\cap\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p})$.

(3)
$$\mathcal{M}_{k'_1+1;\ell'}(\beta';\mathbf{p}_1)^{\mathfrak{c}}\,{}_{\mathrm{ev}_0}\times_{\mathrm{ev}^1_i}\mathcal{M}^{\mathrm{ann;main}}_{(k''_1+1,k_2+1);\ell''}(\beta'';\mathbf{p}_2;\gamma).$$
Here $k'_1+k''_1=k_1+1$, $\ell'+\ell''=\ell$, $\beta'+\beta''=\beta$, $i=1,\dots,k''_1$.

(4)
$$\mathcal{M}_{k'_2+1;\ell'}(\beta';\mathbf{p}_1)^{\mathfrak{c}}\,{}_{\mathrm{ev}_0}\times_{\mathrm{ev}^2_i}\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k''_2+1);\ell''}(\beta'';\mathbf{p}_2;\gamma).$$
Here $k'_2+k''_2=k_2+1$, $\ell'+\ell''=\ell$, $\beta'+\beta''=\beta$, $i=1,\dots,k''_2$.

(5)
$$\mathcal{M}^{\mathrm{ann;main}}_{(k'_1+1,k_2+1);\ell'}(\beta';\mathbf{p}_1)\,{}_{\mathrm{ev}^1_0}\times_{\mathrm{ev}_i}\mathcal{M}_{k''_1+1;\ell''}(\beta';\mathbf{p}_2;\gamma')^{\mathfrak{c}}.$$
Here $k'_1+k''_1=k_1+1$, $\ell'+\ell''=\ell$, $\beta'+\beta''=\beta$, $i=1,\dots,k''_1$.

(6)
$$\mathcal{M}^{\mathrm{ann;main}}_{(k_1+1,k'_2+1);\ell'}(\beta';\mathbf{p}_1)\,{}_{\mathrm{ev}^2_0}\times_{\mathrm{ev}_i}\mathcal{M}_{k''_2+1;\ell''}(\beta'';\mathbf{p}_2;\gamma)^{\mathfrak{c}}.$$
Here $k'_2+k''_2=k_1+1$, $\ell'+\ell''=\ell$, $\beta'+\beta''=\beta$, $i=1,\dots,k''_2$.

We define a multisection $\mathfrak{s}$ on $\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta;\mathbf{p};\gamma)$ so that it is compatible at the above boundaries. We also assume $\mathfrak{s}$ is transversal to 0 and T^n-invariant. Moreover it is symmetric with respect to the cyclic permutation of boundary marked points at each boundary component and permutation of interior marked points.

Now we put

$$\mathfrak{b}_{\mathrm{high}}^{\otimes \ell} = \sum_{I \in \mathfrak{I}_\ell} c_I \mathbf{f}_{i_1} \otimes \cdots \otimes \mathbf{f}_{i_\ell} = \sum_{I \in \mathfrak{I}_\ell} c_{I,\ell} \mathbf{f}_{I,\ell}$$

and apply Stokes's theorem for $(\mathrm{ev}^1, \mathrm{ev}^2)^*([\mathfrak{v}; k_1], [\mathfrak{w}; k_2])$ on $\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta; \mathbf{f}_{I,\ell}; \gamma)^{\mathfrak{s}}$. The rest of the proof is the same as the proof of Lemma 20.16. ☐

Proposition 20.9 follows from Lemmata 20.22, 20.23, and 20.24. ☐

Proof of Proposition 20.10. We begin with the following lemma. Let $g \in T^n$. We let it act on $L(u) \times L(u)$ by $g(x, y) = (gx, y)$. Let dg be the Haar measure of T^n and let T_Δ be an distributional n form on $L(u) \times L(u)$ which is Poincaré dual to the diagonal. We regard $\mathbf{e}_I$ (the generator of $H(L(u); \mathbb{C})$) as a harmonic form on $L(u)$. Let g^{IJ} be as in Theorem 20.1.

Lemma 20.25.

$$\int_{g \in T^n} g_* T_\Delta dg = \sum_{I,J} (-1)^{|I||J|} g^{IJ} \mathbf{e}_I \times \mathbf{e}_J.$$

See Lemma 26.7 for the sign. (Precisely speaking, in Lemma 26.7 we work on chains instead of cochains.) Here the equality is as distributional n forms on $L(u) \times L(u)$.

Proof. The left hand side is a $T^n \times T^n$ equivariant n form. Therefore it is smooth and harmonic. It is also cohomologous to the Poincaré dual to the diagonal. The lemma follows. ☐

Let $g_1, g_2 \in T^n$. We define

$$(g_1 \mathrm{ev}_0, g_2 \mathrm{ev}_{a_1}) : \mathcal{M}_{k_1+2;\ell_1}(\beta_1; \mathbf{p}_1) \to L(u) \times L(u) \tag{20.19}$$

by

$$(g_1 \mathrm{ev}_0, g_2 \mathrm{ev}_{a_1})(\mathbf{x}) = (g_1 \cdot \mathrm{ev}_0(\mathbf{x}), g_2 \cdot \mathrm{ev}_{a_1}(\mathbf{x})).$$

Lemma 20.26. *The two maps* (*20.19*) *and* (*20.15*) *are transversal.*

The proof is the same as Lemma 20.17. We denote the fiber product of (20.19) and (20.15) by $\mathcal{N}(k_1, k_2; a_1, a_2; \ell_1, \ell_2; \beta_1, \beta_2; \mathbf{p}_1; \mathbf{p}_2; g_1, g_2)$.

Lemma 20.27.

$$\begin{aligned} &\frac{1}{\ell!} \int_{\mathcal{M}^{\mathrm{ann;main}}_{(k_1,k_2);\ell}(\beta; \mathfrak{b}_{\mathrm{high}}^{\otimes \ell}) \cap \mathfrak{forget}^{-1}([\Sigma_2])} (\mathrm{ev}^1, \mathrm{ev}^2)^*([\mathfrak{v}; k_1], [\mathfrak{w}; k_2]) \\ &= \sum_{\substack{\beta_1+\beta_2=\beta \\ \ell_1+\ell_2=\ell}} \sum_{\substack{a_1=1,\dots,k_1 \\ a_2=1,\dots,k_2}} \frac{1}{\ell_1! \ell_2!} \int_{\mathcal{N}(k_1,k_2:a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak{b}_{\mathrm{high}}^{\otimes \ell_1};\mathfrak{b}_{\mathrm{high}}^{\otimes \ell_2};e,e)} (\mathrm{ev}^1, \mathrm{ev}^2)^*([\mathfrak{v}; k_1], [\mathfrak{w}; k_2]). \end{aligned}$$

Here e denotes the unit of the group T^n.

Proof. This is immediate from Lemma 20.20 and the definition. ☐

Lemma 20.28.

$$\begin{aligned}
&\sum_{\substack{\beta_1+\beta_2=\beta\\ \ell_1+\ell_2=\ell}} \sum_{\substack{a_1=1,\dots,k_1\\ a_2=1,\dots,k_2}} \frac{T^{(\beta_1+\beta_2)\cap\omega}}{\ell_1!\ell_2!} \exp((\beta_1+\beta_2)\cap\mathfrak b_2)\rho(\partial\beta)\\
&\int_{g\in T^n} dg \int_{\mathcal N(k_1,k_2:a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};g,e)} (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak v;k_1],[\mathfrak w;k_2])\\
&= \sum_{\beta_1,\beta_2}\sum_{\ell_1+\ell_2}\sum_{I,J\in 2^{\{1,\dots,n\}}} g^{IJ}\frac{T^{(\beta_1+\beta_2)\cap\omega}}{\ell_1!\ell_2!}\exp((\beta_1+\beta_2)\cap\mathfrak b_2)\rho(\partial(\beta_1+\beta_2))\\
&\qquad\qquad \langle \mathfrak q^{\mathfrak c,b_+^{\mathfrak c}}_{2,\ell_1;\beta_1}(\mathfrak b_{\rm high}^{\otimes\ell_1};(\mathfrak v\otimes\mathbf e_I)),\mathfrak q^{\mathfrak c,b_+^{\mathfrak c}}_{2,\ell_2;\beta_2}(\mathfrak b_{\rm high}^{\otimes\ell_2};(\mathbf e_J\otimes\mathfrak w))\rangle_{\mathrm{PD}_L}.
\end{aligned}$$

Proof. This is immediate from Lemma 20.25 and the definition. □

Lemma 20.29.

$$\begin{aligned}
&\sum_{\substack{\beta_1+\beta_2=\beta\\ \ell_1+\ell_2=\ell}} \sum_{\substack{a_1=1,\dots,k_1\\ a_2=1,\dots,k_2}} \frac{T^{(\beta_1+\beta_2)\cap\omega}}{\ell_1!\ell_2!} \exp((\beta_1+\beta_2)\cap\mathfrak b_2)\rho(\partial\beta)\\
&\qquad\qquad \int_{\mathcal N(k_1,k_2:a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};g,e)} (\mathrm{ev}^1,\mathrm{ev}^2)^*([\mathfrak v;k_1],[\mathfrak w;k_2])
\end{aligned}$$

is independent of $g\in T^n$.

Proof. Let $g,g'\in T^n$ and $\gamma:[0,1]\to T^n$ be a path joining them. We put

$$\begin{aligned}
&\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};\gamma,e)\\
&= \bigcup_{t\in[0,1]}\{t\}\times\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};\gamma(t),e).
\end{aligned}$$

Note by Lemma 20.26 the $\mathfrak c$-perturbation is transversal for *any*

$$\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};\gamma(t),e).$$

So we have already chosen the perturbation of the moduli space

$$\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};\gamma,e).$$

The boundary of $\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};\gamma,e)$ is

$$\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};g,e)$$

and

$$\mathcal N(k_1,k_2;a_1,a_2;\ell_1,\ell_2;\beta_1,\beta_2;\mathfrak b_{\rm high}^{\otimes\ell_1};\mathfrak b_{\rm high}^{\otimes\ell_2};g',e),$$

and those described by the following Figure 20.4.



Case 1



Case 2



Case 3



Case 4



Figure 20.4

We apply Stokes' theorem to $(\mathrm{ev}^1, \mathrm{ev}^2)^*([\mathfrak{v}; k_1], [\mathfrak{w}; k_2])$. It suffices to show that the contribution from the boundaries of Figure 20.4 becomes zero after taking the weighted sum.

The boundary Case 1 cancels with those of Case 5, after taking sum over β_1, β_2.

Similarly the boundary Case 2 cancels with those of Case 6, after taking sum over β_1, β_2. (We use the T^n-equivariance of the moduli space here.)

The boundaries of Cases 4 and 8 are zero. This is by the same reason as the (3) of Lemma 19.10. (Namely unitality of the element E there and compatibility with the forgetful map of our perturbations.) (Note that there are some other cases where bubble with no boundary marked points occurs at the different position. They all can be handled in the same way.)

Finally the contribution from the boundary components of the Cases 3 and 7 becomes zero after taking weighted sum. This is by the same reason as the Case

(4) of Lemma 19.10. (The vanishing of the boundary operator.) Lemma 20.29 is proved. □

Lemmata 20.27, 20.28, 20.29 imply Proposition 20.10. □

The proof of Theorem 20.1 is now complete. □

Remark 20.30. Theorem 20.1 is stated and proved only for the case of fibers of toric manifolds. However various parts of the proof (especially the idea to use the moduli space of pseudo-holomorphic annuli and cyclic symmetry) can be applied to more general situation.

We choose to prove this statement only for the toric case in this paper since in toric case we can take short cut in several places, while the proof for the general case is rather involved. For example, we always use T^n-invariant perturbations in the present paper and so do not need to use general differential forms on $L(u) = T^n$ other than the T^n-equivariant ones. In the general situation we need to use a canonical model and so the moduli spaces involved become more complicated to define. Also we use the technique of averaging differential forms over the T^n action in our proof of Proposition 20.10. Since this technique is not available for other cases, we need to use more sophisticated homological algebra instead in the non-toric cases.

Another reason why we restrict ourselves to the toric case here is that at the time of writing this article we do not have any other particular examples to which we may apply the ideas used in this paper. However we have little doubt that many non-toric examples, to which many ideas presented here can be applied, will be found in the near future.

21. Poincaré duality is residue pairing

We take

$$\mathrm{vol}_{L(u)} \in H^n(L(u);\Lambda_0) \cong HF((L(u),\mathfrak{b},b^{\mathfrak{c}}),(L(u),\mathfrak{b},b^{\mathfrak{c}});\Lambda_0),$$

which is the volume form of $L(u)$. We normalize it so that $\int_{L(u)} \mathrm{vol}_{L(u)} = 1$.

Remark 21.1. It may be confusing that $\mathrm{vol}_{L(u)} \neq 1$. We note that 1 is the Poincaré dual to the fundamental *homology* class $[L(u)]$. The degree of $\mathrm{vol}_{L(u)}$ is n and the degree of 1 is 0.

The results of Sections 19, 20 imply:

Proposition 21.2. *Let* $(u,b) \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$. *Then we have*

$$\langle i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(\mathrm{vol}_{L(u)}), i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(\mathrm{vol}_{L(u)})\rangle_{\mathrm{PD}_X} = Z(\mathfrak{b},b).$$

Remark 21.3. We still do not assume that (u,b) is nondegenerate. When (u,b) is a degenerate critical point, Proposition 21.2 implies that the left hand side is 0.

Proof. This is immediate from Theorem 20.1 and Corollary 19.9. See Subsection 26.2 for the proof of the sign in the above formula. □

Now we are ready to complete the proof of Theorem 2.32 (2). We assume that $\mathfrak{PO}_{\mathfrak{b}}$ is a Morse function. We first consider two elements (u,b), $(u',b') \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$, $(u,b) \neq (u',b')$ corresponding to the unit elements of some factors

$\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}};\eta)$ in the factorization of $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ given in Proposition 2.15. We denote the corresponding unit elements of $\mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}})$ by $1_{(u,b)}$, $1_{(u',b')}$ respectively. Then

$$\langle 1_{(u,b)}, 1_{(u',b')}\rangle_{\mathrm{PD}_X} = \langle 1_{(u,b)}, 1_{(u',b')} \cup^{\mathfrak{b}} 1\rangle_{\mathrm{PD}_X} = \langle 1_{(u,b)} \cup^{\mathfrak{b}} 1_{(u',b')}, 1\rangle_{\mathrm{PD}_X} = 0.$$

Here $\cup^{\mathfrak{b}}$ is the quantum cup product deformed by $\mathfrak{b}$. (See Section 2.) (The second equality is nothing but the well known Frobenius pairing in Gromov-Witten theory. See for example [Ma].) So it suffices to study the case $(u,b) = (u',b')$. We identify $1_{(u,b)} \in \mathrm{Jac}(\mathfrak{PO}_{\mathfrak{b}}) \otimes_{\Lambda_0} \Lambda$ with an element of $H(X;\Lambda)$ by the isomorphism Theorem 2.32 (1). Since $1_{(u,b)}$ is the idempotent and $\mathfrak{ks}_{\mathfrak{b}}$ is a ring isomorphism it follows from Lemma 17.1 that

$$i^{\#}_{\mathrm{qm},(u,b)}(1_{(u,b)}) = 1 \in H^0(L(u);\Lambda).$$

Therefore Theorem 19.8 and $\langle 1, \mathrm{vol}_{L(u)}\rangle_{\mathrm{PD}_{L(u)}} = 1$ imply

$$\langle 1_{(u,b)}, i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(\mathrm{vol}_{L(u)})\rangle_{\mathrm{PD}_X} = 1. \tag{21.1}$$

Since

$$i^{\#}_{\mathrm{qm},(u,b)}(1_{(u',b')}) = 0$$

for $(u',b') \neq (u,b)$, it follows from (21.1) that

$$i_{\#,\mathrm{qm},(\mathfrak{b},b,u)}(\mathrm{vol}_{L(u)}) = \frac{1}{\langle 1_{(u,b)}, 1_{(u,b)}\rangle_{\mathrm{PD}_X}} 1_{(u,b)}. \tag{21.2}$$

Therefore Proposition 21.2 implies

$$\langle 1_{(u,b)}, 1_{(u,b)}\rangle_{\mathrm{PD}_X} Z(\mathfrak{b}, b) = 1.$$

The proof of Theorem 2.32 (2) is now complete. □

22. Clifford algebra and Hessian matrix

In Sections 22 and 23, we will prove Theorem 2.24.

In [Cho2], Cho proved that Floer cohomology of the toric fiber is isomorphic to the Clifford algebra in the Fano case as a ring (for the case $b = 0$, $\mathfrak{b} = 0$). In this section, we follow Cho's argument to prove the following Theorem 22.2, from which (2),(3) of Theorem 2.24 follows.

Let $X_1, \dots, X_n$ be formal variables and $d_i \in \Lambda$ ($i = 1, \dots, n$). We consider relations

$$\begin{cases} X_iX_j + X_jX_i = 0, & i \neq j \\ X_iX_i = d_i 1. & \end{cases} \tag{22.1}$$

We take a free (noncommutative) Λ algebra generated by $X_1, \dots, X_n$ and divide it by the two-sided ideal generated by (22.1). We denote it by $\mathrm{Cliff}_{\Lambda}(n;\vec{d})$, where we set $\vec{d} = (d_1, \dots, d_n)$.

Let $I = (i_1, \dots, i_k)$, $1 \le i_1 < \cdots < i_k \le n$. We write the set of such I's by $2^{\{1,\dots,n\}}$. We put

$$X_I = X_{i_1}X_{i_2}\cdots X_{i_{k-1}}X_{i_k} \in \mathrm{Cliff}_{\Lambda}(n;\vec{d}).$$

It is well known and can be easily checked that $\{X_I \mid I \in 2^{\{1,\dots,n\}}\}$ forms a basis of $\mathrm{Cliff}_{\Lambda}(n;\vec{d})$ as a Λ vector space.

For $I \in 2^{\{1,\dots,n\}}$, we denote its complement by I^c.

We next consider $(u,b) \in \mathrm{Crit}(\mathfrak{PO}_{\mathfrak{b}})$. We take $(u, b^{\mathfrak{c}})$ which is the weak bounding cochain of the cyclic filtered A_∞ algebra $(H(L(u);\Lambda_0), \{\mathfrak{m}_k^{\mathfrak{c},\mathfrak{b}}\})$ corresponding to b.

When (2) of Condition 2.19 is satisfied, we have $b = b^{\mathfrak c}$. If (1) or (2) of Condition 2.19 is satisfied, we have $b^{\mathfrak c} \in H^1(L(u);\Lambda_0)$.

We put $y(u)_i = e^{x_i}$ and regard $\mathfrak{PO}_{\mathfrak b}$ as a function of x_i. Denote $b = \sum_{i=1}^n x_i(b)\mathbf{e}_i$.

Definition 22.1. If (2) of Condition 2.19 is satisfied, we put $\mathfrak{PO}^{\mathfrak c}_{\mathfrak b} = \mathfrak{PO}_{\mathfrak b}$. If (1) of Condition 2.19 is satisfied, we put

$$\mathfrak{PO}^{\mathfrak c}_{\mathfrak b} = \mathfrak{PO}_{\mathfrak b} \circ \mathfrak{F}_*^{-1} : H^1(L(u);\Lambda_0) \to \Lambda_0. \tag{22.2}$$

We recall that Corollary 18.26 shows $\mathfrak{F}_* =$ identity on $H^1(L(u);\Lambda_0)$ when (2) of Condition 2.19 is satisfied.

Let $(2d_1,\dots,2d_n)$ be the eigenvalues of the Hessian matrix

$$\left[\frac{\partial^2 \mathfrak{PO}^{\mathfrak c}_{\mathfrak b}}{\partial x_i \partial x_j}(x(b^{\mathfrak c}))\right]_{i,j=1}^{i,j=n}$$

counted with multiplicity.

In Sections 21 and 22, we identify

$$H(L(u),\Lambda) \cong HF((L(u);(\mathfrak b, b));(L(u);(\mathfrak b, b));\Lambda).$$

Then by definition, the product $\cup^{\mathfrak c,Q}$ is given by

$$h_1 \cup^{\mathfrak c,Q} h_2 = (-1)^{\deg h_1(\deg h_2+1)}\mathfrak m_2^{\mathfrak c,\mathfrak b,b^{\mathfrak c}}(h_1,h_2)$$

where $\mathfrak m_2^{\mathfrak c,\mathfrak b,b^{\mathfrak c}}$ is as defined in (18.6). Then the (filtered) A_∞ relation implies that $(H(L(u);\Lambda),\cup^{\mathfrak c,Q})$ defines a ring.

Theorem 22.2. *Suppose that one of the two conditions in Condition 2.19 is satisfied. There exist a basis* $\{\mathbf{e}'_i \mid i=1,\dots,n\} \subset H^1(L(u);\Lambda)$ *and a map*

$$\Phi : \mathrm{Cliff}_\Lambda(n;\vec d) \to (H(L(u);\Lambda),\cup^{\mathfrak c,Q})$$

with the following properties:

1. Φ *is a* Λ *algebra isomorphism.*
2. $\Phi(X_i) = \mathbf{e}'_i$.
3. *We put* $\Phi(X_I) = \mathbf{e}'_I$. *Then*

$$\langle \mathbf{e}'_I, \mathbf{e}'_J\rangle_{\mathrm{PD}_{L(u)}} = \begin{cases} (-1)^{*(I)} & J = I^c, \\ 0 & \text{otherwise.}\end{cases}$$

Here

$$*(I) = \#\{(i,j) \in I \times I^c \mid j < i\}. \tag{22.3}$$

4. *If* $I \ne \{1,\dots,n\}$, *then*

$$\mathbf{e}'_I \in \bigoplus_{d<n} H^d(L(u);\Lambda).$$

5. $$\mathbf{e}'_{\{1,\dots,n\}} - \mathrm{vol}_{L(u)} \in \bigoplus_{d<n} H^d(L(u);\Lambda).$$

Proof. We first consider the case $\dim L(u) = 2$. (That is the case where (1) of Condition 2.19 holds.) By Lemma 18.24, we have $b^{\mathfrak c} \in H^1(L(u);\Lambda_0)$.

Let $x \in H^1(L(u);\Lambda_0)$. We have

$$\langle x \cup^{\mathfrak c,Q} x, 1\rangle_{\mathrm{PD}_{L(u)}} = \langle x, x\cup^{\mathfrak c,Q} 1\rangle_{\mathrm{PD}_{L(u)}} = \langle x, x\rangle_{\mathrm{PD}_{L(u)}} = 0.$$

Therefore using $\dim L(u) = 2$ again there exists $\mathfrak{Q} : H^1(L(u);\Lambda_0) \to \Lambda_0$ such that

$$x \cup^{\mathfrak{c},Q} x = \mathfrak{Q}(x)1.$$

It is easy to see that $\mathfrak{Q}$ is a quadratic form. In fact, we have

$$\mathfrak{Q}(x) = \frac{1}{2}\frac{d^2}{dt^2}\mathfrak{PO}^{\mathfrak{c}}_{\mathfrak{b}}(b^{\mathfrak{c}} + tx). \tag{22.4}$$

Therefore there exists a basis $\mathbf{e}'_1$, $\mathbf{e}'_2$ of $H^1(L(u);\Lambda_0)$ such that

$$\mathfrak{Q}(x_1\mathbf{e}'_1 + x_2\mathbf{e}'_2) = d_1x_1^2 + d_2x_2^2,$$

where $2d_1, 2d_2 \in \Lambda_0$ are eigenvalues of the Hessian of $\mathfrak{PO}^{\mathfrak{c}}_{\mathfrak{b}}$ at $b^{\mathfrak{c}}$. By replacing $\mathbf{e}'_1$ by $c\mathbf{e}'_1$ for some $c \in \Lambda_0 \setminus \Lambda_+$ we may assume $\mathbf{e}'_1\mathbf{e}'_2 - \mathrm{vol}_{L(u)} \in \bigoplus_{d<n} H^d(L(u);\Lambda)$. It is easy to see that $\mathbf{e}'_1$, $\mathbf{e}'_2$ have required properties.

We next consider the case (2) of Condition 2.19 is satisfied. We start with proving two lemmata. Consider the classical cup product $h_1 \wedge h_2$.

Lemma 22.3. *Assume* (2) *of Condition 2.19. If* $h_i \in H^{d_i}(L(u);\Lambda)$*, then*

$$h_1 \cup^{\mathfrak{c},Q} h_2 - h_1 \wedge h_2 \in \bigoplus_{d<d_1+d_2} H^d(L(u);\Lambda).$$

Proof. By Lemma 18.19, we have $\dim \mathcal{M}^{\mathrm{main};\mathbf{p}}_{3;\ell}(\beta;\mathbf{p}) \geq n+2$ if degree of $\mathbf{p}(i) = 2$. Therefore

$$\begin{aligned}\deg \mathfrak{q}_{\ell;2;\beta}(\mathfrak{b}^{\otimes\ell}, h_1, h_2) &= \deg \mathrm{ev}_{0*}((\mathrm{ev}_1,\mathrm{ev}_2)^*(h_1\times h_2); \mathcal{M}^{\mathrm{main}}_{3;\ell}(\beta;\mathfrak{b}^{\otimes\ell}))\\ &= \deg h_1 + \deg h_2 + n - \dim \mathcal{M}^{\mathrm{main};}_{3;\ell}(\beta;\mathfrak{b}^{\otimes\ell}) \leq d_1 + d_2 - 2.\end{aligned}$$

The lemma follows. ☐

Lemma 22.4. *Assume* (2) *of Condition 2.19. Then we have*

$$\mathbf{e}_i \cup^{\mathfrak{c},Q} \mathbf{e}_j + \mathbf{e}_j \cup^{\mathfrak{c},Q} \mathbf{e}_i = \left(\frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial x_i\partial x_j}(x(b))\right)1.$$

Proof. By definition

$$\mathfrak{PO}^u_{\mathfrak{b}}(x) = \sum_{\beta,\mathbf{p}} c_{\mathbf{p}}\mathfrak{b}^{\mathbf{p}}T^{\beta\cap\omega/2\pi}c(\beta;\mathbf{p})\prod_{i=1}^n \exp((\mathbf{e}_i\cap\partial\beta)x_i).$$

Here we put

$$\sum_{\ell=0}^{\infty}\frac{1}{\ell!}\mathfrak{b}^{\otimes\ell} = \sum_{\mathbf{p}} c_{\mathbf{p}}\mathbf{p}$$

and

$$c(\beta;\mathbf{p}) = \mathrm{ev}_{0*}(\mathcal{M}_{1;|\mathbf{p}|}(\beta;\mathbf{p})).$$

The summation is taken for all β, $\mathbf{p}$ with $\dim \mathcal{M}_{1;|\mathbf{p}|}(\beta;\mathbf{p}) = n$. (See (7.5) [FOOO5].) It is then easy to see that

$$\frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial x_i\partial x_j} = \sum_{\beta,\mathbf{p}} c_{\mathbf{p}}\mathfrak{b}^{\mathbf{p}}T^{\beta\cap\omega/2\pi}c(\beta;\mathbf{p})(\mathbf{e}_i\cap\partial\beta)(\mathbf{e}_j\cap\partial\beta)\prod_{a=1}^n \exp((\mathbf{e}_a\cap\partial\beta)x_a). \tag{22.5}$$

Let h_i be the harmonic one form whose cohomology class is $\mathbf{e}_i$. We can easily prove that

$$\begin{aligned}&\mathrm{ev}_{0*}((\mathrm{ev}_1,\mathrm{ev}_2)^*(h_1\times h_2) + (\mathrm{ev}_1,\mathrm{ev}_2)^*(h_2\times h_1); \mathcal{M}^{\mathrm{main};\mathbf{p}}_{3;|\mathbf{p}|}(\beta;\mathbf{p}))\\ &= c(\beta;\mathbf{p})(\mathbf{e}_i\cap\partial\beta)(\mathbf{e}_j\cap\partial\beta).\end{aligned} \tag{22.6}$$

(See the proof of Lemma 10.8 [FOOO4] and Appendix 1 of the present paper for the relevant argument.) We remark

$$\begin{aligned}&\mathfrak{q}^{\mathfrak{c}}_{\ell;2;\beta}([\mathbf{p}];h_i\otimes h_j)\\&=\Big(\sum_{\beta,\mathbf{p}}c_{\mathbf{p}}\mathfrak{b}^{\mathbf{p}}T^{\beta\cap\omega/2\pi}\left(\prod_{a=1}^{n}\exp((\mathbf{e}_a\cap\partial\beta)x_i)\right)\\&\qquad\times \mathrm{ev}_{0*}((\mathrm{ev}_1,\mathrm{ev}_2)^*(h_i\times h_j);\mathcal{M}^{\mathrm{main};\mathbf{p}}_{3;|\mathbf{p}|}(\beta;\mathbf{p}))^{\mathfrak{c}}\Big)1.\end{aligned}\tag{22.7}$$

Lemma 22.4 follows immediately from (22.5), (22.6), (22.7). □

By the cyclicity of the $\mathfrak{c}$-perturbation we have cyclic property

$$\langle h_1\cup^{\mathfrak{c},Q}h_2,h_3\rangle_{\mathrm{PD}_{L(u)}}=\langle h_1,h_2\cup^{\mathfrak{c},Q}h_3\rangle_{\mathrm{PD}_{L(u)}}\tag{22.8}$$

for $h_1,h_2,h_3\in H(L(u);\Lambda_0)$.

We now take an $n\times n$ matrix $A=[a_{ij}]_{i,j=1}^{i,j=n}$ ($a_{ij}\in\Lambda$) such that

(1)

$$^{t}A\left[\frac{\partial^2\mathfrak{PO}_{\mathfrak{b}}}{\partial x_i\partial x_j}(x(b))\right]_{i,j=1}^{i,j=n}A=2\begin{bmatrix}d_1&0&\dots&0\\0&d_2&\dots&0\\\vdots&\vdots&\ddots&\vdots\\0&0&\dots&d_n\end{bmatrix}.$$

(2) $\det A=1$.

We put

$$x_i'=\sum_j a_{ji}x_i,\qquad \mathbf{e}_i'=\sum_{ij}a_{ij}\mathbf{e}_i.$$

Lemma 22.4 implies that

$$\mathbf{e}_i'\cup^{\mathfrak{c},Q}\mathbf{e}_j'+\mathbf{e}_j'\cup^{\mathfrak{c},Q}\mathbf{e}_i'=2d_i\delta_{ij}.$$

Therefore there exists a ring homomorphism $\Phi:\mathrm{Cliff}_\Lambda(n;\vec{d})\to(H(L(u);\Lambda),\cup^Q)$ satisfying (2) of Theorem 22.2.

We define a filtration on $\mathrm{Cliff}_\Lambda(n;\vec{d})$ by

$$F^l\mathrm{Cliff}_\Lambda(n;\vec{d})=\bigoplus_{\#I\le l}\Lambda Z_I.$$

The associated graded algebra $gr(\mathrm{Cliff}_\Lambda(n;\vec{d}))$ is the exterior algebra. We define a filtration on $H(L(u);\Lambda)$ by

$$F^lH(L(u);\Lambda)=\bigoplus_{k\le l}H^k(L(u);\Lambda).$$

Lemma 22.3 implies that the quantum product $\cup^Q$ respects this filtration and $gr(H(L(u);\Lambda))$ is isomorphic to the exterior algebra as a ring.

Therefore Φ induces an isomorphism $gr(\mathrm{Cliff}_\Lambda(n;\vec{d}))\to gr(H(L(u);\Lambda))$. It implies that Φ itself is an isomorphism. Hence follows statement (1).

Statement (4) follows immediately from Lemma 22.3 and the definition. By Lemma 22.3 we have

$$\begin{aligned}\mathbf{e}'_{\{1,\dots,n\}}&=\mathbf{e}_1'\cup^{\mathfrak{c},Q}\dots\cup^{\mathfrak{c},Q}\mathbf{e}_n'\equiv\mathbf{e}_1'\wedge\dots\wedge\mathbf{e}_n'\\&\equiv\det A\ \ \mathbf{e}_1\wedge\dots\wedge\mathbf{e}_n=\mathrm{vol}_{L(u)}\qquad\mathrm{mod}\ \bigoplus_{d<n}H^d(L(u);\Lambda).\end{aligned}$$

(5) follows.

For $I = \{i_1, \dots, i_k\} \in 2^{\{1,\dots,n\}}$ we put $d^I = d_{i_1} \cdots d_{i_k}$. By the Clifford relation (22.1), we have

$$\mathbf{e}'_I \cup^{\mathfrak{c},Q} \mathbf{e}'_J = (-1)^* d^{I \cap J} \mathbf{e}'_{I \ominus J} \tag{22.9}$$

where $I \ominus J = (I \cup J) \setminus (I \cap J)$. (We do not calculate the sign $*$ in (22.9) since we do not need it.) (22.9) and (4) imply that $\mathbf{e}'_I \cup^{\mathfrak{c},Q} \mathbf{e}'_J$ has no degree n component if $J \neq I^c$. Therefore using (22.8), we have:

$$\langle \mathbf{e}'_I, \mathbf{e}'_J \rangle_{\mathrm{PD}_{L(u)}} = \langle \mathbf{e}'_I, \mathbf{e}'_J \cup^{\mathfrak{c},Q} 1 \rangle_{\mathrm{PD}_{L(u)}} = \langle \mathbf{e}'_I \cup^{\mathfrak{c},Q} \mathbf{e}'_J, 1 \rangle_{\mathrm{PD}_{L(u)}} = \int_{L(u)} \mathbf{e}'_I \cup^{\mathfrak{c},Q} \mathbf{e}'_J = 0.$$

Remark 22.5. This is another point for which we use the cyclic symmetry in an essential way.

(22.9) implies

$$\begin{aligned} \mathbf{e}'_I \wedge \mathbf{e}'_{I^c} \equiv \mathbf{e}'_I \cup^{\mathfrak{c},Q} \mathbf{e}'_{I^c} &\equiv (-1)^{*(I)} \mathbf{e}'_{\{1,\dots,n\}} \\ &\equiv (-1)^{*(I)} \mathrm{vol}_{L(u)} \mod \bigoplus_{d<n} H^d(L(u);\Lambda). \end{aligned}$$

Therefore we have

$$\langle \mathbf{e}'_I, \mathbf{e}'_{I^c} \rangle_{\mathrm{PD}_{L(u)}} = \int_{L(u)} \mathbf{e}'_I \wedge \mathbf{e}'_{I^c} = (-1)^{*(I)}.$$

Here $*(I)$ is as in (23.6). We have thus proved statement (2). The proof of Theorem 22.2 is now complete. □

23. Residue pairing and Hessian determinant

In this section we complete the proof of Theorem 2.24. We first prove (2) and (3). Using Theorem 22.2 (3) and (22.9) we calculate

$$Z((H(L(u);\Lambda), \langle \cdot, \cdot \rangle_{\mathrm{PD}_{L(u)}}, \cup^{\mathfrak{c},\mathfrak{b},b}, PD[L(u)]))$$

defined as (2.18). We consider the nonzero terms in (2.18). Since $g^{I_3 0} g^{J_3 0} \neq 0$, it follows from Theorem 22.2 (3) that $I_3 = J_3 = \{1, \dots, n\}$. Then $\langle \mathbf{e}_{I_1} \cup^{\mathfrak{c},Q} \mathbf{e}_{I_2}, \mathbf{e}_{I_3} \rangle \neq 0$ implies $I_1 = I_2$, which we put I.

Similarly we have $J_1 = J_2$. $g^{I_1 J_1} \neq 0$ implies $J_1 = J_2 = I^c$. Thus we have

$$\begin{aligned} &Z((H(L(u);\Lambda), \langle \cdot, \cdot \rangle_{\mathrm{PD}_{L(u)}}, \cup^{\mathfrak{c},\mathfrak{b},b}, PD[L(u)])) \\ &= \sum_{I \in 2^{\{1,\dots,n\}}} (-1)^{*_{I,1}} \langle \mathbf{e}_I \cup^{\mathfrak{c},Q} \mathbf{e}_I, \mathbf{e}_{1,\dots,n} \rangle_{PD_{L(u)}} \langle \mathbf{e}_{I^c} \cup^{\mathfrak{c},Q} \mathbf{e}_{I^c}, \mathbf{e}_{1,\dots,n} \rangle_{PD_{L(u)}} \\ &= \sum_{I \in 2^{\{1,\dots,n\}}} (-1)^{*_{I,2}} d_1 \dots d_n. \end{aligned} \tag{23.1}$$

If $\# I = p$, then

$$*_{I,1} = p(n-p) + n(n-1)/2.$$

(This sign comes from the sign $*$ in (2.18).) We then have

$$*_{I,2} = p(p-1)/2 + (n-p)(n-p-1)/2 + *_{I,1} \equiv 0 \mod 2.$$

Therefore (23.1) is equal to

$$2^n d_1 \cdots d_n = \det \left[\frac{\partial^2 \mathfrak{PO}^{\mathfrak{c}}_{\mathfrak{b}}}{\partial x_i \partial x_j}(x(b)) \right]_{i,j=1}^{i,j=n}. \tag{23.2}$$

Thus Theorem 2.24 (3) holds. (2) also follows using equality $\mathfrak{PO}^{\mathfrak{c}}_{\mathfrak{b}} = \mathfrak{PO}_{\mathfrak{b}}$ ((22.2) and Corollary 18.26).

We next prove (1) of Theorem 2.24. Let X be a compact toric manifold and $\mathfrak{b} \in H(X;\Lambda_0)$. Let $(u,b) \in \mathfrak{M}(X,\mathfrak{b})$.

Here we use $\mathfrak{q}$-perturbation (and not $\mathfrak{c}$-perturbation) to define $\mathfrak{M}(X,\mathfrak{b})$. We study the operation $\mathfrak{m}_2^{\mathfrak{b},b}$ (and not $\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b}$) which is defined by $\mathfrak{q}$-perturbation at the first step of the proof. So, at the first step of the proof, we consider $b \in H^1(L(u);\Lambda_0)$.

We assume that b is a nondegenerate critical point of $\mathfrak{PO}_{\mathfrak{b}}$. We define $\cup^Q$ by

$$h_1 \cup^Q h_2 = (-1)^{\deg h_1(\deg h_2+1)}\mathfrak{m}_2^{\mathfrak{b},b}(h_1,h_2).$$

We first prove the following variant of Theorem 22.2.

Proposition 23.1. *There exist a basis* $\{\mathbf{e}'_i \mid i=1,\dots,n\} \subset H^1(L(u);\Lambda)$ *and a map*

$$\Phi : \mathrm{Cliff}_{\Lambda}(n;\vec{d}) \to (H(L(u);\Lambda),\cup^Q)$$

with the following properties:

1. Φ *is a* Λ *algebra isomorphism.*
2. $\Phi(X_i) = \mathbf{e}'_i$.
3. $\{\mathbf{e}'_i \mid i=1,\dots,n\} \subset H^1(L(u);\Lambda_0)$. *Moreover* $\{\mathbf{e}'_i \mod H^1(L(u);\Lambda_+) \mid i = 1,\dots,n\}$ *form a basis of* $H^1(L(u);\mathbb{C})$.
4. $$\mathbf{e}'_{\{1,\dots,n\}} - \mathrm{vol}_{L(u)} \in \bigoplus_{d<n} H^d(L(u);\Lambda_+).$$

Proof. We start with the following lemma.

Lemma 23.2. *Let* H *be a symmetric* $n\times n$ *matrix with* Λ_0 *entries. Then there exists* $\mathbf{e}'_i \in \Lambda_0^n$, $i=1,\dots,n$, *with the following properties:*

1. $\{\mathbf{e}'_i \mid i=1,\dots,n\}$ *form a basis of* Λ^n.
2. $\{\mathbf{e}'_i \mod \Lambda_+ \mid i=1,\dots,n\}$ *form a basis of* $\mathbb{C}^n$.
3. $(\mathbf{e}'_i, H\mathbf{e}'_j) = 0$ *for* $i \ne j$. *Here* $(\cdot,\cdot) : \Lambda^n \times \Lambda^n \to \Lambda$ *is the standard quadratic form.*

Proof. We take $\lambda \ge 0$ the largest number such that all the entries of $T^{-\lambda}H$ are in Λ_0. We may assume $\lambda = 0$ without loss of generality.

Let $\overline{H} \equiv H \mod \Lambda_+$ be the complex $n\times n$ matrix. It is nonzero and symmetric. We can take a basis $\bar{\mathbf{f}}_i$ of $\mathbb{C}^n$ such that $(\bar{\mathbf{f}}_i, \overline{H}\bar{\mathbf{f}}_j) = 0$ for $i \ne j$. We may assume furthermore that $(\bar{\mathbf{f}}_i, \overline{H}\bar{\mathbf{f}}_i) = 1$ for $i = 1,\dots,k$ and $(\bar{\mathbf{f}}_i, \overline{H}\bar{\mathbf{f}}_i) = 0$ for $i = k+1,\dots,n$. ($k > 0$.) It is easy to lift $\bar{\mathbf{f}}_i$ to $\mathbf{f}_i \in \Lambda_0^n$ such that $(\mathbf{f}_i, H\mathbf{f}_j) = 0$ holds for any $i = 1,\dots,k$ and $j \ne i$. Then, with respect to the splitting $\Lambda_0^n \cong \Lambda_0^k \oplus \Lambda_0^{n-k}$, H also splits so that its restriction to the first factor has the required basis $\mathbf{f}_i$, $i=1,\dots,k$. Thus the lemma is proved by induction on n. □

We apply Lemma 23.2 to the Hessian matrix $\mathrm{Hess}_b(\mathfrak{PO}_{\mathfrak b})$ and obtain $\mathbf{e}'_i$. We may replace $\mathbf{e}'_1$ by $c\mathbf{e}'_1$ for some $c \in \Lambda_0 \setminus \Lambda_+$ so that we have the following in addition:

$$\det\,[\mathbf{e}'_1 \dots \mathbf{e}'_n] = 1. \tag{23.3}$$

(Here we regard each of $\mathbf{e}'_i$ as $1 \times n$ matrix.)

Now statement (3) is satisfied.

We next observe that the operator $\mathfrak{m}_2^{\mathfrak b, b}$, that is constructed by $\mathfrak q$-perturbation, satisfies the conclusion of Lemma 22.3, in our situation where Condition 2.19 (2),(3) is *not* assumed. In fact we can use [FOOO5] Corollary 6.6 in place of Lemma 18.6.

It implies

$$\mathbf{e}'_I - \mathbf{e}'_{i_1} \wedge \cdots \wedge \mathbf{e}'_{i_k} \in \bigoplus_{\ell < k} H^\ell(L(u);\Lambda_+) \tag{23.4}$$

for $I = (i_1,\dots,i_k)$. Here we put $\mathbf{e}'_I = \mathbf{e}'_{i_1} \cup^Q \cdots \cup^Q \mathbf{e}'_{i_k}$. We use the fact that the contribution of pseudo-holomorphic discs to $\mathfrak{m}_2$ is of positive energy to show that the difference in (23.4) is 0 modulo Λ_+.

Lemma 23.2 (3) and (23.4) imply

$$\mathbf{e}'_i \cup^Q \mathbf{e}'_j + \mathbf{e}'_j \cup^Q \mathbf{e}'_i = \begin{cases} 2d_i & \text{if } i = j \\ 0 & \text{if } i \neq j, \end{cases}$$

for some $d_i \in \Lambda_+$. Then if we define the map Φ by (2), statement (1) follows.

(4) follows from (23.3) and (23.4). The proof of Proposition 23.1 is complete. □

We remark that d_i are eigenvalues of the Hessian of $\mathfrak{PO}^{\mathfrak b}$ with respect to the coordinate $\mathbf{e}'_i$. Namely if we write $b \in H^1(L(u);\Lambda_0)$ as $b = \sum x'_i \mathbf{e}'_i$ then

$$\det\left[\frac{\partial^2 \mathfrak{PO}^{\mathfrak b}}{\partial x'_i \partial x'_j}(b)\right]_{i,j=1}^{i,j=n} = d_1 \dots d_n.$$

In fact the matrix in the left hand side is the diagonal matrix with diagonal entries d_i. Then using (23.3) we have

$$\det\left[\frac{\partial^2 \mathfrak{PO}^{\mathfrak b}}{\partial x_i \partial x_j}(b)\right]_{i,j=1}^{i,j=n} = d_1 \dots d_n. \tag{23.5}$$

Here x_i is a coordinate of $H^1(L(u);\Lambda)$ with respect to the basis $\mathbf{e}_i$.

Remark 23.3. Since we do not have cyclic symmetry, Proposition 23.1 does *not* imply $\langle \mathbf{e}'_I, \mathbf{e}'_J \rangle_{PD_{L(u)}} = 0$ for $J \neq I^c$. We only have

$$\langle \mathbf{e}'_I, \mathbf{e}'_J \rangle_{\mathrm{PD}_{L(u)}} \equiv \begin{cases} (-1)^{*(I)} & \mod \Lambda_+ \quad \text{for } J = I^c, \\ 0 & \mod \Lambda_+ \quad \text{otherwise.} \end{cases}$$

Here we denote

$$*\,(I) = \#\{(i,j) \in I \times I^c \mid j < i\}. \tag{23.6}$$

Now we move from $\mathfrak{m}_2^{\mathfrak b, b}$ to $\mathfrak{m}_2^{\mathfrak c, \mathfrak b, b^{\mathfrak c}}$, where $b^{\mathfrak c}$ is obtained from b as in Definition 18.27. Then by Corollary 18.22 we have a ring isomorphism

$$\mathfrak{F}_1 : (H(L(u);\Lambda_0), \cup^Q) \to (H(L(u);\Lambda_0), \cup^{\mathfrak c, Q}).$$

Here we remark that $\cup^Q$ is induced from $\mathfrak{m}_2^{\mathfrak b, b}$ and $\cup^{\mathfrak c, Q}$ is induced from $\mathfrak{m}_2^{\mathfrak c, \mathfrak b, b^{\mathfrak c}}$.

We now define

$$\mathbf{e}''_i = \mathfrak{F}_1(\mathbf{e}'_i) \tag{23.7}$$

and

$$\mathbf{e}''_I = \mathbf{e}''_{i_1} \cup^{\mathfrak{c},Q} \cdots \cup^{\mathfrak{c},Q} \mathbf{e}''_{i_k}. \tag{23.8}$$

Now we have

Proposition 23.4. *There exists a map*

$$\Psi : \mathrm{Cliff}_\Lambda(n; \vec{d}) \to (H(L(u);\Lambda), \cup^{\mathfrak{c},Q})$$

with the following properties:

1. Ψ *is a* Λ *algebra isomorphism.*
2. $\Psi(X_i) = \mathbf{e}''_i$.
3. $$\mathbf{e}''_{\{1,\dots,n\}} - \mathrm{vol}_{L(u)} \in \bigoplus_{d \le n} H^d(L(u);\Lambda_+).$$

This is immediate from Proposition 23.1 and $\mathfrak{F}_1 - id \equiv 0 \mod \Lambda_+$.

Now we are in the position to complete the proof of Theorem 2.24 (1). We have $\mathfrak{o} \in \Lambda_+$ such that

$$(1+\mathfrak{o})\mathrm{vol}_{L(u)} - \mathbf{e}''_{1,\dots,n} \in \bigoplus_{k<n} H^k(L(u);\Lambda).$$

Using Corollary 19.9 we find that the formula (20.2) for $\mathfrak{v} = \mathfrak{w} = (1+\mathfrak{o})\mathrm{vol}_{L(u)}$ is equal to (20.2) for $\mathfrak{v} = \mathfrak{w} = \mathbf{e}''_{1,\dots,n}$. Namely

$$\begin{aligned}(1+\mathfrak{o})^2 &\langle i_{\#,\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}})}(\mathrm{vol}_{L(u)}), i_{\#,\mathrm{qm},(\mathfrak{b},b^{\mathfrak{c}})}(\mathrm{vol}_{L(u)})\rangle_{\mathrm{PD}_X} \\ = &\sum_{I,J \in 2^{\{1,\dots,n\}}} (-1)^{*_{1,I,J}} g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_I, \mathbf{e}''_{1,\dots,n}), \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n})\rangle_{\mathrm{PD}_{L(u)}}.\end{aligned} \tag{23.9}$$

Using cyclic symmetry and Proposition 23.4, Formula (23.9) is equal to

$$\begin{aligned}&\sum_{I,J \in 2^{\{1,\dots,n\}}} (-1)^{*_{2,I,J}} g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_{1,\dots,n}, \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n})), \mathbf{e}''_I\rangle_{\mathrm{PD}_{L(u)}} \\ &= \sum_{I,J \in 2^{\{1,\dots,n\}}} (-1)^{*_{3,I,J}} g^{IJ} d_J \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_{1,\dots,n}, \mathbf{e}''_{J^c}), \mathbf{e}''_I\rangle_{\mathrm{PD}_{L(u)}} \\ &= \sum_{I,J \in 2^{\{1,\dots,n\}}} (-1)^{*_{4,I,J}} g^{IJ} d_1 \cdots d_n \langle \mathbf{e}''_J, \mathbf{e}''_I\rangle_{\mathrm{PD}_{L(u)}} = 2^n d_1 \cdots d_n.\end{aligned}$$

Here $d_I = d_{i_1} \dots d_{i_k}$ for $I = (i_1, \dots, i_k)$.

The calculation of the sign is as follows. We put $|I| = k$, $|J| = \ell$. Note we only need to calculate sign in case $\ell \equiv n-k \mod 2$. Now

$$\begin{aligned} *_{1,I,J} &= \frac{n(n-1)}{2} \\ *_{2,I,J} &= *_{1,I,J} + \ell(k+1). \end{aligned}$$

Here we get the sign $(n+k)(\ell+n+1)$ to go from $\langle \cdot,\cdot \rangle_{PD_{L(u)}}$ to $\langle \cdot,\cdot\rangle_{\mathrm{cyc}}$ and $(k+1)(n+1+n+\ell+1)$ for the cyclic symmetry and $\ell(k+1)$ to back from $\langle \cdot,\cdot\rangle_{\mathrm{cyc}}$ to $\langle \cdot,\cdot \rangle_{PD_{L(u)}}$. The sum of these three is $\ell(k+1)$.

We next have

$$*_{3,I,J} = *_{2,I,J} + \ell(n+1) + \frac{\ell(\ell-1)}{2} + *(J).$$

Here $*(J)$ is as in (23.6). In fact, we get $\ell(n+1)$ to go from $\mathfrak{m}_2$ to the product, and

$$\mathbf{e}''_J \cup^{\mathfrak{c},Q} \mathbf{e}''_{1,\dots,n} = (-1)^{\frac{\ell(\ell-1)}{2}+*(J)}\mathbf{e}''_{J^c}$$

by Clifford relation. Similarly we have

$$*_{4,I,J} = *_{3,I,J} + n(k+1) + \frac{(n-\ell)(n-\ell-1)}{2} + *(J).$$

We can easily find that $*_{4,I,J} \equiv 0 \mod 2$ by using $k+\ell \equiv n \mod 2$.

The above calculation and (23.5) imply Theorem 2.24 (1). □

Remark 23.5. We remark that $\mathbf{e}''_i \in H^1(L(u);\Lambda)$ may *not* hold. $\mathbf{e}''_i$, $(i=1,\dots,n)$ form a basis of the tangent space $T_{b^{\mathfrak{c}}}(\mathfrak{F}_*(H^1(L(u);\Lambda_0)))$.

24. CYCLIC HOMOLOGY AND VARIATION OF THE INVARIANT Z

The purpose of this section is to put the invariant $Z(H,\{\mathfrak{m}_k\},\langle\cdot\rangle,e)$ (Definition 2.21) of cyclic filtered A_∞ algebra $(H,\{\mathfrak{m}_k\},\langle\cdot\rangle,e)$ in the perspective of several other stories such as cyclic cohomology.

We begin with a review of cyclic (co)homology of A_∞ algebra which is a minor modification of the story of cyclic (co)homology of associative algebras. ([Con], [Lo].)

Let K be a field of characteristic 0 and $(C,\{\mathfrak{m}_k\}_{k=0}^{\infty},e)$ an A_∞ algebra with unit e. We assume the following condition.

Condition 24.1. (1) C is a finite dimensional K vector space.

(2) $\mathfrak{m}_0(1) = ce$ for some $c \in K$.

(3) $\mathfrak{m}_1 = 0$.

Hereafter in this section, we replace $\mathfrak{m}_0$ by 0. Namely we put $\mathfrak{m}_0(1)=0$. Thanks to item (2) above, we still have a filtered A_∞ algebra, after this replacement .

We define $B_k(C[1])$ in the same way. We define cyc on them as in (17.6). Let $B_k^{\rm cyc}(C[1])$ be the quotient space as in Section 17. We put

$$B^{\rm cyc}(C[1]) = \bigoplus_{k=0}^{\infty} B_k^{\rm cyc}(C[1]). \tag{24.1}$$

The A_∞ operation $\mathfrak{m}_k$ defines a differential $\delta^{\rm cyc}$ on them by (17.8).

Definition 24.2. The *cyclic homology* $HC_*(C)$ is the homology group of the chain complex $(B^{\rm cyc}(C[1]),\delta^{\rm cyc})$.

The *cyclic cohomology* $HC^*(C)$ is the homology group of the cochain complex $(Hom(B^{\rm cyc}(C[1]),K),(\delta^{\rm cyc})^*)$.

We put

$$F_k B^{\rm cyc}(C[1]) = \bigoplus_{\ell=0}^{k} B_\ell^{\rm cyc}(C[1]) \tag{24.2}$$

and

$$F^k Hom(B^{\rm cyc}(C[1]),K) = \prod_{\ell=k}^{\infty} Hom(B_\ell^{\rm cyc}(C[1]),K). \tag{24.3}$$

By Condition 24.1, that is $\mathfrak{m}_0 = 0$, we have

$$\delta^{\rm cyc}\left(F_k B^{\rm cyc}(C[1])\right) \subset F_{k-1}B^{\rm cyc}(C[1]),$$
$$(\delta^{\rm cyc})^*\left(F^k Hom(B^{\rm cyc}(C[1]),K)\right) \subset F^{k+1}Hom(B^{\rm cyc}(C[1]),K).$$

Therefore we have homomorphisms

$$H(F_kB^{\rm cyc}(C[1]),\delta^{\rm cyc}) \to HC_*(C), \quad H(Hom(F^kB^{\rm cyc}(C[1]),K),(\delta^{\rm cyc})^*) \to HC^*(C).$$

We denote their images by $F_kHC_*(C)$, $F^kHC^*(C)$.

For a unital A_∞ algebra $(C,\{\mathfrak{m}_k\}_{k=0}^\infty,e)$ satisfying Condition 24.1, $(C,\mathfrak{m}_2,e)$ becomes an (associative) algebra with unit. We call it the *underlying algebra.*

Proposition 24.3 (Compare [Ka] section 6)**.** *If the underlying algebra of $(C,\{\mathfrak{m}_k\}_{k=0}^\infty,e)$ is isomorphic to the Clifford algebra associated to a non-degenerate quadratic form, then*

$$\frac{F_kHC_*(C)}{F_{k-1}HC_*(C)} \cong \begin{cases} K & k \text{ is even,} \\ 0 & k \text{ is odd.}\end{cases} \tag{24.4}$$

Moreover all the nonzero elements in the cyclic homology $HC_*(C)$ have degree $* \equiv n-1 \mod 2$. (Here we use the shifted degree.)

Proof. Following [Ka], we first consider Hochschild homology. We consider $BC = \bigoplus_{k=0}^\infty B_k(C[1])$ and define $\delta^{\rm cyc} : B(C[1]) \to B(C[1])$ by (17.8).

Definition 24.4. We call the homology group $H(B(C[1]),\delta^{\rm cyc})$, the *Hochschild homology* and denote it by $HH_*(C,C)$.

The boundary operator $\delta^{\rm cyc}$ respects the number filtration $F_kB(C[1])$. Therefore it induces a filtration $F_kHH_*(C,C)$ on Hochschild homology.

Lemma 24.5. *If the underlying algebra of C is isomorphic to the Clifford algebra associated to a non-degenerate quadratic form, then $HH_*(C,C) \cong K$. It is generated by the element*

$$e_{1\cdots n} \in C = B_1(C[1]). \tag{24.5}$$

In particular, $HH_*(C,C) = F_1HH_*(C,C)$.

Proof. Let C' be the underlying Clifford algebra regarded as an A_∞ algebra. (Namely the $\mathfrak{m}_2$ of C' is the same as that of C and all higher $\mathfrak{m}_k$ of C' is zero.) We consider the spectral sequence $E_k(C)$ induced by the number filtration $F_kB(C[1])$. It is easy to see that $E_2(C) \cong E_2(C') \cong E_\infty(C')$. On the other hand, as in [Ka] page 696, $E_\infty(C') \cong K$ and generated by the element (24.5). Therefore the spectral sequence for C also degenerates at E_2 term. The lemma follows. □

Now we use the following variant of the famous result by Connes [Con]. This A_∞ version is given, for example, in [CL], [HL].

Theorem 24.6. *There exists a long exact sequence*

$$\longrightarrow HC_*(C,C) \xrightarrow{S} HC_*(C,C) \xrightarrow{B} HH_*(C,C) \xrightarrow{I} HC_*(C,C) \xrightarrow{S} \tag{24.6}$$

such that

$$S(F_kH_*C(C,C)) \subset F_{k-2}HC_*(C,C), \tag{24.7a}$$
$$B(F_kHC_*(C,C)) \subset F_{k+1}HH_*(C,C), \tag{24.7b}$$
$$I(F_kHH_*(C,C)) \subset F_kHC_*(C,C). \tag{24.7c}$$

Proof. (Sketch) We consider the diagram:

$$\begin{tikzcd}
{} \arrow[d] & {} \arrow[d] & {} \arrow[d] & \\
F_3B(C[1]) \arrow[d, "\delta^{\mathrm{cyc}}"] & F_3B(C[1]) \arrow[l, "1-\mathrm{cyc}"'] \arrow[d, "\widehat{d}"] & F_3B(C[1]) \arrow[l, "N"'] \arrow[d, "\delta^{\mathrm{cyc}}"] & \ldots \arrow[l, "1-\mathrm{cyc}"'] \\
F_2B(C'[1]) \arrow[d, "\delta^{\mathrm{cyc}}"] & F_2B(C[1]) \arrow[l, "1-\mathrm{cyc}"'] \arrow[d, "\widehat{d}"] & F_2B(C[1]) \arrow[l, "N"'] \arrow[d, "\delta^{\mathrm{cyc}}"] & \ldots \arrow[l, "1-\mathrm{cyc}"'] \\
C[1] & C[1] \arrow[l, "1-\mathrm{cyc}"'] & C[1] \arrow[l, "N"'] & \ldots \arrow[l, "1-\mathrm{cyc}"']
\end{tikzcd} \tag{24.8}$$

See [Lo] page 54. Here $N = \sum_{i=0}^{k-1} \mathrm{cyc}^i$ on $B_k(C'[1])$.

We take infinitely many copies of $B(C[1])$ and denote them by $B(C[1])_\ell$, $\ell = 1, 2, \ldots$. We define a map

$$B(C[1])_{2\ell} \longrightarrow B(C[1])_{2\ell-1}$$

as $1-\mathrm{cyc}$ and

$$B(C[1])_{2\ell+1} \longrightarrow B(C[1])_{2\ell}$$

as N. Moreover we define a map

$$B(C[1])_{2\ell} \longrightarrow B(C[1])_{2\ell}$$

as $-\widehat{d}$ and

$$B(C[1])_{2\ell-1} \longrightarrow B(C[1])_{2\ell-1}$$

as δ^{cyc}. The four assignments combined define

$$\widehat{\widehat{d}} : \bigoplus_{\ell=1}^{\infty} B(C[1])_\ell \to \bigoplus_{\ell=1}^{\infty} B(C[1])_\ell.$$

The commutativity of (24.8) implies $\widehat{\widehat{d}} \circ \widehat{\widehat{d}} = 0$.

In the same way as the proof of [Lo] Theorem 2.1.5, we can prove that

$$HC(C) \cong H\left(\bigoplus_{\ell=1}^{\infty} B(C[1])_\ell, \widehat{\widehat{d}}\right).$$

We also remark that the $\widehat{d}$ cohomology always vanishes.

On the other hand, we have an exact sequence of chain complecies

$$0 \to \bigoplus_{\ell=3}^{\infty} B(C[1])_\ell \to \bigoplus_{\ell=1}^{\infty} B(C[1])_\ell \to \bigoplus_{\ell=1}^{2} B(C[1])_\ell \to 0.$$

The homology of the second complex is the cyclic homology. The homology of the first complex is the cyclic homology with degree shifted by 2. The homology of the third complex is the homology of δ^{cyc} complex, that is the Hochschild homology. The theorem follows. □

Now we are ready to complete the proof of Proposition 24.3. By Lemma 24.5 we have $F_0HH(C,C) \cong K$. By (24.6) and (24.7b) $I : HH_*(C,C) \to HC_*(C,C)$ is injective on $F_0HH(C,C)$. Therefore the generator of $F_0HH(C,C)$ induces a nontrivial element of $HH_*(C,C)$. Then using $HH(C,C) \cong F_0HH(C,C) \cong K$ and (24.6) we can prove that S induces an isomorphism

$$F_kHC_*(C,C)/F_{k-1}HC_*(C,C) \cong F_{k-2}HC_*(C,C)/F_{k-3}HC_*(C,C)$$

for $k > 0$. (24.4) follows easily.

We remark that

$$\deg' e_{1\dots n} = \deg e_{1\dots n} - 1 = n-1.$$

Moreover S does not change the parity of the degree. The statement on degree follows. □

It is well known among the experts that cyclic cohomology controls deformations of cyclic A_∞ algebras. Explanation of this is now in order. (In a similar way, deformations of A_∞ algebras are controlled by Hochschild cohomology. See [FOOO3] Subsection 7.4.6, for example.)

Let $(C, \{\mathfrak{m}_k\}, e)$ be a unital A_∞ algebra satisfying Condition 24.1. Let $\langle \cdot \rangle : C^k \otimes C^{n-k} \to K$ be a nondegenerate bilinear form such that

$$\begin{aligned} &\langle \mathfrak{m}_k(x_1,\dots,x_k), x_0\rangle \\ &= (-1)^{\deg' x_0(\deg' x_1+\dots+\deg' x_k)}\langle \mathfrak{m}_k(x_0,x_1,\dots,x_{k-1}), x_k\rangle. \end{aligned} \tag{24.9}$$

In other words, $(C, \{\mathfrak{m}_k\}, \langle \cdot \rangle, e)$ is a *cyclic* unital A_∞ algebra.

If $\mathfrak{m}_k$ is defined only for $k \le K$ and if the A_∞ relation

$$\sum_{\substack{k_1,k_2\\ k_1+k_2=k}} \sum_i (-1)^{\deg' x_1+\dots+\deg' x_{i-1}} \mathfrak{m}_{k_1}(x_1,\dots,\mathfrak{m}_{k_2}(x_i,\dots),\dots) = 0 \tag{24.10}$$

is satisfied only for $k \le K$, we call it a *cyclic unital A_K algebra.*

We next consider the ring

$$K_m = \begin{cases} \dfrac{K[\theta]}{(\theta^{m+1})} & \text{if } m < \infty, \\ K[[\theta]] & \text{if } m = \infty. \end{cases} \tag{24.11}$$

Definition 24.7. An A_K *deformation of order* m of $(C, \{\mathfrak{m}_k\}, \langle\cdot\rangle)$ is an operations $\mathfrak{m}_k^\theta$ on $C \otimes A_m$ such that (24.9) and (24.10) hold for $k \le K$ and that $\mathfrak{m}_k^\theta \equiv \mathfrak{m}_k$ mod θ.

Remark 24.8. We do not deform $\langle\cdot\rangle$ or e here but only the operation $\mathfrak{m}_k$.

A pseudo-isotopy of cyclic A_∞ algebra is defined by a family of operators

$$\mathfrak{m}_k^t : B_k(C[1]) \to C[1], \quad \mathfrak{c}_k^t : B_k(C[1]) \to C[1]$$

parametrized by $t \in [0,1]$ such that:

(1) They are smooth with respect to t in the sense of Definition 18.15.
(2) For each t, $(C, \{\mathfrak{m}_k^t\}, \langle\cdot\rangle)$ is a cyclic A_∞ algebra.
(3)

$$\langle \mathfrak{c}_{k,\beta}^t(x_1,\dots,x_k), x_0\rangle = (-1)^*\langle \mathfrak{c}_{k,\beta}^t(x_0,x_1,\dots,x_{k-1}), x_k\rangle \tag{24.12}$$

$* = (\deg \mathbf{x}_0 + 1)(\deg \mathbf{x}_1 + \ldots + \deg \mathbf{x}_k + k)$.

(4) For each $x_i \in \overline{C}[1]$

$$\begin{aligned}&\frac{d}{dt}\mathfrak{m}^t_{k,\beta}(x_1,\dots,x_k)\\ &+\sum_{k_1+k_2=k}\sum_{\beta_1+\beta_2=\beta}\sum_{i=1}^{k-k_2+1}(-1)^*\mathfrak{c}^t_{k_1,\beta_1}(x_1,\dots,\mathfrak{m}^t_{k_2,\beta_2}(x_i,\dots),\dots,x_k)\\ &-\sum_{k_1+k_2=k}\sum_{\beta_1+\beta_2=\beta}\sum_{i=1}^{k-k_2+1}\mathfrak{m}^t_{k_1,\beta_1}(x_1,\dots,\mathfrak{c}^t_{k_2,\beta_2}(x_i,\dots),\dots,x_k)\\ &=0.\end{aligned} \tag{24.13}$$

Here $* = \deg' x_1 + \cdots + \deg' x_{i-1}$.

We can define a pseudo-isotopy of cyclic A_K structure and also of an algebra over K_m in the same way.

Definition 24.9. Two A_K deformations of order m are said to be *pseudo-isotopic* if there exists a pseudo-isotopy of cyclic A_K structure over K_m such that its operators $\mathfrak{m}^t_k$ and $\mathfrak{c}^t_k$ satisfy:

$$\frac{d\mathfrak{m}^t_k}{dt} \equiv 0 \mod \theta, \quad \mathfrak{c}^t_k \equiv 0 \mod \theta.$$

We can prove that a pseudo-isotopy is an equivalence relation in the same way as in [Fu2] Lemma 8.2.

Proposition 24.10. *The set of pseudo-isotopy classes of A_N deformations of order 1 of $(C, \{\mathfrak{m}_k\}, \langle\cdot\rangle)$ corresponds one to one to the set*

$$\begin{aligned}&HC^*(C)/F^{N+2}HC^*(C) \quad (\textit{if } N<\infty),\\ &\operatorname*{proj\,lim}_m HC^*(C)/F^{m+2}HC^*(C) \quad (\textit{if } N=\infty),\end{aligned}$$

with $* \equiv n+1 \mod 2$.

Proof. Let $\mathfrak{m}^\theta_k = \mathfrak{m}_k + \theta\Delta\mathfrak{m}_k$ be an A_K deformation of order one. We put

$$\Delta^+\mathfrak{m}_k(x_1,\dots,x_k,x_0) = \langle \Delta\mathfrak{m}_k(x_1,\dots,x_k), x_0\rangle.$$

Then

$$(\Delta^+\mathfrak{m}_0,\dots,\Delta^+\mathfrak{m}_N) \in \frac{\prod_{k=0}^\infty Hom(B^{\mathrm{cyc}}_k(C[1]),K)}{\prod_{k=N+2}^\infty Hom(B^{\mathrm{cyc}}_k(C[1]),K)}. \tag{24.14}$$

The A_K relation among them implies that the left hand side of (24.14) is a $(\delta^{\mathrm{cyc}})^*$ cocycle.

Suppose $\mathfrak{m}_k + \theta\Delta\mathfrak{m}_k$ is pseudo-isotopic to $\mathfrak{m}_k + \theta\Delta'\mathfrak{m}_k$. Take a pseudo-isotopy $\mathfrak{m}^t_k$, $\mathfrak{c}^t_k$ among them. Since $\mathfrak{c}^t_k \equiv 0 \mod \theta$ we have $\mathfrak{c}^t_k = \overline{\mathfrak{c}}^t_k\theta$. We put

$$\mathfrak{e}_k = \int_0^1 \overline{\mathfrak{c}}^t_k dt,$$

and define $\mathfrak{e}^+_k$ in the same way. It follows easily from (24.13) that

$$(\Delta\mathfrak{m}_0,\dots,\Delta\mathfrak{m}_N) - (\Delta'\mathfrak{m}_0,\dots,\Delta'\mathfrak{m}_N) = \delta^{\mathrm{cyc}}(\mathfrak{e}^+_0,\dots,\mathfrak{e}_N).$$

We have thus defined a homomorphism to $HC^*(C)/F^{N+2}HC^*(C)$ from the set of pseudo-isotopy classes of A_m-deformations of $(C, \{\mathfrak{m}_k\}, \langle\cdot\rangle)$ with order 1. It is straightforward to check that this map gives the required isomorphism.

The proof in the case $N = \infty$ is similar.

We remark that if $\Delta^+\mathfrak{m}_k(x_1,\dots,x_k,x_0) \ne 0$ then

$$\sum_{i=1}^{k} \deg' x_i + 1 + \deg' x_0 \equiv n+2 \mod 2.$$

Therefore $\deg \Delta^+\mathfrak{m}_k \equiv n+1 \mod 2$. □

Proposition 24.11. *Let $(C,\{\mathfrak{m}_k\},\langle\cdot\rangle,e)$ be a unital and cyclic A_∞ algebra satisfying Condition 24.1. Let $\alpha \in HC^*(C)$. If $[\alpha] \in HC^*(C)/F^4HC^*(C)$ lifts to an A_3 deformation of order 1, then α lifts to an A_∞ deformation of order ∞.*

Proof. The proof is by obstruction theory, similar to, for example, [FOOO3] Lemma 7.2.74. We put $\alpha = (\alpha_k)_{k=1}^\infty$, $\alpha_k = \sum_{j=1}^\infty \theta^j \alpha_{k,j}$, $\alpha_{k,j} \in Hom(B_k^{\rm cyc}C,K)$. We are given $\alpha_{2,1}$ and will find other $\alpha_{k,j}$. The degree of $\alpha_{k,\ell}$ is $n-1$ modulo 2.

For each step we consider $Hom(B_k^{\rm cyc}C,K)$ part of

$$\sum_{k_1+k_2=k-1}\ \sum_{j_1+j_2=j} \widehat{\alpha}_{k_1,j_1} \circ \widehat{\alpha}_{k_2,j_2}$$

which is a $\delta^{\rm cyc}$ cycle of degree $\equiv n \mod 2$. Namely the obstruction to find them inductively lies in the group $F^kHC^*(C)/F^{k+1}HC^*(C)$ with $* \equiv n \mod 2$. This group vanishes by Proposition 24.3. Hence the proof. □

Remark 24.12. Two cyclic A_∞ algebras which are pseudo-isotopic are homotopy equivalent as cyclic A_∞ algebras. (This is proved in [Fu2] Theorem 8.2.) However the homotopy equivalence may not be strict in the sense of [FOOO3] Definition 3.2.29 (2). In fact, we have the following example.

Let us consider $C = \Lambda e_0 \oplus \Lambda e_1$ with $\deg e_i = i$. Here e_0 is a strict unit and $\mathfrak{m}_2(e_1,e_1) = e_0$. We put $\langle e_0,e_1\rangle = 1$ and $\mathfrak{m}_k = 0$ for $k \ge 3$.

Let $e^{i_1,\dots,i_k} \in Hom(B_kC,K)$ $(i_1,\dots,i_k \in \{0,1\})$ be the elements such that $e^{i_1,\dots,i_k}(e_{i_1}\otimes\cdots\otimes e_{i_k}) = 1$ and that $e^{i_1,\dots,i_k}$ are zero on other basis. We put

$$e^{(i_1,\dots,i_k)} = \sum_{j=1}^{k} \pm e^{i_j,\dots,i_k,i_1,\dots,i_{j-1}}.$$

Here $\pm$ is chosen so that this element is cyclically symmetric. It is easy to see that $\delta^{\rm cyc}e^{(11)} = 0$. However $e^{(11)}$ is zero in cyclic cohomology. Namely $\delta^{{\rm cyc}*}e^{(0)} = e^{(11)}$. This element $e^{(11)}$ corresponds to the deformation of the boundary operator $\mathfrak{m}_1$ to $\mathfrak{m}_1^\theta$ where $\mathfrak{m}_1^\theta(e_1) = \theta e_0$. The (nonstrict) isomorphism between $(C,\mathfrak{m})$ and $(C,\mathfrak{m}^\theta)$ is given by φ such that $\varphi_0(1) = \theta e_0$. Having nonzero $\varphi_0(1)$ corresponds to having nonzero bounding chain b. In fact if $b = \theta e_1/2$, then

$$\mathfrak{m}_1^b(e_1) = \mathfrak{m}_2(\theta e_1/2, e_1) + \mathfrak{m}_2(e_1,\theta e_1/2) = \theta e_0.$$

We also remark that in the above example we have $\delta^{{\rm cyc}*}e^{(01)} = e^{(001)} + e^{(111)}$, and $e^{(111)}$ gives a nonzero element in cyclic cohomology. This corresponds to the deformation to $\mathfrak{m}_2^\theta$ where

$$\mathfrak{m}_2^\theta(e_1,e_1) = e_0 + \theta e_0.$$

We can show nontriviality of this deformation by using the trace Z, for example.

On the other hand, the element $e^{(1111)}$ is a $\delta^{{\rm cyc}*}$ coboundary and is so zero in cyclic cohomology. (This follows from Proposition 24.3. We can also check it directly.)

We remark that the first nontrivial element of the cyclic cohomology lies in $F^0HC^*(C)/F^1HC^*(C)$. It deforms the operator $\mathfrak{m}_0(1)$, that is the critical value. (This comes from Hochschild cohomology and so is an invariant of A_∞ algebra, whether or not it carries a compatible inner product.)

We next discuss the relationship between the trace $Z(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$ and cyclic homology of $(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$. This relationship becomes most transparent in case the following condition is satisfied.

Definition 24.13. Let $(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$ be a unital and cyclic A_∞ algebra satisfying Condition 24.1.

We say $\mathbf{e}_1,\dots,\mathbf{e}_n$ is a *cyclic Clifford basis* if the following conditions are satisfied.

(1)
$$\mathbf{e}_i \cup^Q \mathbf{e}_j + \mathbf{e}_j \cup^Q \mathbf{e}_i = \begin{cases} 2d_i 1 & i=j \\ 0 & i\ne j, \end{cases}$$
where $d_i \in K\setminus 0$.

(2) $\mathbf{e}_I$, $I\in 2^{\{1,\dots,n\}}$ is a basis of C. Moreover
$$\langle \mathbf{e}_I,\mathbf{e}_J\rangle = \begin{cases} (-1)^* & J=I^c \\ 0 & \text{otherwise}. \end{cases}$$
Here $\mathbf{e}_I = \mathbf{e}_{i_1}\cup^Q\cdots\cup^Q\mathbf{e}_{i_k}$ if $I=\{i_1,\dots,i_k\}$. $*=\#\{(i,j)\in I\times J \mid j<i\}$.

The basis $\{\mathbf{e}'_i\}$ in Theorem 22.2 is a cyclic Clifford basis.

Lemma 24.14. *If $\{\mathbf{e}_1,\dots,\mathbf{e}_n\}$ is a cyclic Clifford basis of $(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$, then we have*
$$Z(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e) = 2^n d_1\cdots d_n.$$

The proof is the same as (23.1), (23.2).

Let $\{\mathfrak{m}_k^\theta\}$ be a first order A_3 deformation of $(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$ and consider
$$Z(C,\{\mathfrak{m}_k^\theta\},\langle\,\cdot\,\rangle,e)\in K.$$
We put
$$Z(C,\{\mathfrak{m}_k^\theta\},\langle\,\cdot\,\rangle,e) = Z(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)+\theta\Delta Z$$
where $\Delta Z\in K$. We have thus defined
$$\Delta Z \in Hom\left(\frac{F^3HC^*(C)}{F^4HC^*(C)},K\right). \tag{24.15}$$
Let $\{\mathfrak{m}_k^\theta\}$ be a first order A_3 deformation of $(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)$ and consider
$$Z(C,\{\mathfrak{m}_k^\theta\},\langle\,\cdot\,\rangle,e)\in K.$$
We put
$$Z(C,\{\mathfrak{m}_k^\theta\},\langle\,\cdot\,\rangle,e) = Z(C,\{\mathfrak{m}_k\},\langle\,\cdot\,\rangle,e)+\theta\Delta Z$$
where $\Delta Z\in K$. We have thus defined
$$\Delta Z \in Hom\left(\frac{F^3HC^*(C)}{F^4HC^*(C)},K\right). \tag{24.16}$$

Remark 24.15. Here we use the fact that any isomorphism between unital algebras preserves unit, automatically.

Theorem 24.16. *If b is a nondegenerate critical point of $\mathfrak{PO}_{\mathfrak{b}}$ which corresponds to $b^{\mathfrak{c}}$ by Definition 18.27, then $(H(L(u);\Lambda),\{\mathfrak{m}_k^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}\},\langle\cdot\rangle,e)$ has cyclic Clifford basis.*

Proof. We will prove that the basis $\mathbf{e}''_i$ in Proposition 23.4 is a cyclic Clifford basis up to scaler multiplication. The next proposition is the main part of its proof.

Proposition 24.17. *If* $J \ne I^c$*, then* $\langle \mathbf{e}''_I, \mathbf{e}''_J \rangle_{PD_{L(u)}} = 0$.

Proof. We prove the following lemma.

Lemma 24.18. *If* $K \subset 2^{\{1,\dots,n\}}$*,* $K \ne \{1,\dots,n\}$*, then*

$$\sum_{I,J\in 2^{\{1,\dots,n\}}} (-1)^* g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_I, \mathbf{e}''_K), \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n}) \rangle_{\mathrm{PD}_{L(u)}} = 0. \tag{24.17}$$

Here $* = n(n-1)/2$.

Proof. We calculate the left hand side of (24.17) using cyclic symmetry to obtain

$$\sum_{I,J\in 2^{\{1,\dots,n\}}} (-1)^* g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_K, \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n})), \mathbf{e}''_I \rangle_{\mathrm{PD}_{L(u)}}. \tag{24.18}$$

We do not calculate the sign here but only remark that, for given J, K, n, it depends only on the parity of $|I|$.

We put

$$\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_K, \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n})) = \sum_{S\in 2^{\{1,\dots,n\}}} (-1)^* h_S(J,K)\mathbf{e}''_S.$$

Then (24.18) is equal to

$$\sum_{I,J} (-1)^* h_S(J,K) g^{IJ} g_{SI} = \sum_J (-1)^* h_J(J,K).$$

Here we use the fact that the sum is taken for I such that the parity of $|I|$ is the same as $n - |J|$

On the other hand,

$$\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_K, \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathbf{e}''_{1,\dots,n})) = \pm c\,\mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_K, \mathbf{e}''_{J^c}) = \pm c' \mathbf{e}''_{K\ominus J^c}$$

for $c, c' \in \Lambda$. So $h_J(J,K) = 0$ for $J \ne K \ominus J^c$. $J = K \ominus J^c$ does not hold for $K \ne \{1,\dots,n\}$. The proof of Lemma 24.18 is complete. □

Corollary 24.19. $\mathbf{e}''_K \in \bigoplus_{k<n} H^k(L(u);\Lambda)$.

Proof. We put

$$\mathbf{e}''_K - c_1 \mathrm{vol}_{L(u)} \in \bigoplus_{k<n} H^k(L(u);\Lambda), \qquad \mathbf{e}''_{1,\dots,n} - c_2 \mathrm{vol}_{L(u)} \in \bigoplus_{k<n} H^k(L(u);\Lambda).$$

Here $c_2 \equiv 1 \mod \Lambda_+$ so $c_2 \ne 0$. By Corollary 19.9 the left hand side of (24.17) is

$$c_1 c_2 \sum_{I,J\in 2^{\{1,\dots,n\}}} (-1)^* g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_I, \mathrm{vol}_{L(u)}), \mathfrak{m}_2^{\mathfrak{c},\mathfrak{b},b^{\mathfrak{c}}}(\mathbf{e}''_J, \mathrm{vol}_{L(u)}) \rangle_{\mathrm{PD}_{L(u)}}.$$

By Theorem 20.2 this is nonzero unless $c_1 = 0$. Therefore the corollary follows from Lemma 24.18. □

Now we have

$$\begin{aligned} \langle \mathbf{e}''_I, \mathbf{e}''_J \rangle_{PD_{L(u)}} = \langle \mathbf{e}''_I, \mathbf{e}''_J \cup^{\mathfrak{c},Q} e \rangle_{PD_{L(u)}} &= \langle \mathbf{e}''_I, \cup^{\mathfrak{c},Q} \mathbf{e}''_J, e \rangle_{PD_{L(u)}} \\ &= c \langle \mathbf{e}''_{I\ominus J}, e \rangle_{PD_{L(u)}} = 0. \end{aligned}$$

Proposition 24.17 follows. □

Propositions 23.4 and 24.17 imply that appropriate scaler multiplication of $\mathbf{e}''_I$ gives cyclic Clifford basis. □

Theorem 24.20. *Let* $(C, \{\mathfrak{m}_k^\theta\}, \langle\cdot\rangle, e)$ *be a unital cyclic* A_∞ *algebra that admits a cyclic Clifford basis. Then the homomorphism*

$$\Delta Z : \frac{F^3HC^*(C)}{F^4HC^*(C)} \to K$$

is an isomorphism.

Proof. We consider the following deformation: We define $\mathfrak{m}_2^\theta$ such that the induced product structure is

$$\mathbf{e}_i \cup^Q \mathbf{e}_j + \mathbf{e}_j \cup^Q \mathbf{e}_i = \begin{cases} (2d_1 + \theta)1 & i = j = 1 \\ 2d_i 1 & i = j \neq 1 \\ 0 & i \neq j. \end{cases}$$

This is A_3 deformation. So it lifts to a deformation of A_∞ structure by Proposition 24.11. It gives an element of

$$X \in \frac{F^3HC^*(C)}{F^4HC^*(C)}.$$

By Lemma 24.14

$$Z(C, \{\mathfrak{m}_k^\theta\}, \langle\cdot\rangle, e) = 2^n(d_1 + \theta)d_2 \dots d_n.$$

Therefore

$$\Delta Z(X) = 2^n d_2 \dots d_n \neq 0.$$

Namely $\Delta Z \neq 0$. Since $\frac{F^3HC^*(C)}{F^4HC^*(C)} \cong K$ in our case, we have proved Theorem 24.20. □

It is likely that the components of a cyclic homology in $F_{2k}HC_*(C)/F_{2k-1}HC_*(C)$ for $k > 1$ can also be realized as a higher loop analogue of the trace Z attached to the unital and cyclic filtered A_∞ algebra. Namely we expect that a k-loop analogue of Z for unital and cyclic filtered A_∞ algebra can be constructed by taking a sum of appropriate products of the structure constants of $\mathfrak{m}_{k'}$ $(k' \leq k)$ over certain Feynman diagrams into each of whose exterior vertices the unit is inserted.

Thus the current situation looks very similar to those appearing in the contexts of

1. higher residue pairing and primitive form. [Sa].
2. S^1 equivariant cohomology and gravitational descendent. [Gi2]
3. asymptotic expansion and perturbative Chern-Simons gauge theory.

We hope to explore the relationship of our story with these in a sequel of this paper.

Remark 24.21. We however remark that our invariant Z and its higher-loop cousins are slightly different from the invariant of cyclic A_∞ algebra described in [Ko1]. In fact, in our work the unit plays a significant role as well as the A_∞ operations.

The 0-loop (or genus 0) case of the invariant of [Ko1] looks closer to the invariant defined in [Fu3], where the Maslov class is assumed to vanish.

25. Orientation

The aim of this section is to compare the orientations of the moduli spaces $\mathcal{M}_{1,1}(\beta_a)\ _{\mathrm{ev}^+}\times_{\mathrm{ev}^+}\mathcal{M}_{1,1}(\beta_b)$ and $\mathcal{M}_{3,0}(\beta_{a'})\ _{(\mathrm{ev}_0,\mathrm{ev}_1)}\times_{(\mathrm{ev}_0,\mathrm{ev}_1)}\mathcal{M}_{3,0}(\beta_{b'})$, which appear as codimension two strata of the moduli space of $\mathcal{M}^{\mathrm{ann;main}}_{(1,1);0}(\beta)$.

We recall from Lemma 20.6 that we have two points $[\Sigma_1]$ and $[\Sigma_2]$ in the moduli space of annuli $\mathcal{M}^{\mathrm{ann;\ main}}_{(1,1);0}$. Let γ be a path in $\mathcal{M}^{\mathrm{ann;main}}_{(1,1),0}$ joining $[\Sigma_1]$ and $[\Sigma_2]$. Taking the inverse image of γ under the forgetful map $\mathcal{M}^{\mathrm{ann;main}}_{(1,1),0}(\beta)\to\mathcal{M}^{\mathrm{ann;main}}_{(1,1),0}$ in (20.3), we obtain a compact space with oriented Kuranishi structure, which bounds the inverse images of $[\Sigma_1]$ and $[\Sigma_2]$. Note that the inverse images of $[\Sigma_1]$ and $[\Sigma_2]$ correspond to $\mathcal{M}_{1,1}(\beta_a)\ _{\mathrm{ev}^+}\times_{\mathrm{ev}^+}\mathcal{M}_{1,1}(\beta_b)$ and $\mathcal{M}_{3,0}(\beta_{a'})\ _{(\mathrm{ev}_0,\mathrm{ev}_1)}\times_{(\mathrm{ev}_0,\mathrm{ev}_1)}\mathcal{M}_{3,0}(\beta_{b'})$, respectively. Then our main result of this section is as follows.

Proposition 25.1. *Let γ be a path in $\mathcal{M}^{\mathrm{ann;\ main}}_{(1,1);0}$ joining $[\Sigma_1]$ and $[\Sigma_2]$. Then γ intertwines the orientations between*

$$(-1)^{\frac{n(n-1)}{2}}\bigcup_{\beta_a+\beta_b=\beta}\mathcal{M}_{1,1}(\beta_a)\ _{ev^+}\times_{ev^+}\mathcal{M}_{1,1}(\beta_b)$$

and

$$\bigcup_{\beta_{a'}+\beta_{b'}=\beta}\mathcal{M}_{3,0}(\beta_{a'})\ _{(ev_0,ev_1)}\times_{(ev_0,ev_1)}\mathcal{M}_{3,0}(\beta_{b'}).$$

The orientation of the moduli spaces appearing in Proposition 25.1 will be defined in the subsections below. To compare the orientations of these moduli spaces, we follow the argument in [FOOO3] and reduce the problem to the one for linearized operators. As in the proof of Proposition 8.1.4 in [FOOO3], we further reduce the problem to the case of product bundle pairs. We discuss this point in detail below.

25.1. Operators D_1 and D_2.

Firstly, we explain the orientation of $\mathcal{M}^{\mathrm{ann;main}}_{(1,1);0}(\beta)$.

Since L is a principal homogeneous space under the action of T^n, its tangent bundle TL is trivialized by the fundamental vector fields of the T^n-action. We fix this trivialization in this section. We reduce the problem of determining orientation to the one in the linearized problem.

Let $\rho>1$, $A=A_\rho=[1,\rho]\times S^1$ and $u:A\to X$ a smooth map with $u(\partial_i A)\subset L$ for $i=1,2$. Here we denote $\partial_1 A=\{\rho\}\times S^1$ and $\partial_2 A=\{1\}\times S^1$. Pick $p>2$ and consider the linearization of the Cauchy-Riemann equation at u:

$$D_u\overline{\partial}:W^{1,p}((A;\partial_1A,\partial_2A),u^*(TX;TL,TL))\to L^p(A;\Lambda^{0,1}A\otimes u^*TX).$$

We adopt the argument in Subsection 8.1.1 in [FOOO3] as follows. We pick a boundary parallel circles $C_1=\{\rho-\epsilon\}\times S^1\subset A, C_2=\{1+\epsilon\}\times S^1$ with a sufficiently small $\epsilon>0$. Denote by $Z\cong D^2_{(1)}\cup\mathbb{C}P^1\cup D^2_{(2)}$ the quotient space of A by collapsing C_1 and C_2 to points. Here we identify the origin $O_{(1)}$ of the disk $D^2_{(1)}$ with the "south pole" $S\in S^2\cong\mathbb{C}P^1$ and the origin $O_{(2)}$ of the disk $D^2_{(2)}$ with the "north pole" $N\in S^2\cong\mathbb{C}P^1$. We identify $\partial D^2_{(i)}=\partial_i A$. Using the trivialization of TL, we can descend the vector bundle $u^*TX\to A$ to $E\to Z$. Note that $(u|_{\partial_i A})^*TL\subset E|_{\partial_i A}$, $i=1,2$, as totally real subbundles. Then the orientation of the index $D_u\overline{\partial}$ is described as the fiber product orientation of the index problems on $D^2_{(1)}, \mathbb{C}P^1$ and $D^2_{(2)}$ as follows. We identify $E|_{D^2_{(i)}}$ with a trivial bundle $D^2_{(i)}\times\mathbb{C}^n$ such that the totally real subbundle $(u|_{\partial_i A})^*TL$ is identified with $\partial_i A\times\mathbb{R}^n$.

By deforming the operators in the space of Fredholm operators keeping the boundary condition, we can assume our the operators on $D^2_{(1)}, \mathbb{C}P^1$ and $D^2_{(2)}$ are Dolbeault operators:

$$\begin{aligned} \overline{\partial}_{(i)} = \overline{\partial} \ &: \ W_i = W^{1,p}(D^2_{(i)}, \partial D^2_{(i)}; \underline{\mathbb{C}}^n, \underline{\mathbb{R}}^n) \to L^p(D^2_{(i)}, \Lambda^{0,1} D^2_{(i)} \otimes \underline{\mathbb{C}}^n), \ i = 1,2 \\ \overline{\partial}_{\mathbb{C}P^1} = \overline{\partial} \ &: \ W_{\mathbb{C}P^1} = W^{1,p}(\mathbb{C}P^1, E|_{\mathbb{C}P^1}) \to L^p(\mathbb{C}P^1, \Lambda^{0,1}\mathbb{C}P^1 \otimes E). \end{aligned}$$

Here and hereafter we denote by $\underline{\mathbb{C}}^n$ or $\underline{\mathbb{R}}^n$, the trivial n dimensional, complex or real vector bundles, respectively. Denote by **Dom** the triple $(\xi_1, \zeta, \xi_2) \in W_1 \times W_{\mathbb{C}P^1} \times W_2$ such that

$$\xi_1(O_{(1)}) = \zeta(S) \in E_{O_{(1)}=S}, \quad \xi_2(O_{(2)}) = \zeta(N) \in E_{O_{(2)}=N}.$$

Then we find that the index of $D_u\overline{\partial}$ is identified with the index of the restriction of $\overline{\partial}_{(1)} \oplus \overline{\partial}_{\mathbb{C}P^1} \oplus \overline{\partial}_{(2)}$ to **Dom**.

We now study the orientation of $\mathcal{M}_{1,1}(\beta_a) \ {}_{\mathrm{ev}^+}\times_{\mathrm{ev}^+} \mathcal{M}_{1,1}(\beta_b)$ as a fiber product. We first give orientations on $\mathcal{M}_{1,1}(\beta_a)$ and $\mathcal{M}_{1,1}(\beta_b)$ by following the argument in Subsection 8.1.1 [FOOO3] again. That is, we collapse boundary parallel circles C_i of discs in each factor and reduce the orientation problem to the one for the Dolbeault operator on $\mathbb{C}P^1_{(i)}$ with holomorphic vector bundle $E_{(i)}$ and the one on $(D^2, \partial D^2)$ with $(\underline{\mathbb{C}}^n, \underline{\mathbb{R}}^n)$. By homotopy, we may assume that the origin $O_{(i)} \in D^2_{(i)}$ is located between C_i and $\partial D^2_{(i)}$. Since the Dolbeault operators on $\mathbb{C}P^1_{(i)}$ are complex Fredholm operators and the fiber products with their indices are taken over complex vector spaces, the orientation problem is reduced to the case that $E_{(i)} \cong D^2_{(i)} \times \mathbb{C}^n$ and the totally real subbundle over $\partial D^2_{(i)}$ are $\partial D^2_{(i)} \times \mathbb{R}^n$. Namely, it is enough to study the orientation of the index of the operator D_1, which is the restriction of $\overline{\partial}_{(1)} \oplus \overline{\partial}_{(2)}$ to

$$\mathbf{Dom}_1 = \{(\xi_1, \xi_2) \in W_1 \oplus W_2 \mid \xi_1(O_{(1)}) = \xi_2(O_{(2)})\}.$$

Next, we explain the orientation of $\mathcal{M}_{3,0}(\beta_a) \ {}_{(\mathrm{ev}_0,\mathrm{ev}_1)}\times_{(\mathrm{ev}_0,\mathrm{ev}_1)} \mathcal{M}_{3,0}(\beta_b)$. Denote by $z_0^{(i)}, z_1^{(i)}, z_2^{(i)}$ are boundary marked points on $\partial D^2_{(i)}$ respecting the counter clockwise orientation of the boundary circles. As in the previous case, the orientation problem is reduced to the orientation of the index of the operator D_2, which is the restriction of the Dolbeault operator $\overline{\partial}_{(1)} \oplus \overline{\partial}_{(2)}$ to

$$\mathbf{Dom}_2 = \{(\xi_1, \xi_2) \in W_1 \oplus W_2 \mid \xi_1(z_i^{(1)}) = \xi_2(z_i^{(2)}) \text{ for } i = 1,2\}.$$

To prove Proposition 25.1 it suffices to compare the orientations of the indices of these operators D_1 and D_2.

Notation 25.2. (1) In the rest of this section, $\langle v_1, \dots, v_n \rangle$ stands for the n dimensional real vector space spanned by v_i equipped with the orientation given by the ordering $v_1, \dots, v_n$.

(2) We fix an orientation on $\mathbb{R}^n$. (In geometric setting, we fix an orientation of the Lagrangian submanifold L.) Pick an oriented basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ of $\mathbb{R}^n$ so that $\mathbb{R}^n = \langle \mathbf{e}_1, \dots, \mathbf{e}_n \rangle$. Then $\langle \mathbf{e}_1, \sqrt{-1}\mathbf{e}_1, \dots, \mathbf{e}_n, \sqrt{-1}\mathbf{e}_n \rangle$ gives the orientation on $\mathbb{C}^n$, which will be identified with the tangent space of the ambient toric manifold X.

25.2. **Orientation of Index** D_1**.** For the bundle pair $(\underline{\mathbb{C}}^n, \underline{\mathbb{R}}^n)$, the Dolbeault operators $\overline{\partial}_{(i)}$ $(i = 1, 2)$ are surjective.

To study orientation of IndexD_1, we need to pick finite dimensional reductions of $\overline{\partial}_{(i)}$ such that the coincidence condition for values at $O_{(1)}$ and $O_{(2)}$ is transversal.

Let $A'_i = \{z \in \mathbb{C} \mid 1/4 \le |z| \le 3/4\} \subset D^2_{(i)}$. Pick and fix a decreasing smooth function $\sigma : [0, 1] \to \mathbb{R}$ such that $\sigma(r) \equiv 1$ for $r \le 1/4$ and $\sigma(r) \equiv 0$ for $r \ge 3/4$. We put

$$f_j^{(i)} = \sqrt{-1}\sigma(|z|)\mathbf{e}_j, \qquad g_j^{(i)} = \mathbf{e}_j, \qquad i = 1, 2,\ j = 1, \dots, n$$

and define $2n$-dimensional oriented real vector spaces U_i by

$$U_i = \langle f_1^{(i)}, \dots, f_n^{(i)}, g_1^{(i)}, \dots, g_n^{(i)} \rangle.$$

We define E_i by

$$E_i = \langle \nu_1^{(i)}, \dots, \nu_n^{(i)} \rangle$$

where

$$\nu_j^{(i)} = \sqrt{-1}\sigma'(|z|)(dr - \sqrt{-1}r d\theta) \otimes \mathbf{e}_j, \quad j = 1, \dots, n.$$

Here σ' is the first derivative of σ and (r, θ) is the polar coordinate so that $z = re^{\sqrt{-1}\theta}$. The restriction of the Dolbeault operator $\overline{\partial}_{(i)}$ is the following mapping $s_i : U_i \to \underline{E}_i = U_i \times E_i$ where

$$s_i(f_j^{(i)}) = (f_j^{(i)}, \nu_j^{(i)}), \quad s_i(g_j^{(i)}) = (g_j^{(i)}, 0).$$

By Convention 8.2.1 (4) in [FOOO3], the fiber product of the linearized Kuranishi models $(s_i; \underline{E}_i \to U_i)$ with respect to the evaluation maps $\mathrm{ev}^+_{(i)}$ at the points $O_{(i)} \in D^2_{(i)}$ is given by

$$(-1)^n(s = s_1 \oplus s_2; \underline{E}_1 \oplus \underline{E}_2 \to U_1 \ {}_{\mathrm{ev}^+_{(1)}}\times_{\mathrm{ev}^+_{(2)}} U_2). \tag{25.1}$$

Since the evaluation maps are given by $\mathrm{ev}^+_{(i)}(f_j^{(i)}) = \sqrt{-1}\mathbf{e}_j$ and $\mathrm{ev}^+_{(i)}(g_j^{(i)}) = \mathbf{e}_j$ in this model, we find that

$$U_1 \ {}_{\mathrm{ev}^+_{(1)}}\times_{\mathrm{ev}^+_{(2)}} U_2 = \langle \mathbf{f}_1, \dots, \mathbf{f}_n,\ \mathbf{g}_1, \dots, \mathbf{g}_n \rangle$$

as an oriented vector space. Here

$$\mathbf{f}_j = (f_j^{(1)}, f_j^{(2)}), \quad \mathbf{g}_j = (g_j^{(1)}, g_j^{(2)}), \quad j = 1, \dots, n.$$

We note that for each $i = 1, 2$ the orientation given by

$$\begin{aligned}&\langle \mathrm{ev}^+_{(i)}(f_1^{(i)}), \dots, \mathrm{ev}^+_{(i)}(f_n^{(i)}),\ \mathrm{ev}^+_{(i)}(g_1^{(i)}), \dots, \mathrm{ev}^+_{(i)}(g_n^{(i)}) \rangle \\ &= \langle \sqrt{-1}\mathbf{e}_1, \dots, \sqrt{-1}\mathbf{e}_n, \mathbf{e}_1, \dots, \mathbf{e}_n \rangle\end{aligned}$$

differs from the complex orientation of $\mathbb{C}^n$ by $(-1)^{n(n+1)/2}$. (See Notation 25.2 (2).) Here we regard $\mathbb{C}^n$ as the tangent space of the ambient toric manifold X which is the target space of the evaluation maps.

Therefore, by combining this with (25.1) and canceling $s(\mathbf{f}_j) = (\nu_j^{(1)}, \nu_j^{(2)})$ out from the front of the basis, we obtain a finite dimensional reduction of the operator D_1:

$$(-1)^{n(n-1)/2}(0; \underline{E}_2 \to U). \tag{25.2}$$

Here we define U by

$$U = \langle \mathbf{g}_1, \dots, \mathbf{g}_n \rangle.$$

We identify $E_2 \times U$ with $\underline{E_2} = E_2 \times U_2$ in an obvious way. This is the obstruction bundle of the above Kuranishi structure.

25.3. **Orientation of Index D_2.** For convenience of description, we use $\mathbb{R} \times [0,1] \subset \mathbb{C}$ instead of $D^2 \subset \mathbb{C}$. Let x, y be coordinates such that $z = x + \sqrt{-1}y$ on $\mathbb{C}$. We identify z_0 (resp. z_1) with the limit $x \to +\infty$ (resp. $x \to -\infty$). Pick and fix an increasing smooth function $\rho : \mathbb{R} \to \mathbb{R}$ such that $\rho(x) \equiv 0$ for $x \le -1$ and $\rho(x) \equiv 1$ for $x \ge 1$. We put

$$h_j^{(i)} = \rho(x)\mathbf{e}_j, \qquad k_j^{(i)} = \mathbf{e}_j, \qquad i = 1, 2,\ j = 1, \dots, n$$

and define the oriented vector spaces

$$\begin{aligned} V_i &= \langle h_1^{(i)}, \dots, h_n^{(i)},\ k_1^{(i)}, \dots, k_n^{(i)} \rangle \\ F_i &= \langle \chi_1^{(i)}, \dots, \chi_n^{(i)} \rangle, \end{aligned}$$

where

$$\chi_j^{(i)} = \rho'(x)(dx - \sqrt{-1}dy), \quad j = 1, \dots, n.$$

Then the Dolbeault operator $\overline{\partial}_{(i)}$ restricts to a mapping $t_i : V_i \to \underline{F}_i = V_i \times F_i$ which is defined by

$$t_i(h_j^{(i)}) = (h_j^{(i)}, \chi_j^{(i)}), \quad t_i(k_j^{(i)}) = (k_j^{(i)}, 0).$$

Namely $t_i : V_i \to \underline{F}_i$ is a finite dimensional reduction of $\overline{\partial}_{(i)}$.

By Convention 8.2.1 (4) in [FOOO3] again, the fiber product of the linearized Kuranishi models $(t_i; \underline{F}_i \to V_i)$ with respect to the evaluation maps $(\mathrm{ev}_0, \mathrm{ev}_1)$ at z_0 and z_1 is given by

$$(-1)^n (t = t_1 \oplus t_2; \underline{F}_1 \oplus \underline{F}_2 \to V_1 {}_{(\mathrm{ev}_0,\mathrm{ev}_1)} \times_{(\mathrm{ev}_0,\mathrm{ev}_1)} V_2). \tag{25.3}$$

Since the evaluation maps are given by $\mathrm{ev}_0(h_j^{(i)}) = \mathrm{ev}_0(k_j^{(i)}) = \mathbf{e}_j$, $\mathrm{ev}_1(h_j^{(i)}) = 0$ and $\mathrm{ev}_1(k_j^{(i)}) = \mathbf{e}_j$, we find that

$$V_1 {}_{(\mathrm{ev}_0,\mathrm{ev}_1)} \times_{(\mathrm{ev}_0,\mathrm{ev}_1)} V_2 = \langle \mathbf{h}_1, \dots, \mathbf{h}_n,\ \mathbf{k}_1, \dots, \mathbf{k}_n \rangle$$

where

$$\mathbf{h}_j = (h_j^{(1)}, h_j^{(2)}), \quad \mathbf{k}_j = (k_j^{(1)}, k_j^{(2)}), \quad j = 1, \dots, n.$$

By canceling $t(\mathbf{h}_j) = (\chi_j^{(0)}, \chi_j^{(1)})$ out from the front of the basis, we obtain a finite dimensional reduction of the operator D_2:

$$(-1)^n (0; \underline{F}_2 \to V := \langle \mathbf{k}_1, \dots, \mathbf{k}_n \rangle). \tag{25.4}$$

Here the obstruction bundle is $\underline{F}_2$, which we identify with $F_2 \times V$.

25.4. **Continuation of linear Kuranishi models.** In order to compare the orientations of the index bundles of D_1 and D_2, we will find a continuous family of linear Kuranishi models joining them.

Firstly, we use of gluing to reduce the problems to the ones on the annulus. For $\rho > 1$ we denote the annulus by $A_\rho = \{z \in \mathbb{C} \mid 1 \le |z| \le \rho\}$. We use the polar coordinates (r, θ).

25.4.1. *Linear Kuranishi model for* $\widehat{D}_1$. Gluing the two discs $D^2_{(1)}$ and $D^2_{(2)}$ by $O_{(1)} = O_{(2)}$ and smoothening the double point, we obtain the annulus $A = A_{\rho_1}$. ($\rho_1 > 1$). Here we identify

$$\partial D^2_{(1)} = \{z \in \mathbb{C} \mid |z| = \rho_1\}, \quad \partial D^2_{(2)} = \{z \in \mathbb{C} \mid |z| = 1\}.$$

(This makes the function φ introduced below increasing.) Let $\widehat{D}_1$ be the operator obtained from D_1 after this gluing. Below we describe a finite dimensional model of $\widehat{D}_1$.

Note that the elements of the obstruction bundle $\underline{E}_2$ are supported in the annular region in $D^2_{(2)}$ away from $O_{(2)}$. Therefore after we glue two discs the support of $\underline{E}_2$ becomes to a subannulus $A' = \{z \in \mathbb{C} \mid r_1 \le |z| \le r_2\}$ in A. The elements of $\underline{E}_2$ are also identified with $\mathbb{C}^n$-valued $(0,1)$-forms with support on A'. In other words, there exist $1 < r_1 < r_2 < \rho_1$ and a increasing smooth function $\varphi(r)$ with the property that $\varphi(r) \equiv 0$ for $1 \le r \le r_1$ and $\varphi(r) \equiv 1$ for $r_2 \le r \le \rho_1$ such that after we glue two discs, the vector space E_2 becomes:

$$E = \left\langle \varphi'(r) \left(\frac{\sqrt{-1}}{r} dr + d\theta \right) \otimes \mathbf{e}_1, \dots, \varphi'(r) \left(\frac{\sqrt{-1}}{r} dr + d\theta \right) \otimes \mathbf{e}_n \right\rangle.$$

Denote by $g_j : A = A_{\rho_1} \to \mathbb{C}^n$ the constant map with the value $\mathbf{e}_j$ and define an n dimensional oriented real vector space

$$\overline{V} := \langle g_1, \dots, g_n \rangle.$$

Then the linear Kuranishi model $(-1)^{n(n-1)/2}(0; \underline{E}_2 \to U)$ in (25.2) becomes the following

$$(-1)^{n(n-1)/2}(0; \underline{E} \to \overline{V}), \tag{25.5}$$

after gluing. Here $\underline{E} = E \times \overline{V}$. This is a linear Kuranishi model for $\widehat{D}_1$.

25.4.2. *Linear Kuranishi model for* $\widehat{D}_2$. Recall that we identified D^2 with $\mathbb{R} \times [0,1]$ with z_0 (resp. z_1). Here z_0 (resp. z_1) corresponds to the limit $x \to +\infty$ (resp. $x \to -\infty$). We pick $z_2 = 0 \in \mathbb{R} \times [0,1] \subset \mathbb{C}$. For a sufficiently large $R > 0$, we glue two copies of $([-R,R] \times [0,1])_{(i)}$ by identifying $(\{R\} \times [0,1])_{(1)} = (\{R\} \times [0,1])_{(2)}$ and $(\{-R\} \times [0,1])_{(1)} = (\{-R\} \times [0,1])_{(2)}$ to obtain an annulus A.

We pick an identification of A with A_{ρ_2} ($\rho_2 > 1$) such that $z_2^{(1)}$ corresponds to a point on $\partial_1 A_{\rho_2}$ and $z_2^{(2)}$ corresponds to a point on the unit circle $\partial_2 A_{\rho_2}$.

Let $\widehat{D}_2$ be the operator obtained from D_2 after this gluing. Below we describe a finite dimensional model of $\widehat{D}_2$.

Recall that elements of the obstruction bundle $\underline{F}_2$ are $\mathbb{C}^n$-valued $(0,1)$-form supported in the region $\{-1 \le x \le 1\} \subset \mathbb{R} \times [0,1]$. Hence, after the gluing, these elements correspond to $\mathbb{C}^n$-valued $(0,1)$-forms on the annulus A_{ρ_2} with support in $\{c_1 \le \theta \le c_2\} \subset A$, where $\pi/2 < c_1 < c_2 < 3\pi/2$. Moreover, there exists a decreasing smooth function $\psi : [0, 2\pi) \to \mathbb{R}$ with the property that $\psi(\theta) \equiv 1$ for $\theta \le c_1$ and $\psi(\theta) \equiv 0$ for $\theta \ge c_2$ such that F_2 becomes the space

$$F = \left\langle \psi'(\theta) \left(\frac{\sqrt{-1}}{r} dr + d\theta \right) \otimes \mathbf{e}_1, \dots, \psi'(\theta) \left(\frac{\sqrt{-1}}{r} dr + d\theta \right) \otimes \mathbf{e}_n \right\rangle$$

after gluing. (The function ψ is decreasing because of the way we identify A with A_{ρ_2}.)

Then the linear Kuranishi model $(-1)^n(0;\underline{F}_2\to V)$ described in Subsection 25.3 becomes

$$(-1)^n(0;\underline{F}\to\overline{V}=\langle g_1,\dots,g_n\rangle),\tag{25.6}$$

after gluing. Here $g_j : A = A_{\rho_2}\to\mathbb{C}^n$ is the constant map with the value $\mathbf{e}_j$ and $\underline{F}=F\times\overline{V}$. This is our linear Kuranishi model for $\widehat{D}_2$.

25.4.3. *Continuation of the linear Kuranishi models.* Let $A = A_{\rho_1} = A_{\rho_2}$. We take functions $\varphi(r)$ (resp. $\psi(\theta)$) as in Subsection 25.4.1 (resp. Subsection 25.4.2).

We consider the following families of oriented vector spaces:

$$\begin{aligned}E(\tau)=&\Big\langle(\tau+(1-\tau)\varphi'(r))\Big(\frac{\sqrt{-1}}{r}dr+d\theta\Big)\otimes\mathbf{e}_1,\\&\qquad\dots,(\tau+(1-\tau)\varphi'(r))\Big(\frac{\sqrt{-1}}{r}dr+d\theta\Big)\otimes\mathbf{e}_n\Big\rangle\end{aligned}$$

and

$$\begin{aligned}F(\tau)=&\Big\langle(-\tau+(1-\tau)\psi'(\theta))\Big(\frac{\sqrt{-1}}{r}dr+d\theta\Big)\otimes\mathbf{e}_1,\\&\qquad\dots,(-\tau+(1-\tau)\psi'(\theta))\Big(\frac{\sqrt{-1}}{r}dr+d\theta\Big)\otimes\mathbf{e}_n\Big\rangle.\end{aligned}$$

They are subspaces of $C^\infty(A;\Lambda^{0,1}\otimes\underline{\mathbb{C}}^n)$.

We remark

$$F(0)=F,\quad E(0)=E,\quad F(1)=(-1)^nE(1).\tag{25.7}$$

Note that $\varphi(r)$ (resp. $\psi(\theta)$) is an increasing (resp. decreasing) function. Therefore for any $\tau\in[0,1]$ we have

$$\begin{aligned}\tau+(1-\tau)\varphi'(r)\ge 0\\-\tau+(1-\tau)\psi'(r)\le 0,\end{aligned}\tag{25.8}$$

both of which are not identically zero.

We show the following:

Lemma 25.3. *Let $P=\overline{\partial}: W^{1,p}(A,\partial A;\mathbb{C}^n,\mathbb{R}^n)\to L^p(A;\Lambda^{0,1}A\otimes\mathbb{C}^n)$ be the Dolbeault operator on the annuls A with coefficients in the trivial bundle pair $(\underline{\mathbb{C}}^n,\underline{\mathbb{R}}^n)$. Here $p>2$. For any $\tau\in[0,1]$, we have the following:*

1. Im $\overline{\partial}\oplus E(\tau)=L^p(A;\Lambda^{0,1}A\otimes\mathbb{C}^n)$.
2. Im $\overline{\partial}\oplus F(\tau)=L^p(A;\Lambda^{0,1}A\otimes\mathbb{C}^n)$.

Proof. For the Dolbeault operator P on the annulus $(A,\partial A)$ with coefficients in the trivial bundle pair $(\underline{\mathbb{C}}^n,\underline{\mathbb{R}}^n)$, we have index $P=0$. We also find that Ker $P\cong\mathbb{R}^n$ consisting of constant sections with values in $\mathbb{R}^n$. Thus Coker P is n-dimensional. In order to prove the lemma, it is enough to show Im $\overline{\partial}\cap E(\tau)=0$ and Im $\overline{\partial}\cap F(\tau)=0$, respectively. Since the bundle pair $(\underline{\mathbb{C}}^n,\underline{\mathbb{R}}^n)$ is the direct product of n-copies of $(\underline{\mathbb{C}},\underline{\mathbb{R}})$, we consider the case that $n=1$ from now on.

(1) Suppose that there exists $f:(A,\partial A)\to(\mathbb{C},\mathbb{R})$ such that

$$\overline{\partial}f=(\tau+(1-\tau)\varphi'(r))\left(\frac{\sqrt{-1}}{r}dr+d\theta\right).$$

Pick a function $g(r):[1,\rho_1]\to\mathbb{R}$ such that $g(1)=0$ and

$$rg'(r)=\tau+(1-\tau)\varphi'(r).$$

Then we have

$$\overline{\partial}(\sqrt{-1}g(r)) = (\tau + (1-\tau)\varphi'(r))\left(\frac{\sqrt{-1}}{r}dr + d\theta\right).$$

Then $f - \sqrt{-1}g$ is a holomorphic function on A such that

$$\operatorname{Im}\,(f - \sqrt{-1}g) = \begin{cases} 0 & \text{on } \partial_1 A \\ g(\rho_1) & \text{on } \partial_2 A. \end{cases}$$

Here $\partial_1 A = \{z \in \mathbb{C} \mid |z| = \rho_1\}$, $\partial_2 A = \{z \in \mathbb{C} \mid |z| = 1\}$.

By the Schwarz reflection principle, we extend $f - \sqrt{-1}g$ to a holomorphic function on $\mathbb{C} \setminus \{0\}$, the real part of which is bounded. Since the region $\{z \in \mathbb{C} \mid \operatorname{Re} z < R\}$ for a constant $R > 0$ is conformally equivalent to the unit disc, the removable singularity theorem implies that $f - \sqrt{-1}g$ extends across the origin in $\mathbb{C}$ and it is a constant function. On the other hand, by (25.8), $\tau + (1-\tau)\varphi'(r)$ is non-negative and positive somewhere for $\tau \in [0,1]$. Thus $g(\rho_1) > 0$. This is a contradiction. Hence we have $\operatorname{Im}\overline{\partial} \cap E(\tau) = 0$.

(2) Let $\mathrm{pr} : \widetilde{A} = [1, \rho_2] \times \mathbb{R} \to A$ be the universal cover of A. By abuse of notation, we denote by (r, θ) the coordinates on $[1, \rho_2] \times \mathbb{R}$. We define $\widetilde{f}(r,\theta) = \widetilde{f}(\theta)$ by

$$\widetilde{f}(\theta) = \int_0^\theta (-\tau + (1-\tau)\psi'(\theta))d\theta,$$

which satisfies

$$\overline{\partial}\widetilde{f} = (-\tau + (1-\tau)\psi'(\theta))\left(\frac{\sqrt{-1}}{r}dr + d\theta\right).$$

Suppose that there exists $h : (A, \partial A) \to (\mathbb{C}, \mathbb{R})$ such that

$$\overline{\partial}h = (-\tau + (1-\tau)\psi'(\theta))\left(\frac{\sqrt{-1}}{r}dr + d\theta\right).$$

Then $\widetilde{k} = \widetilde{f} - h \circ \mathrm{pr} : (\widetilde{A}, \partial\widetilde{A}) \to (\mathbb{C}, \mathbb{R})$ is a holomorphic function such that

$$\widetilde{k}(r, 2\pi) - \widetilde{k}(r, 0) = \widetilde{f}(2\pi) - \widetilde{f}(0).$$

By the Schwarz reflection principle, we extend $\widetilde{k}$ to an entire holomorphic function on $\mathbb{C}$, the imaginary part of which is bounded. Hence $\widetilde{k}$ is a constant function.

On the other hand, by (25.8), $-\tau + (1-\tau)\psi'(\theta)$ is non-positive and negative somewhere for $\tau \in [0,1]$. Hence $\widetilde{f}(2\pi) - \widetilde{f}(0) \neq 0$. This is a contradiction and we have $\operatorname{Im}\overline{\partial} \cap F(\tau) = 0$. □

Lemma 25.3 and (25.7) imply that the linear Kuranishi models $(0; \underline{E} \to \overline{V})$ and $(-1)^n(0; \underline{F} \to \overline{V})$ are joined by a one parameter family of linear Kuranishi models $\{(0; \underline{E}(\tau) \to \overline{V})\}_{0 \le \tau \le 1}$ and $\{(0; \underline{F}(\tau) \to \overline{V})\}_{0 \le \tau \le 1}$ and the deformation of the complex structure on the annuli as ρ varies from ρ_1 to ρ_2. Therefore by (25.5) and (25.6) the orientations of the linear Kuranishi models of the indices of D_1 and D_2 differ by the multiplication by $(-1)^{n(n-1)/2}$. This implies Proposition 25.1. □

26. Sign in Theorem 20.1 and Proposition 21.2

Using Proposition 25.1, we check the signs in the formulas in Theorem 20.1 and Proposition 21.2. Throughout this section, space means one with oriented Kuranishi structure. (See Definition A1.17 in [FOOO3]). The dimension of a space with Kuranishi structure means the virtual dimension. We use the same notation as in [FOOO3] and follow the convention. Especially see Section 8.2 of [FOOO3]. Other than the convention of the pairing we use the same convention as [FOOO3] on orientation and sign. See Remark 26.5.

26.1. **Some lemmas.** In this subsection we prepare some general lemmas on orientation of fiber product and Poincaé duality which will be used later. Let Z be a smooth closed oriented manifold. For the later argument we consider the case either Z is a toric manifold X of dimension $2n$, or a Lagrangian submanifold L of X. Let S and T be spaces with oriented Kuranishi structures. Let $f : S \to Z$ and $g : T \to Z$ be weakly submersive maps. (See Definition A1.13 [FOOO3] for the definition of weakly submersive maps.) Then we can define the fiber product $S \times_Z T := S_f \times_g T$ over Z in the sense of Kuranishi structure. We put

$$\deg S = \dim Z - \dim S, \quad \deg T = \dim Z - \dim T.$$

By Convention 8.2.1 (4) in [FOOO3], we define the orientation on $S \times_Z T$. On the other hand, we denote by Δ the diagonal set in $Z \times Z$. Let $i : \Delta \to Z \times Z$ be an obvious inclusion map. By Remark A1.44 in [FOOO3], we can also define the fiber product $\Delta \times_{Z\times Z} (S \times T)$. Clearly $S \times_Z T = \Delta \times_{Z\times Z} (S \times T)$ as a set, but as for the orientations we have

Lemma 26.1. *$S \times_Z T = (-1)^{\dim Z \deg S} \Delta \times_{Z\times Z} (S \times T)$.*

Proof. First we assume that S, T are smooth manifolds and f_S, f_T are submersion. Put $S = S^\circ \times Z$ and $T = Z \times {}^\circ T$. See [FOOO3] Convention 8.2.1 for this notation. Then Convention 8.2.1 (3) in [FOOO3] gives the orientation on $S \times_Z T$ by $S \times_Z T = S^\circ \times Z \times {}^\circ T = (-1)^{\dim Z \deg S} Z \times S^\circ \times {}^\circ T$. On the other hand, if we put $S \times T = (Z \times Z) \times {}^\circ(S \times T)$, we find $S^\circ \times {}^\circ T = {}^\circ(S \times T)$. By identifying $\Delta = Z$ as oriented spaces, we obtain $S \times_Z T = (-1)^{\dim Z \deg S} \Delta \times_{Z\times Z} (S \times T)$. For general cases we can prove the lemma by using Convention 8.2.1 (4) in [FOOO3]. □

Next, let $f_i : S \to Z$ and $g_i : T \to Z$ $(i = 1, 2)$ be weakly submersive maps. We can define the fiber product $S \times_{Z\times Z} T := S_{(f_1,f_2)} \times_{(g_1,g_2)} T$ over $Z \times Z$. Then we can show the following.

Lemma 26.2.

$$S \times_{Z\times Z} T = (-1)^{\dim Z(\deg S+\dim T)} \big(\Delta_i \times_{f_1\times g_1} (S \times T)\big)_{f_2\times g_2} \times_i \Delta.$$

Remark 26.3. By the associative property of the fiber product (Lemma 8.2.3 in [FOOO3]), this is equivalent to

$$S \times_{Z\times Z} T = (-1)^{\dim Z(\deg S+\dim T)} \Delta_i \times_{f_1\times g_1} \big((S \times T)_{f_2\times g_2} \times_i \Delta\big).$$

If no confusion can occur, we simply denote the right hand side by

$$(-1)^{\dim Z(\deg S+\dim T)} \Delta \times_{Z\times Z} (S \times T) \times_{Z\times Z} \Delta.$$

Proof. We put $Z_i^S = Z_i^T = Z$ $(i = 1, 2)$ and

$$S = Z_1^S \times {}^{\circ}S^{\circ} \times Z_2^S = (-1)^{\dim Z \dim S}{}^{\circ}S^{\circ} \times Z_1^S \times Z_2^S$$
$$T = Z_1^T \times {}^{\circ}T^{\circ} \times Z_2^T = (-1)^{\dim Z \dim T} Z_1^T \times Z_2^T \times {}^{\circ}T^{\circ}.$$

Then by the definition of fiber product we have

$$S \times_{Z \times Z} T = (-1)^{\dim Z(\dim S + \dim T)}{}^{\circ}S^{\circ} \times Z \times Z \times {}^{\circ}T^{\circ}.$$

On the other hand, we find that

$$\begin{aligned}
&\Delta \times_{Z\times Z} (S \times T) \times_{Z \times Z} \Delta \\
&= \Delta \times_{Z \times Z} (Z_1^S \times {}^{\circ}S^{\circ} \times Z_2^S \times Z_1^T \times {}^{\circ}T^{\circ} \times Z_2^T) \times_{Z\times Z} \Delta \\
&= (-1)^{\dim Z(\deg S + \dim T)} \Delta \times_{Z\times Z} (Z_1^S \times Z_1^T \times {}^{\circ}S^{\circ} \times {}^{\circ}T^{\circ} \times Z_2^S \times Z_2^T) \times_{Z \times Z} \Delta \\
&= (-1)^{\dim Z(\deg S + \dim T)} \Delta \times {}^{\circ}S^{\circ} \times {}^{\circ}T^{\circ} \times \Delta \\
&= (-1)^{\dim Z(\deg S + \dim T)} S \times_{Z \times Z} T.
\end{aligned}$$

□

Let $f : \Delta^p \to P \subset Z$ and $g : \Delta^q \to Q \subset Z$ be smooth singular simplicies of Z. For simplicity of notation we sometimes denote them by P and Q respectively. We denote $\deg P = \dim Z - \dim P$ and $\deg Q = \dim Z - \dim Q$. In the sense of Kuranishi structure, we may assume that f and g are weakly submersive and we can take the fiber product $P \times_Z Q$ as above.

Definition 26.4. We define the Poincaré pairing in Z by

$$\langle P, Q \rangle_{PD_Z} := (-1)^{\deg P \deg Q} P \times_Z Q.$$

Remark 26.5. We note that we used a different pairing, which is denoted by $\langle \cdot, \cdot \rangle_{\text{book}}$ here, in [FOOO3] from $\langle \cdot, \cdot \rangle_{PD_Z}$. See Definition 8.4.6 in [FOOO3] for the definition of $\langle \cdot, \cdot \rangle_{\text{book}}$. The relation between $\langle \cdot, \cdot \rangle_{PD_Z}$ and $\langle \cdot, \cdot \rangle_{\text{book}}$ is given by

$$\langle P, Q \rangle_{PD_Z} = (-1)^{\deg P \deg Q} \langle P, Q \rangle_{\text{book}}.$$

This is the only point where sign convention of this paper is different from one in [FOOO3]. In fact, by Remark 8.4.7 in [FOOO3], we have

$$\langle P, Q \rangle_{PD_Z} = \# P \cap Q = \int_Z PD(P) \wedge PD(Q).$$

Here $PD(P)$ and $PD(Q)$ are differential forms (currents) which are Poincaré dual to P and Q satisfying

$$\int_P v = \int_Z PD(P) \wedge v, \quad \int_Q v = \int_Z PD(Q) \wedge v \tag{26.1}$$

for any differential form v. See Remark 3.5.8 (1) in [FOOO3].

The following lemma is an immediate consequence of Lemma 26.1.

Lemma 26.6. *We have*

$$\langle P, Q \rangle_{PD_Z} = (-1)^{\deg P \dim Q} \Delta \times_{Z \times Z} (P \times Q).$$

Next, we decompose the diagonal set Δ into sum of $\mathbf{e}_I \times \mathbf{e}_J$. (Here we note that in this section we use homological notation. That is, $\mathbf{e}_I$ denotes a chain which is the Poincaré dual via (26.1) to $\mathbf{e}_I$ used in Section 2, Section 20 – Section 23 and Section 24.) Namely we show the following.

Lemma 26.7. $\Delta = \sum_{I,J}(-1)^{|I||J|}g^{IJ}\mathbf{e}_I \times \mathbf{e}_J$, *where* $|I| = \deg \mathbf{e}_I$, $|J| = \deg \mathbf{e}_J$, $g_{IJ} = \langle \mathbf{e}_I, \mathbf{e}_J\rangle_{PD_L}$ *and* g^{IJ} *is its inverse matrix.*

Remark 26.8. This is a standard fact. See [BT] Lemma 11.22, for example. Note that the formula in [BT] is written in terms of differential forms, while ours above is written as the equality of homology classes. To translate the equality of homology classes into one of differential forms, we need to use the Poincaré duality. There are different sigh conventions of the Poincaré duality. In fact, our convention (26.1) is different form Bott-Tu's given by (6.21) in [BT]. We denote by $PD^{BT}(P)$ the Poincaré dual to P in the sense of Bott-Tu's text book. Then we can easily see that

$$PD^{BT}(P) = (-1)^{\deg P \dim P} PD(P). \tag{26.2}$$

By noticing this difference, we can see that Lemma 26.7 is the same as Bott-Tu's Lemma 11.22. (Anyway, our equality itself is an equality of homology classes. So the sign here does not depend on the choice of conventions.)

Now when $Z = L$, the Poincaré pairing $\langle \cdot, \cdot \rangle_{PD_L}$ itself does not satisfy the cyclic symmetry like (2.20). However, if we use $\overline{\mathfrak m}_2(P,Q)$ for the pairing, it satisfies this cyclic property. Using the cohomological notation, we define

$$\langle P, Q\rangle_{\rm cyc} := \int_L \overline{\mathfrak m}_2(P,Q). \tag{26.3}$$

We also extend it over Λ coefficients naturally and denote it by the same symbol. The difference between $\langle P,Q\rangle_{PD_L}$ and $\langle P,Q\rangle_{\rm cyc}$ is described by the following lemma which is nothing but Corollary 8.6.4 in [FOOO3].

Lemma 26.9. $\langle P,Q\rangle_{PD_L} = (-1)^{\deg P(\deg Q+1)}\langle P,Q\rangle_{\rm cyc}$.

We define

$$h_{IJ} := \langle \mathbf{e}_I, \mathbf{e}_J\rangle_{\rm cyc} = (-1)^{|I|(|J|+1)}\langle \mathbf{e}_I, \mathbf{e}_J\rangle_{PD_L} = (-1)^{|I|(|J|+1)} g_{IJ}, \tag{26.4}$$

with $|I| = \deg \mathbf{e}_I$, $|J| = \deg \mathbf{e}_J$. Then we have

$$h^{IJ} = (-1)^{(|I|+1)|J|} g^{IJ}. \tag{26.5}$$

Lemma 26.10.

$$\begin{aligned} &g^{IJ}\langle \overline{\mathfrak m}_2(\mathbf{e}_I, pt),\ \overline{\mathfrak m}_2(\mathbf{e}_J, pt)\rangle_{PD_L} \\ &= \sum_{I,J,A,B,C,D} (-1)^{|A||J|} h^{IJ}h^{AB}h^{C0}h^{D0}\ \langle \mathbf{e}_A \cup^{\mathfrak b,b} \mathbf{e}_I, \mathbf{e}_C\rangle_{\rm cyc}\ \langle \mathbf{e}_B \cup^{\mathfrak b,b} \mathbf{e}_J, \mathbf{e}_D\rangle_{\rm cyc}. \end{aligned}$$

Proof. We first note that the following general formula.

Sublemma 26.11.

$$\langle P,Q\rangle_{PD_L} = \sum_{A,B}(-1)^{|A||B|}g^{AB}\langle P, \mathbf{e}_A\rangle_{PD_L}\langle Q, \mathbf{e}_B\rangle_{PD_L}.$$

Proof. Put $P = \sum P^C\mathbf{e}_C$ and $Q = \sum Q^D\mathbf{e}_D$. Then we have

$$\begin{aligned} \langle P, \mathbf{e}_A\rangle_{PD_L} &= \sum P^C\langle \mathbf{e}_C, \mathbf{e}_A\rangle_{PD_L} = \sum P^C g_{CA} \\ \langle Q, \mathbf{e}_B\rangle_{PD_L} &= \sum Q^D\langle \mathbf{e}_D, \mathbf{e}_B\rangle_{PD_L} = \sum Q^D g_{DB} \\ \langle P, Q\rangle_{PD_L} &= \sum P^C Q^D\langle \mathbf{e}_C, \mathbf{e}_D\rangle_{PD_L} = \sum P^C Q^D g_{CD}. \end{aligned}$$

Therefore we have

$$\begin{aligned} g^{AB}\langle P, \mathbf{e}_A\rangle_{PD_L}\langle Q, \mathbf{e}_B\rangle_{PD_L} &= \sum g^{AB}P^C g_{CA}Q^D g_{DB} \\ &= \sum P^C Q^D g^{AB} g_{CA} g_{DB}. \end{aligned}$$

Here we note that $g_{DB} = (-1)^{|D||B|}g_{BD}$. Thus we get

$$g^{AB}g_{CA}g_{DB} = (-1)^{|D||B|}g_{CA}g^{AB}g_{BD} = (-1)^{|A||B|}g_{CD},$$

because $|A| = |D|$. Hence we obtain Sublemma 26.11. ☐

By this sublemma, we have

$$\begin{aligned} &\sum_{I,J} g^{IJ}\langle \overline{\mathfrak{m}}_2(\mathbf{e}_I, pt),\ \overline{\mathfrak{m}}_2(\mathbf{e}_J, pt)\rangle_{PD_L} \\ &= \sum_{I,J,A,B} (-1)^{|A||B|} g^{IJ} g^{AB} \langle \overline{\mathfrak{m}}_2(\mathbf{e}_I, pt),\ \mathbf{e}_A\rangle_{PD_L}\langle \overline{\mathfrak{m}}_2(\mathbf{e}_J, pt),\ \mathbf{e}_B\rangle_{PD_L}. \end{aligned}$$

Now in order to use the cyclic symmetry, we replace the pairing $\langle\cdot,\cdot\rangle_{PD_L}$ by $\langle\cdot,\cdot\rangle_{\rm cyc}$ defined by (26.3). Then by using Lemma 26.9 and (26.5), we find that the above is equal to

$$\sum_{I,J,A,B} (-1)^{|A||B|+\gamma_1} h^{IJ}h^{AB}\ \langle \overline{\mathfrak{m}}_2(\mathbf{e}_I, pt),\ \mathbf{e}_A\rangle_{\rm cyc}\ \langle \overline{\mathfrak{m}}_2(\mathbf{e}_J, pt),\ \mathbf{e}_B\rangle_{\rm cyc}.$$

Here

$$\begin{aligned} \gamma_1 &= \deg \overline{\mathfrak{m}}_2(\mathbf{e}_I, pt)(|A|+1) + \deg \overline{\mathfrak{m}}_2(\mathbf{e}_J, pt)(|B|+1) + (|I|+1)|J| + (|A|+1)|B| \\ &= (|I|+n)(|A|+1) + (|J|+n)(|B|+1) + (|I|+1)|J| + (|A|+1)|B|, \end{aligned}$$

where $n = \dim L$. We note that

$$\begin{aligned} |A|+|B| = |I|+|J| = n \\ |I|+|A| = |J|+|B| = 0. \end{aligned} \tag{26.6}$$

In particular, we have $|A| \equiv |I|$ and $|B| \equiv |J|$. Therefore it follows that

$$\gamma_1 \equiv n \qquad \mod 2. \tag{26.7}$$

Then by using the cyclic symmetry (2.20), we have

$$\sum_{I,J,A,B} (-1)^{|A||B|+\gamma_1+\gamma_2} h^{IJ}h^{AB}\ \langle \overline{\mathfrak{m}}_2(\mathbf{e}_A, \mathbf{e}_I),\ pt\rangle_{\rm cyc}\ \langle \overline{\mathfrak{m}}_2(\mathbf{e}_B, \mathbf{e}_J),\ pt\rangle_{\rm cyc}$$

with

$$\gamma_2 = (|A|+1)(|I|+n) + (|B|+1)(|J|+n). \tag{26.8}$$

Furthermore from (2.21), we obtain

$$\sum_{I,J,A,B,C,D} (-1)^{|A||B|+\gamma_1+\gamma_2+\gamma_3} h^{IJ}h^{AB}h^{C0}h^{D0}\ \langle \mathbf{e}_A \cup^{\mathfrak{b},b} \mathbf{e}_I,\ \mathbf{e}_C\rangle_{\rm cyc}\ \langle \mathbf{e}_B \cup^{\mathfrak{b},b} \mathbf{e}_J,\ \mathbf{e}_D\rangle_{\rm cyc},$$

where

$$\gamma_3 = |A|(|I|+1) + |B|(|J|+1). \tag{26.9}$$

By taking (26.6) into account, (26.7), (26.8) and (26.9) yield

$$|A||B| + \gamma_1 + \gamma_2 + \gamma_3 \equiv |A||B| \equiv |A||J|.$$

This finishes the proof of Lemma 26.10. ☐

26.2. **Proofs of signs in Theorem 20.1 and Proposition 21.2.** Let P and Q be smooth singular simplicies of L. At the final stage, we will consider the case $P = Q = pt$, a point class of L which corresponds to vol_L.

Step 1: Firstly we explore orientations on various spaces of fiber products using the evaluation maps at the boundary marked points.

We apply Lemma 26.2 to the situation that

$$\mathrm{ev}_i^{\beta_1} : \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \to L, \quad \mathrm{ev}_i^{\beta_2} : \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2) \to L, \quad (i = 0, 1).$$

Note that since the Maslov index $\mu(\beta)$ is even, we have $\dim \mathcal{M}_{3,0}^{\mathrm{main}}(\beta) = n + \mu(\beta) \equiv n$ and $\deg \mathcal{M}_{3,0}^{\mathrm{main}}(\beta) \equiv 0 \mod 2$ for any β. Thus we have

$$\begin{aligned}&\Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1)_{(\mathrm{ev}_0^{\beta_1}, \mathrm{ev}_1^{\beta_1})} \times_{(\mathrm{ev}_0^{\beta_2}, \mathrm{ev}_1^{\beta_2})} \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big)_{(\mathrm{ev}_2^{\beta_1}, \mathrm{ev}_2^{\beta_2})} \times \big(P \times Q\big) \\ &= (-1)^{\epsilon_1} \Big(\Delta \times_{L \times L} \Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big)_{L \times L} \times \Delta\Big)_{(\mathrm{ev}_2^{\beta_1}, \mathrm{ev}_2^{\beta_2})} \times \big(P \times Q\big),\end{aligned}$$

where

$$\epsilon_1 \equiv n(\deg \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) + \dim \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)) \equiv n \mod 2.$$

By the associativity (Lemma 8.2.3 in [FOOO3]) and Lemma 26.7, we have

$$\begin{aligned}&\Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1)_{(\mathrm{ev}_0^{\beta_1}, \mathrm{ev}_1^{\beta_1})} \times_{(\mathrm{ev}_0^{\beta_2}, \mathrm{ev}_1^{\beta_2})} \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big)_{(\mathrm{ev}_2^{\beta_1}, \mathrm{ev}_2^{\beta_2})} \times \big(P \times Q\big) \\ &= \sum_{I,J} (-1)^{\epsilon_1 + |I||J|} g^{IJ} \\ &\qquad \Delta \times_{L \times L} \Big(\Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big) \times_{L \times L} \big(\mathbf{e}_I \times \mathbf{e}_J\big)_{(\mathrm{ev}_2^{\beta_1}, \mathrm{ev}_2^{\beta_2})} \times \big(P \times Q\big)\Big).\end{aligned} \tag{26.10}$$

We show the following lemma.

Lemma 26.12.

$$\begin{aligned}&\Delta \times_{L \times L} \Big(\Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big) \times_{L \times L} \big(\mathbf{e}_I \times \mathbf{e}_J\big)_{(ev_2^{\beta_1}, ev_2^{\beta_2})} \times \big(P \times Q\big)\Big) \\ &= (-1)^{n + |J| \deg P} \Delta \times_{L \times L} \Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1; \mathbf{e}_I, P) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2; \mathbf{e}_J, Q)\Big).\end{aligned}$$

Proof. We write the LHS in Lemma 26.12 as

$$\Delta \times_{L \times L} \Big(\Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big) \times_{L_1 \times L_2} \big(\mathbf{e}_I \times \mathbf{e}_J\big) \times_{L_3 \times L_4} \big(P \times Q\big)\Big)$$

and put

$$\begin{aligned}&\mathbf{e}_I = L_1 \times {}^\circ\mathbf{e}_I, \quad \mathbf{e}_J = L_2 \times {}^\circ\mathbf{e}_J, \\ &P = L_3 \times {}^\circ P, \quad Q = L_4 \times {}^\circ Q, \\ &\mathcal{X} = \Delta \times_{L \times L} \Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big) \\ &\quad = \mathcal{X}^\circ \times L_1 \times L_2 \times L_3 \times L_4 = \mathcal{X}^\circ \times L_3 \times L_4 \times L_1 \times L_2.\end{aligned} \tag{26.11}$$

By the associativity again, it is equal to

$$\Big(\Delta \times_{L \times L} \Big(\mathcal{M}_{3,0}^{\mathrm{main}}(\beta_1) \times \mathcal{M}_{3,0}^{\mathrm{main}}(\beta_2)\Big)\Big) \times_{L_1 \times L_2} \big(\mathbf{e}_I \times \mathbf{e}_J\big) \times_{L_3 \times L_4} \big(P \times Q\big).$$

Using the notation introduced in (26.11), we can rewrite it as

$$\begin{aligned}&\big(\mathcal{X}^\circ \times L_3 \times L_4 \times L_1 \times L_2\big) \times_{L_1\times L_2} \big(L_1 \times {}^\circ\mathbf{e}_I \times L_2 \times {}^\circ\mathbf{e}_J\big)\\ &\qquad\qquad \times_{L_3\times L_4} \big(L_3 \times {}^\circ P \times L_4 \times {}^\circ Q\big)\\ &= (-1)^{\epsilon_2}\big(\mathcal{X}^\circ \times L_3 \times L_4 \times L_1 \times L_2\big) \times_{L_1\times L_2} \big(L_1 \times L_2 \times {}^\circ\mathbf{e}_I \times {}^\circ\mathbf{e}_J\big)\\ &\qquad\qquad \times_{L_3\times L_4} \big(L_3 \times L_4 \times {}^\circ P \times {}^\circ Q\big),\end{aligned} \tag{26.12}$$

where ϵ_2 is given by

$$\epsilon_2 = n(|I| + \deg P). \tag{26.13}$$

If we put $\mathbf{e}_I \times \mathbf{e}_J = L_1 \times L_2 \times {}^\circ(\mathbf{e}_I \times \mathbf{e}_J)$ and $P \times Q = L_3 \times L_4 \times {}^\circ(P \times Q)$, we can easily see

$$\begin{aligned}{}^\circ(\mathbf{e}_I \times \mathbf{e}_J) &= (-1)^{n|I|}\,{}^\circ\mathbf{e}_I \times {}^\circ\mathbf{e}_J,\\ {}^\circ(P \times Q) &= (-1)^{n\deg P}\,{}^\circ P \times {}^\circ Q.\end{aligned} \tag{26.14}$$

Therefore (26.12) is equal to

$$\begin{aligned}&(-1)^{\epsilon_2+\epsilon_3+\epsilon_4}\Big(\mathcal{X}^\circ \times L_3 \times L_4 \times L_1 \times L_2 \times {}^\circ(\mathbf{e}_I \times \mathbf{e}_J)\Big) \times_{L_3\times L_4} \Big(L_3 \times L_4 \times {}^\circ(P \times Q)\Big)\\ &= (-1)^{\epsilon_2+\epsilon_3+\epsilon_4}\Big(\mathcal{X}^\circ \times L_1 \times L_2 \times {}^\circ(\mathbf{e}_I \times \mathbf{e}_J) \times L_3 \times L_4\Big) \times_{L_3\times L_4} \Big(L_3 \times L_4 \times {}^\circ(P \times Q)\Big)\\ &= (-1)^{\epsilon_2+\epsilon_3+\epsilon_4}\mathcal{X}^\circ \times L_1 \times L_2 \times {}^\circ(\mathbf{e}_I \times \mathbf{e}_J) \times L_3 \times L_4 \times {}^\circ(P \times Q),\end{aligned} \tag{26.15}$$

where

$$\epsilon_3 = n|I|, \qquad \epsilon_4 = n\deg P.$$

Next we study $\mathcal{X}^\circ$. We put

$$\begin{aligned}\mathcal{M}^{\text{main}}_{3,0}(\beta_1) &= \mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times L_1 \times L_3,\\ \mathcal{M}^{\text{main}}_{3,0}(\beta_2) &= \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ \times L_2 \times L_4.\end{aligned}$$

Then we find that

$$\begin{aligned}\mathcal{X} &= \Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times L_1 \times L_3 \times \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ \times L_2 \times L_4\Big)\\ &= (-1)^n\Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ \times L_1 \times L_2 \times L_3 \times L_4\Big)\\ &= (-1)^n\Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ\Big) \times L_1 \times L_2 \times L_3 \times L_4.\end{aligned}$$

Hence we have

$$\mathcal{X}^\circ = (-1)^n\Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ\Big). \tag{26.16}$$

We substitute (26.16) into (26.15). Then the LHS in Lemma 26.12 is equal to

$$\begin{aligned}&(-1)^{\epsilon_2+\epsilon_3+\epsilon_4+n}\Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times\\ &\qquad\qquad \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ \times L_1 \times L_2 \times {}^\circ(\mathbf{e}_I \times \mathbf{e}_J) \times L_3 \times L_4 \times {}^\circ(P \times Q)\Big)\\ &= (-1)^{\epsilon_2+n+\epsilon_5}\Delta \times_{L\times L} \Big(\mathcal{M}^{\text{main}}_{3,0}(\beta_1)^\circ \times\\ &\qquad\qquad L_1 \times {}^\circ\mathbf{e}_I \times L_3 \times {}^\circ P \times \mathcal{M}^{\text{main}}_{3,0}(\beta_2)^\circ \times L_2 \times {}^\circ\mathbf{e}_J \times L_4 \times {}^\circ Q\Big).\end{aligned}$$

Here an elementary calculation shows that

$$\epsilon_5 = (n + \deg P)\dim \mathbf{e}_J. \tag{26.17}$$

By Remark 8.2.6 in [FOOO3] it is equal to the following fiber product

$$(-1)^{\epsilon_2+n+\epsilon_5}\Delta\times_{L\times L}$$
$$\left(\left(\mathcal{M}_{3,0}^{\rm main}(\beta_1)\times_{L_1\times L_3}(\mathbf{e}_I\times P)\right)\times\left(\mathcal{M}_{3,0}^{\rm main}(\beta_2)\times_{L_2\times L_4}(\mathbf{e}_J\times Q)\right)\right).$$

Then by Definition 8.4.1 in [FOOO3] we get

$$(-1)^{\epsilon_2+n+\epsilon_5+\epsilon_6}\Delta\times_{L\times L}\left(\mathcal{M}_{3,0}^{\rm main}(\beta_1;\mathbf{e}_I,P)\times\mathcal{M}_{3,0}^{\rm main}(\beta_2;\mathbf{e}_J,Q)\right),$$

where

$$\epsilon_6=(n+1)(|I|+|J|)=(n+1)n\equiv 0\mod 2. \tag{26.18}$$

Therefore, from (26.13), (26.17) and (26.18) we obtain Lemma 26.12. Here we also used (26.6) again. ☐

Next task is to rewrite the RHS in Lemma 26.12 by using the Poincaré pairing which is introduced in Definition 26.4.

Lemma 26.13.

$$\begin{aligned}&\Delta\times_{L\times L}\left(\mathcal{M}_{3,0}^{\rm main}(\beta_1;\mathbf{e}_I,P)\times\mathcal{M}_{3,0}^{\rm main}(\beta_2;\mathbf{e}_J,Q)\right)\\&=(-1)^{\epsilon_7}\langle\mathfrak{m}_{2,\beta_1}(\mathbf{e}_I,P),\ \mathfrak{m}_{2,\beta_2}(\mathbf{e}_J,Q)\rangle_{PD_L},\end{aligned}$$

where

$$\epsilon_7=(|I|+\deg P)(n+|J|+\deg Q).$$

Proof. By Lemma 26.6, $\epsilon_7=\deg\mathcal{M}_{3,0}^{\rm main}(\beta_1;\mathbf{e}_I,P)\dim\mathcal{M}_{3,0}^{\rm main}(\beta_2;\mathbf{e}_J,Q)\equiv(|I|+\deg P)(n+|J|+\deg Q)\mod 2$. ☐

An easy calculation shows that

$$\begin{aligned}\epsilon&:=\epsilon_1+|I||J|+n+|J|\deg P+\epsilon_7\\&\equiv(\deg P+|I|)(\deg Q+n)\mod 2.\end{aligned}$$

Hence combing it with (26.10), Lemma 26.12 and Lemma 26.13, we obtain the following.

Proposition 26.14.

$$\begin{aligned}&\left(\mathcal{M}_{3,0}^{\rm main}(\beta_1)_{({\rm ev}_0^{\beta_1},{\rm ev}_1^{\beta_1})}\times_{({\rm ev}_0^{\beta_2},{\rm ev}_1^{\beta_2})}\mathcal{M}_{3,0}^{\rm main}(\beta_2)\right)_{({\rm ev}_2^{\beta_1},{\rm ev}_2^{\beta_2})}\times(P\times Q)\\&=\sum_{I,J}(-1)^{\epsilon}g^{IJ}\langle\mathfrak{m}_{2,\beta_1}(\mathbf{e}_I,P),\ \mathfrak{m}_{2,\beta_2}(\mathbf{e}_J,Q)\rangle_{PD_L},\end{aligned}$$

where

$$\epsilon=(\deg P+|I|)(\deg Q+n).$$

Step 2: Next, we explore orientations on various spaces of fiber products using the evaluation maps at the interior marked points. In this case we first apply Lemma 26.1 to the situation that

$${\rm ev}_{\beta'}^+:\mathcal{M}_{1,1}(\beta')\to X,\quad {\rm ev}_{\beta''}^+:\mathcal{M}_{1,1}(\beta'')\to X.$$

Then we have

$$\begin{aligned}&\left(\mathcal{M}_{1,1}(\beta')_{{\rm ev}_{\beta'}^+}\times_{{\rm ev}_{\beta''}^+}\mathcal{M}_{1,1}(\beta'')\right)\times_{L_1\times L_2}(P\times Q)\\&=(-1)^{\delta_1}\Delta_X\times_{X\times X}\left(\mathcal{M}_{1,1}(\beta')\times\mathcal{M}_{1,1}(\beta'')\right)\times_{L_1\times L_2}(P\times Q)\end{aligned}\tag{26.19}$$

with

$$\delta_1 = \dim X \deg \mathcal{M}_{1,1}(\beta') \equiv 0 \mod 2.$$

Now we show the following.

Lemma 26.15.

$$\begin{aligned}&\Delta_X \times_{X\times X} \Big(\mathcal{M}_{1,1}(\beta') \times \mathcal{M}_{1,1}(\beta'')\Big) \times_{L_1\times L_2} (P\times Q)\\ &= \Delta_X \times_{X\times X} \Big(\mathcal{M}_{1,1}(\beta';P) \times \mathcal{M}_{1,1}(\beta'';Q)\Big).\end{aligned}$$

Proof. Put

$$\begin{aligned}\mathcal{M}_{1,1}(\beta') &= X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times L_1,\\ \mathcal{M}_{1,1}(\beta'') &= X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ \times L_2,\\ P &= L_1 \times {}^\circ P,\\ Q &= L_2 \times {}^\circ Q.\end{aligned}$$

We note that $\dim {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \equiv \dim {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ \equiv 0 \mod 2$. Then the LHS in Lemma 26.15 can be written as

$$\begin{aligned}&\Delta_X \times_{X\times X} \Big(\mathcal{M}_{1,1}(\beta') \times \mathcal{M}_{1,1}(\beta'')\Big) \times_{L_1\times L_2} (P\times Q)\\ &= \Delta_X \times_{X\times X} \Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times L_1 \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ \times L_2\Big)\\ &\qquad \times_{L_1\times L_2} \Big(L_1 \times {}^\circ P \times L_2 \times {}^\circ Q\Big)\\ &= (-1)^{\delta_2}\Delta_X \times_{X\times X} \Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ \times L_1 \times L_2\Big)\\ &\qquad \times_{L_1\times L_2} \Big(L_1 \times L_2 \times {}^\circ P \times {}^\circ Q\Big)\end{aligned}$$

where

$$\delta_2 = n \deg P.$$

It is equal to

$$\begin{aligned}&= (-1)^{\delta_2}\Delta_X \times_{X\times X} \Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ\\ &\qquad \times L_1 \times L_2 \times {}^\circ P \times {}^\circ Q)\Big)\\ &= (-1)^{\delta_2}\Delta_X \times_{X\times X} \Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times L_1 \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ\\ &\qquad \times L_2 \times {}^\circ P \times {}^\circ Q)\Big)\\ &= (-1)^{\delta_2+\delta_3}\Delta_X \times_{X\times X} \Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times L_1 \times {}^\circ P \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ\\ &\qquad \times L_2 \times {}^\circ Q)\Big),\end{aligned}$$

where $\delta_3 = n \deg P$. Since $\delta_2 + \delta_3 \equiv 0 \mod 2$ and

$$\begin{aligned}&\Big(X \times {}^\circ\mathcal{M}_{1,1}(\beta')^\circ \times L_1 \times {}^\circ P \times X \times {}^\circ\mathcal{M}_{1,1}(\beta'')^\circ \times L_2 \times {}^\circ Q)\Big)\\ &= \Big(\mathcal{M}_{1,1}(\beta') \times_{L_1} P\Big) \times \Big(\mathcal{M}_{1,1}(\beta'') \times_{L_2} Q\Big),\end{aligned}$$

we find that the LHS in Lemma 26.15 is equal to

$$\Delta_X \times_{X\times X} \left(\Big(\mathcal{M}_{1,1}(\beta') \times_{L_1} P\Big) \times \Big(\mathcal{M}_{1,1}(\beta'') \times_{L_2} Q\Big)\right).$$

Moreover, Definition 8.10.2 in [FOOO3] shows

$$\Big(\mathcal{M}_{1,1}(\beta') \times_{L_1} P\Big) \times \Big(\mathcal{M}_{1,1}(\beta'') \times_{L_2} Q\Big) = \mathcal{M}_{1,1}(\beta'; P) \times \mathcal{M}_{1,1}(\beta''; Q).$$

Thus this finishes the proof of Lemma 26.15. □

By using the Poincaré pairing on X (which is defined in Definition 26.4 generally), we find from Lemma 26.6 that

$$\begin{aligned} &\Delta_X \times_{X\times X} \Big(\mathcal{M}_{1,1}(\beta'; P) \times \mathcal{M}_{1,1}(\beta''; Q)\Big) \\ =&(-1)^{\delta_4} \langle \mathcal{M}_{1,1}(\beta'; P),\ \mathcal{M}_{1,1}(\beta''; Q)\rangle_{PD_X}, \end{aligned} \tag{26.20}$$

where

$$\delta_4 = \deg \mathcal{M}_{1,1}(\beta'; P) \dim \mathcal{M}_{1,1}(\beta''; Q) \equiv \dim P \dim Q.$$

Note that $\deg \mathcal{M}_{1,1}(\beta'; P) \equiv \dim \mathcal{M}_{1,1}(\beta'; P)$ because $\dim X$ is even. Therefore, by (26.19), Lemma 26.15 and (26.20), we obtain

Proposition 26.16.

$$\begin{aligned} &\Big(\mathcal{M}_{1,1}(\beta')_{ev^+_{\beta'}} \times_{ev^+_{\beta''}} \mathcal{M}_{1,1}(\beta'')\Big) \times_{L_1\times L_2} (P \times Q) \\ &= (-1)^{\delta} \langle \mathcal{M}_{1,1}(\beta'; P),\ \mathcal{M}_{1,1}(\beta''; Q)\rangle_{PD_X}, \end{aligned}$$

where

$$\delta = \dim P \dim Q.$$

Step 3: We suppose that $\beta_1 + \beta_2 = \beta' + \beta''$. We use Proposition 25.1 together with Proposition 26.14 and Proposition 26.16 to get the following.

Proposition 26.17.

$$\begin{aligned} &\langle \mathfrak{p}_{1,\beta'}(P),\ \mathfrak{p}_{1,\beta''}(Q)\rangle_{PD_X} \\ &= \sum_{I,J} (-1)^{\epsilon+\delta+\frac{n(n-1)}{2}} g^{IJ} \langle \mathfrak{m}_{2,\beta_1}(\mathbf{e}_I, P),\ \mathfrak{m}_{2,\beta_2}(\mathbf{e}_J, Q)\rangle_{PD_L}. \end{aligned}$$

Here

$$\epsilon = (\deg P + |I|)(\deg Q + n), \quad \delta = \dim P \dim Q,$$

which are given in Proposition 26.14 and Proposition 26.16 respectively.

Now, we consider the case

$$P = Q = pt,$$

which corresponds to vol_L. Then since $\dim P = \dim Q = 0$, we have

$$\begin{aligned} &\langle \mathfrak{p}_{1,\beta'}(pt),\ \mathfrak{p}_{1,\beta''}(pt)\rangle_{PD_X} \\ &= \sum_{I,J} (-1)^{\frac{n(n-1)}{2}} g^{IJ} \langle \mathfrak{m}_{2,\beta_1}(\mathbf{e}_I, pt),\ \mathfrak{m}_{2,\beta_2}(\mathbf{e}_J, pt)\rangle_{PD_L}. \end{aligned} \tag{26.21}$$

This finishes verification of sign in Theorem 20.1.

Combining (26.21) and Lemma 26.10, we obtain

$$
\begin{aligned}
&\langle \mathfrak{p}_{1,\beta'}(pt),\ \mathfrak{p}_{1,\beta''}(pt)\rangle_{PD_X} \\
&= \sum_{I,J,A,B,C,D} (-1)^{|A||J|+\frac{n(n-1)}{2}} h^{IJ}h^{AB}h^{C0}h^{D0} \\
&\qquad\qquad \langle \mathbf{e}_A \cup^{\mathfrak{b},b} \mathbf{e}_I,\ \mathbf{e}_C\rangle_{\rm cyc}\langle \mathbf{e}_B \cup^{\mathfrak{b},b} \mathbf{e}_J,\ \mathbf{e}_D\rangle_{\rm cyc} \qquad (26.22)\\
&= \sum_{I_1,I_2,I_3,J_1,J_2,J_3} (-1)^{|I_1||J_2|+\frac{n(n-1)}{2}} h^{I_1J_1}h^{I_2J_2}h^{I_30}h^{J_30} \\
&\qquad\qquad \langle \mathbf{e}_{I_1} \cup^{\mathfrak{b},b} \mathbf{e}_{I_2},\ \mathbf{e}_{I_3}\rangle_{\rm cyc}\langle \mathbf{e}_{J_1} \cup^{\mathfrak{b},b} \mathbf{e}_{J_2},\ \mathbf{e}_{J_3}\rangle_{\rm cyc}.
\end{aligned}
$$

Here $A = I_1, B = J_1, C = I_3, D = J_3, I = I_2$ and $J = J_2$. By Definition 2.21, the right hand side of (26.22) is nothing but $Z(\mathfrak{b}, b)$ and hence we have verified Proposition 21.2 with sign.

Remark 26.18. If we rewrite the RHS in (26.22) in terms of the Poincaré pairing $\langle \cdot,\ \cdot\rangle_{PD_L}$, we use Lemma 26.9 and (26.5) to obtain

$$
\begin{aligned}
&\langle \mathfrak{p}_{1,\beta'}(pt),\ \mathfrak{p}_{1,\beta''}(pt)\rangle_{PD_X} \\
&= \sum_{I_1,I_2,I_3,J_1,J_2,J_3} (-1)^{\zeta} g^{I_1J_1}g^{I_2J_2}g^{I_30}g^{J_30} \qquad (26.23)\\
&\qquad\qquad \langle \mathbf{e}_{I_1} \cup^{\mathfrak{b},b} \mathbf{e}_{I_2},\ \mathbf{e}_{I_3}\rangle_{PD_L}\langle \mathbf{e}_{J_1} \cup^{\mathfrak{b},b} \mathbf{e}_{J_2},\ \mathbf{e}_{J_3}\rangle_{PD_L}
\end{aligned}
$$

where

$$
\begin{aligned}
\zeta &= |I_1||J_2| + \frac{n(n-1)}{2} + (|I_1|+1)|J_1| + (|I_2|+1)|J_2| \\
&\quad + |I_3| + 1 + |J_3| + 1 + (|I_1|+|I_2|)|I_3| + (|J_1|+|J_2|)|J_3| \qquad (26.24)\\
&\equiv |I_1||J_2| + \frac{n(n-1)}{2} \qquad \mod 2
\end{aligned}
$$

by taking (26.6) into account.

Part 3. **Appendix**

27. APPENDIX 1: COINCIDENCE OF THE TWO DEFINITIONS OF $\delta^{\mathfrak{b},b}$

Let

$$b = \sum_{i=1}^{n} \mathfrak{x}_i \mathbf{e}_i \in H^1(L(u);\Lambda_0).$$

Decompose $b = b_0 + b_+$ with $b_0 \in H^1(L(u);\mathbb{C})$ and $b_+ \in H^1(L(u);\Lambda_+)$. We start with comparing the two representations

$$\begin{array}{rcl} \rho^{b_0} : H_1(L(u);\mathbb{Z}) & \to & \mathbb{C}^*; \quad \gamma \mapsto e^{\gamma\cap b_0} \\ \rho^{b} : H_1(L(u);\mathbb{Z}) & \to & \Lambda_0 \setminus \Lambda_+; \quad \gamma \mapsto e^{\gamma\cap b}. \end{array}$$

If we put $\mathfrak{y}_i = e^{\mathfrak{x}_i} \in \Lambda_0 \setminus \Lambda_+$ and $\gamma = \sum k_i \mathbf{e}_i^* \in H_1(L(u);\mathbb{Z})$, then we can write $\rho^b : H_1(L(u);\mathbb{Z}) \to \Lambda_0 \setminus \Lambda_+$ as

$$\rho^b(\gamma) = e^{\gamma\cap b} = \mathfrak{y}_1^{k_1}\dots\mathfrak{y}_n^{k_n}$$

as in Section 6. With these notations, we also have

$$\rho^b = \rho^{b_0}\rho^{b_+}.$$

Now we unravel the definitions of $\mathfrak{m}_1^{\mathfrak{b},b}$ given in Definition 6.14 and $\delta^{\mathfrak{b},b}$ in Definition 6.16. By definition, we have

$$\begin{aligned} &\mathfrak{m}_1^{\mathfrak{b},b}(h) \\ = &\sum_{l_0,\,l_1\in\mathbb{Z}_{\geq 0}} \mathfrak{m}^{\mathfrak{b},\rho}_{1+l_0+l_1;\beta}(b_+^{l_0},h,b_+^{l_1}) \\ = &\sum_{l_0,\,l_1\in\mathbb{Z}_{\geq 0}} \sum_{\beta}\sum_{\ell=0}^{\infty} T^{\beta\cap\omega/2\pi}\rho^{b_0}(\partial\beta)\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathfrak{b}^{\ell};b_+^{l_0},h,b_+^{l_1}) \\ = &\sum_{\beta}\sum_{\ell=0}^{\infty} T^{\beta\cap\omega/2\pi}\rho^{b_0}(\partial\beta)\sum_{l_0,\,l_1\in\mathbb{Z}_{\geq 0}} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathfrak{b}^{\ell};b_+^{l_0},h,b_+^{l_1}) \\ = &\sum_{\beta}\sum_{\ell=0}^{\infty} T^{\beta\cap\omega/2\pi}\rho^{b_0}(\partial\beta)\sum_{k=0}^{\infty}\sum_{l_0+l_1=k} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathfrak{b}^{\ell};b_+^{l_0},h,b_+^{l_1}). \end{aligned} \tag{27.1}$$

Here we used the definition of $\mathfrak{m}_k^{\rho}$ given in (6.12). On the other hand, $\delta^{\mathfrak{b},b}$ is defined as

$$\begin{aligned} \delta^{\mathfrak{b},b} &= \mathfrak{m}_{1,0} + \sum_{\beta,\ell}\frac{1}{\ell!}T^{\beta\cap\omega/2\pi}\rho^{b}(\partial\beta)\delta^{\mathfrak{b}}_{\beta,\ell} \\ &= \mathfrak{m}_{1,0} + \sum_{\beta,\ell}T^{\beta\cap\omega/2\pi}\rho^{b_0}(\partial\beta)\left(\frac{1}{\ell!}\rho^{b_+}(\partial\beta)\delta^{\mathfrak{b}}_{\beta,\ell}\right) \end{aligned} \tag{27.2}$$

in Definition 6.16. And by the definition (6.16) of $\delta^{\mathfrak{b}}_{\beta,\ell}$, we have

$$\delta^{\mathfrak{b}}_{\beta,\ell}(h) = (-1)^{\deg h}\mathfrak{q}_{\ell;1;\beta}(\mathfrak{b}^{\ell};h). \tag{27.3}$$

By comparing (27.1)-(27.3), the proof boils down to proving the following identity

$$\sum_{l_0+l_1=k}\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathfrak{b}^{\ell};b_+^{l_0},h,b_+^{l_1}) = \frac{(-1)^{\deg h}}{k!}\langle\partial\beta,b_+\rangle^k\mathfrak{q}_{\ell;1;\beta}(\mathfrak{b}^{\ell};h). \tag{27.4}$$

In fact, we will prove the following identity

$$\sum_{l_0+l_1=k} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1}) = \frac{(-1)^{\deg h}}{k!}\langle \partial\beta, b_+\rangle^k \mathfrak{q}_{\ell;1;\beta}(\mathbf{p}; h) \tag{27.5}$$

for all $\mathbf{p} = (\mathbf{p}(\mathbf{1}), \dots, \mathbf{p}(\ell))$ with $\mathbf{p}(i)$ being a T^n-invariant cycle in $\bigoplus_{k>0}\mathcal{A}^k(\mathbb{Z})$ from which (27.4) immediately follows.

But we can rewrite

$$\sum_{l_0+l_1=k} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1}) = \sum_{l_0=1}^{k+1} \mathfrak{q}_{\ell;k+1;\beta}(\mathbf{p}; b_+^{l_0-1}, h, b_+^{k-l_0+1}).$$

Let

$$C_k = \{(t_1, \dots, t_k) \mid 0 \le t_1 \le \cdots \le t_k \le 1\} \tag{27.6}$$

and $\widehat{C}_k$ its iterated blow up of C_k considered in [FOOO4] which we recall here now.

Let $S = \partial D$ be the boundary of the unit disc $D = D^2 \subset \mathbb{C}$ and $\beta_D \in H_2(\mathbb{C}, S)$ be the homology class of the unit disc. We consider the moduli space $\mathcal{M}_{k+1}(\mathbb{C}, S; \beta_D)$ and the evaluation map $\mathrm{ev} = (\mathrm{ev}_0, \dots, \mathrm{ev}_k) : \mathcal{M}_{k+1}(\mathbb{C}, S; \beta_D) \to (S^1)^{k+1}$. We fix a point $p_0 \in S \subset \mathbb{C}$ and put

$$\widehat{C}_k := \mathrm{ev}_0^{-1}(p_0) \subset \mathcal{M}_{k+1}(\mathbb{C}, S; \beta_D).$$

We make the identification $S^1 \setminus \{p_0\} \cong (0,1)$. Then $\vec{\mathrm{ev}}$ induces a diffeomorphism

$$\widehat{C}_k \cap \mathcal{M}_{k+1}^{\mathrm{reg}}(\mathbb{C}, S; \beta_D) \to \mathrm{Int}\ C_k$$

given by

$$[w, z_0, \dots, z_k] \mapsto (w(z_1) - w(z_0), \dots, w(z_k) - w(z_0))$$

where

$$\mathrm{Int}\ C_k = \{(t_1, \dots, t_k) \mid 0 < t_1 < \cdots < t_k < 1\} \subset C_k.$$

We identify $\partial D = S \cong \mathbb{R}/\mathbb{Z} \cong S^1$.

We choose the $\mathfrak{q}$-multisections on $\mathcal{M}^{\mathrm{main}}_{k+1,\ell}(\beta; \mathbf{p})$ that satisfies Condition 6.3. For given $\mathbf{p}$, consider the perturbed moduli spaces $\mathcal{M}^{\mathrm{main}}_{k+1,\ell}(\beta; \mathbf{p})^{\mathfrak{q}}$. Then we have

$$\mathcal{M}^{\mathrm{main}}_{k+1,\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \cong \mathcal{M}^{\mathrm{main}}_{1,\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \times \widehat{C}_k. \tag{27.7}$$

In fact, $\mathcal{M}^{\mathrm{main}}_{1,\ell}(\beta; \mathbf{p})^{\mathfrak{q}}$ consists of finitely many free T^n orbits (with multiplicity $\in \mathbb{Q}$) and $\mathcal{M}^{\mathrm{main}}_{1}(\beta; \mathbf{p})^{\mathfrak{q}} = \mathcal{M}^{\mathrm{main,reg}}_{1}(\beta; \mathbf{p})^{\mathfrak{q}}$. By Condition 6.3 (4) we have a map $\mathcal{M}^{\mathrm{main,reg}}_{k+1;\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \to \mathcal{M}^{\mathrm{main,reg}}_{1;\ell}(\beta; \mathbf{p})^{\mathfrak{q}}$. It is easy to see that the fiber can be identified with $\widehat{C}_k$.

Under this identification, Proposition 7.3 [CO] implies that the evaluation map ev is induced by

$$\mathrm{ev}_i(\mathfrak{u}; t_1, \dots, t_k) = t_i[\partial\beta] \cdot \mathrm{ev}(\mathfrak{u}) \tag{27.8}$$

for $(\mathfrak{u}; t_1, \dots, t_k) \in \mathcal{M}^{\mathrm{main}}_{1;\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \times \mathrm{Int}\ C_k \subset \mathcal{M}^{\mathrm{main}}_{1;\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \times \widehat{C}_k$. Here $\partial\beta \in H_1(L(u); \mathbb{Z}) \cong \mathbb{R}^n$, $\mathbb{R}^n \cong \mathrm{Lie}(T^n)$, is identified with an element of the universal cover $\widetilde{L}(u) \cong \mathbb{R}^n$ of $L(u) \cong T^n$ and $[t_i\partial\beta] \in L(u)$ acts as a multiplication on the torus. $\mathrm{ev}(\mathfrak{u})$ is defined by the evaluation map $\mathrm{ev} : \mathcal{M}^{\mathrm{main}}_{1;\ell}(\beta; \mathbf{p})^{\mathfrak{q}} \to L(u)$. We also have:

$$\mathrm{ev}_0(\mathfrak{u}; t_1, \dots, t_k) = \mathrm{ev}(\mathfrak{u}). \tag{27.9}$$

To prove (27.5), we take the wedge product with a T^n-invariant form g and integrate it on $L(u)$. By definition, we have

$$\begin{aligned}
&\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1}) \\
&= \sum_{I_0\in\{1,\dots,b\}^{l_0}} \sum_{I_1\in\{1,\dots,b\}^{l_1}} (x_{i_1}(b_+)\dots x_{i_{l_0}}(b_+))(x_{j_1}(b_+)\dots x_{j_{l_1}}(b_+)) \\
&\quad\times \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; \mathbf{f}_{i_1},\dots,\mathbf{f}_{i_{l_0}}, h, \mathbf{f}_{j_1},\dots,\mathbf{f}_{j_{l_1}}),
\end{aligned} \tag{27.10}$$

where $I_0 = (i_1,\dots,i_{l_0})$, $I_1 = (j_1,\dots,j_{l_1})$. We denote

$$x_{I_0}(b_+) = x_{i_1}(b_+)\cdots x_{i_{l_0}}(b_+), \quad x_{I_1}(b_+) = x_{j_1}(b_+)\cdots x_{j_{l_1}}(b_+).$$

Then we obtain

$$\begin{aligned}
&k!\int_{L(u)} g\wedge\left(\sum_{l_0+l_1=k}\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1})\right) \\
= &\int_{L(u)} g\wedge k!\sum_{I_0\in\{1,\dots,b\}^{l_0}} \sum_{I_1\in\{1,\dots,b\}^{l_1}} x_{I_0}(b_+)x_{I_1}(b_+)\times \\
&\qquad \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; \mathbf{f}_{i_1},\dots,\mathbf{f}_{i_{l_0}}, h, \mathbf{f}_{j_1},\dots,\mathbf{f}_{j_{l_1}}) \\
= &\ k!\sum_{I_0,\,I_1} x_{I_0}(b_+)x_{I_1}(b_+)\times \\
&\qquad \int_{L(u)} g\wedge\left(\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; \mathbf{f}_{i_1},\dots,\mathbf{f}_{i_{l_0}}, h, \mathbf{f}_{j_1},\dots,\mathbf{f}_{j_{k+1-l_0}})\right).
\end{aligned}$$

Now we have

$$\begin{aligned}
&\int_{L(u)} g\wedge\left(\mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; \mathbf{f}_{i_1},\dots,\mathbf{f}_{i_{l_0}}, h, \mathbf{f}_{j_1},\dots,\mathbf{f}_{j_{k+1-l_0}})\right) \\
= &\int_{\mathcal{M}^{\text{main}}_{k+1,\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \text{ev}_0^* g\wedge\Big(\text{ev}_0^*(t_1[\partial\beta])^*\mathbf{f}_{i_1}\wedge\cdots\wedge\text{ev}_0^*(t_{l_0}[\partial\beta])^*\mathbf{f}_{i_{l_0}}\wedge \\
&\text{ev}_0^*(t_{l_0}[\partial\beta])^*h\wedge\text{ev}_0^*(t_{l_0+2}[\partial\beta])^*\mathbf{f}_{j_1}\wedge\cdots\wedge\text{ev}_0^*(t_{k+1}[\partial\beta])^*\mathbf{f}_{j_{k+1-l_0}}\Big) \\
= &\int_{\mathcal{M}^{\text{main}}_{1,\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \text{ev}_0^* g\wedge\int_{\widehat{C}_{k+1}}\text{ev}_0^*(t_1[\partial\beta])^*\mathbf{f}_{i_1}\wedge\cdots\wedge\text{ev}_0^*(t_{l_0}[\partial\beta])^*\mathbf{f}_{i_{l_0}}\wedge \\
&\text{ev}_0^*(t_{l_0}[\partial\beta])^*h\wedge\text{ev}_0^*(t_{l_0+2}[\partial\beta])^*\mathbf{f}_{j_1}\wedge\cdots\wedge\text{ev}_0^*(t_{k+1}[\partial\beta])^*\mathbf{f}_{j_{k+1-l_0}}.
\end{aligned}$$

But we have

$$\begin{aligned}
&\int_{\widehat{C}_{k+1}}\text{ev}_0^*(t_1[\partial\beta])^*\mathbf{f}_{i_1}\wedge\cdots\wedge\text{ev}_0^*(t_{l_0}[\partial\beta])^*\mathbf{f}_{i_{l_0}}\wedge \\
&\text{ev}_0^*(t_{l_0+1}[\partial\beta])^*h\wedge\text{ev}_0^*(t_{l_0+2}[\partial\beta])^*\mathbf{f}_{j_1}\wedge\cdots\wedge\text{ev}_0^*(t_{k+1}[\partial\beta])^*\mathbf{f}_{j_{k+1-l_0}} \\
= &\ (-1)^{(\deg h)^{l_0}}\text{ev}_0^*\Big(\int_{C_{k+1}}(t_{l_0+1}[\partial\beta])^*h\wedge(t_1[\partial\beta])^*\mathbf{f}_{i_1}\wedge\cdots\wedge(t_{l_0}[\partial\beta])^*\mathbf{f}_{i_{l_0}}\wedge \\
&(t_{l_0+2}[\partial\beta])^*\mathbf{f}_{j_1}\wedge\cdots\wedge(t_{k+1}[\partial\beta])^*\mathbf{f}_{j_{k+1-l_0}}\Big) \\
= &\ (-1)^{(\deg h)^{l_0}}\int_{C_{k+1}}(t_{l_0+1}[\partial\beta])^*h\wedge(t_1[\partial\beta])^*\mathbf{f}_{i_1}\wedge\cdots\wedge(t_{l_0}[\partial\beta])^*\mathbf{f}_{i_{l_0}}\wedge \\
&(t_{l_0+2}[\partial\beta])^*\mathbf{f}_{j_1}\wedge\cdots\wedge(t_{k+1}[\partial\beta])^*\mathbf{f}_{j_{k+1-l_0}}
\end{aligned}$$

where the last equality follows from T^n-invariance of the integrands. By changing the variables

$$(t_{l_0+1}, t_1, \dots, t_{l_0}, \widehat{t_{l_0+1}}, t_{l_0+2}, \dots, t_{k+1}) \mapsto (s, s_1, \dots, s_k)$$

the term in the big parenthesis right above becomes

$$\begin{aligned}&(-1)^{(\deg h)^{l_0}} \int_{C_k} \int_{s_{l_0} \le s \le s_{l_0+1}} (s[\partial\beta])^* h \wedge (s_1[\partial\beta])^* \mathbf{f}_{i_1} \wedge \\ &\cdots \wedge (s_{l_0}[\partial\beta])^* \mathbf{f}_{i_{l_0}} \wedge (s_{l_0+2}[\partial\beta])^* \mathbf{f}_{j_1} \wedge \cdots \wedge (s_{k+1}[\partial\beta])^* \mathbf{f}_{j_{k+1-l_0}}.\end{aligned}$$

Therefore we obtain

$$\begin{aligned}&k! \sum_{I \in \{1,\dots,b\}^{k+1}} x_{I_0}(b_+) x_{I_1}(b_+) \times \\ &\int_{\mathcal{M}^{\text{main}}_{k+1,\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \big(\mathfrak{q}_{\ell;k+1;\beta}(\mathbf{p}; \mathbf{f}_{i_1}, \dots, \mathbf{f}_{i_{l_0}}, h, \mathbf{f}_{j_1}, \dots, \mathbf{f}_{j_{k+1-l_0}})\big) \\ = \;& \int_{\mathcal{M}_{1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \\ &\Big(\sum_{l_0=1}^{k+1} (-1)^{(\deg h)^{l_0}} k! \Big(\int_{C_k} \int_{s_{l_0} \le s \le s_{l_0+1}} (s[\partial\beta])^* h \wedge (s_1[\partial\beta])^* b_+ \wedge \cdots \wedge (s_k[\partial\beta])^* b_+ \Big) \\ = \;& \int_{\mathcal{M}_{1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \\ &\quad \Big(\sum_{l_0=1}^{k+1} (-1)^{\deg h} \int_{s_{l_0} \le s \le s_{l_0+1}} k! \int_{C_k} (s[\partial\beta])^* h \wedge (s_1[\partial\beta])^* b_+ \wedge \cdots \wedge (s_k[\partial\beta])^* b_+ \Big).\end{aligned} \tag{27.11}$$

Here we set $s_0 := 0$, $s_{k+2} := 1$.

Now it is easy to see that (27.11) becomes

$$(-1)^{\deg h} \int_{\mathcal{M}_{1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \int_0^1 (s[\partial\beta]\mathrm{ev}_0)^* h \wedge \left(k! \int_{C_k} (s_1[\partial\beta])^* b_+ \wedge \cdots \wedge (s_k[\partial\beta])^* b_+ \right).$$

On the other hand, a straightforward computation shows $(s_k[\partial\beta])^* b_+ = \langle b_+, [\partial\beta] \rangle \, ds_k$ on C^k and so

$$(s_1[\partial\beta])^* b_+ \wedge \cdots \wedge (s_k[\partial\beta])^* b_+ = \langle b_+, [\partial\beta] \rangle^k \, ds_1 \wedge ds_2 \wedge \cdots \wedge ds_k$$

thereon. Using this, we evaluate the above integral

$$\begin{aligned} &= (-1)^{\deg h} \int_{\mathcal{M}_{1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \int_0^1 (s[\partial\beta]\mathrm{ev}_0)^* h \wedge \left(k! \int_{C^k} \langle b_+, [\partial\beta] \rangle^k \, ds_1 ds_2 \dots ds_k \right) \\ &= (-1)^{\deg h} \int_{\mathcal{M}_{1;\ell}(\beta;\mathbf{p})^{\mathfrak{q}}} \mathrm{ev}_0^* g \wedge \int_0^1 (s[\partial\beta]\mathrm{ev}_0)^* h \cdot (b_+ \cap \partial\beta)^k \\ &= (-1)^{\deg h} \int_{L(u)} g \wedge \mathfrak{q}_{\ell;1;\beta}(\mathfrak{b}^{\ell}; h) \cdot (b_+ \cap \partial\beta)^k \end{aligned}$$

where we used the equality

$$k! \int_{C^k} \langle b_+, [\partial\beta] \rangle^k \, ds_1 ds_2 \dots ds_k = k! \frac{1}{k!} (b_+ \cap \partial\beta)^k = (b_+ \cap \partial\beta)^k$$

for the second equality. Therefore we have proved

$$\begin{aligned} & k! \int_{L(u)} g \wedge \left(\sum_{l_0+l_1=k} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1}) \right) \\ = \quad & (-1)^{\deg h} \int_{L(u)} g \wedge q_{\ell;1;\beta}(\mathfrak{b}^{\ell}; h) \cdot (b_+ \cap \partial\beta)^k \end{aligned}$$

for any T^n-invariant form g. Since T^n-invariant form g spans $H^*(T^n;\Lambda_0)$, this implies

$$\sum_{l_0+l_1=k} \mathfrak{q}_{\ell;1+l_0+l_1;\beta}(\mathbf{p}; b_+^{l_0}, h, b_+^{l_1}) = (-1)^{\deg h} \frac{(b_+ \cap \partial\beta)^k}{k!} q_{\ell;1;\beta}(\mathbf{p}; h).$$

This finishes the proof of (27.5) and hence the proof of the coincidence of the two definitions of $\delta^{\mathfrak{b},b}$.

28. Appendix 2: Interpolation between component-wise and T^n equivariant Kuranishi structures

In this section we clarify the points mentioned in Remarks 9.14 and 20.13. We first explain the problem in more detail. We review the notion 'component-wise Kuranishi structure' and 'disk component-wise Kuranishi structure' introduced in [Fu2] for this purpose.

We consider the moduli space $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta;\mathbf{p})$ introduced in Section 6. Here $\mathbf{p} = (P_1,\dots,P_\ell)$ and $P_i \in \mathcal{A}(X)$ are T^n invariant cycles. We consider the following two kinds of stratifications of $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta;\mathbf{p})$.

(1) The stratum of this stratification is a fiber product whose factors are

$$\mathcal{M}^{\text{main}}_{k_i+1;\ell_i}(\beta_i;\mathbf{p}_i), \qquad i=1,\dots,m$$

where $\sum k_i = k+m$, $\sum \beta_i = \beta$, $\sum \ell_i = \ell$, $\mathbf{p}_1 \cup \dots \cup \mathbf{p}_m = \mathbf{p}$ (disjoint union). Elements of the interior of this stratum have m disk components. We include the tree of sphere components to the disk component on which it is rooted.

(2) The stratum of this stratification is a fiber product whose factors are

$$\mathcal{M}^{\text{main}}_{k_i+1;\ell_i}(\beta_i;\mathbf{p}_i), \qquad i=1,\dots,m$$

or

$$\mathcal{M}_{\ell'_j}(\alpha_j;\mathbf{p}_j), \qquad j=1,\dots,m'$$

where $\sum k_i = k+m$, $\sum \beta_i + \sum \alpha_j = \beta$, $\sum \ell_i + \sum \ell'_j = \ell + m'$, $\mathbf{p}_1 \cup \dots \cup \mathbf{p}_m \cup \mathbf{p}'_1 \cup \dots \cup \mathbf{p}'_{m'} = \mathbf{p}$ (disjoint union).

We call the stratification (1) the *disk stratification* and (2) the *disk-sphere stratification.* See [FOOO3] Subsection 7.1.1 for stratification (1).

Definition 28.1. (1) We say a system of Kuranishi structures of $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta;\mathbf{p})$ *disk component-wise* if it is compatible with the fiber product description of each of the strata of disk stratification.

(2) We say a system of Kuranishi structures of $\mathcal{M}^{\text{main}}_{k+1;\ell}(\beta;\mathbf{p})$ and of $\mathcal{M}_{\ell}(\alpha;\mathbf{p})$ is *component-wise* if it is compatible with the fiber product description of each of the strata of disk-sphere stratification.

We observe that the Kuranishi structures of $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})$ which we use in this paper and in [FOOO4, FOOO5] are disk component-wise and T^n equivariant. In fact Condition 6.3 (3) implies that it is disk component-wise. (Actually this condition is stated for multisection. We need compatibility of Kuranishi structures also, since otherwise the compatibility of multisection does not make sense.) The detail of the construction of T^n equivariant Kuranishi structure is given in [FOOO4] §B.

On the other hand, the Kuranishi structures of $\mathcal{M}^{\rm main}_{k+1;\ell}(\beta;\mathbf{p})$ we use in this paper and in [FOOO4, FOOO5] is not necessarily component-wise. The reason is discussed in [FOOO4] Remark 11.4. The difficulty to make our Kuranishi structure both component-wise and T^n equivariant lies in the fact that the T^n action for the moduli space of pseudo-holomorphic *spheres* is not necessarily free though T^n action for the moduli space of pseudo-holomorphic *disks* with at least one boundary marked point is free.

In the rest of this section, we discuss how we go around this trouble and prove Lemmas 9.15.

Let $\mathbf{p}^{(\ell)} = (P_1,\dots,P_\ell)$ be as above. We consider the fiber product

$$\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p}^{(\ell)}) \ {}_{{\rm ev}^{\rm int}_3,\dots,{\rm ev}^{\rm int}_{\ell+2}}\times_{X^\ell} (P_1\times\cdots\times P_\ell). \tag{28.1}$$

The evaluation maps at the boundary marked points give a map

$${\rm ev} = ({\rm ev}_0,\dots,{\rm ev}_k) : \mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p}^{(\ell)}) \to L^{k+1} \tag{28.2}$$

and the evaluation maps at the 1st and 2nd interior marked points give:

$${\rm ev}_{\rm int} = ({\rm ev}^{\rm int}_1,{\rm ev}^{\rm int}_2) : \mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p}^{(\ell)}) \to X^2. \tag{28.3}$$

We also consider the forgetful map

$$\mathfrak{forget} : \mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p}^{(\ell)}) \to \mathcal{M}^{\rm main}_{1;2} \tag{28.4}$$

as in (9.3). We consider the point $[\Sigma_0]\in\mathcal{M}^{\rm main}_{1;2}$ as in Lemma 9.3 and take its small neighborhood U_0 in $\mathcal{M}^{\rm main}_{1;2}$. We consider

$$\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p}^{(\ell)}) \cap \mathfrak{forget}^{-1}(U_0),$$

which we write

$$\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p};U_0)$$

for simplicity. We can define the notion of disk component-wise (resp. component-wise) system of Kuranishi structures on it by modifying Definition 28.1 in an obvious way.

Its codimension one stratum is a union of fiber product (28.5) or (28.6)

$$\mathcal{M}^{\rm main}_{k_1+1;\ell_1}(\beta_1;\mathbf{p}_1) \ {}_{{\rm ev}_0}\times_{{\rm ev}_i} \mathcal{M}^{\rm main}_{k+1;\ell_2+2}(\beta;\mathbf{p}_2;U_0) \tag{28.5}$$

where $k_1+k_2=k$, $\ell_1+\ell_2=\ell$, $\mathbf{p}_1\cup\mathbf{p}_2=\mathbf{p}$ (disjoint union).

$$\mathcal{M}^{\rm main}_{k_1+1;\ell_1+2}(\beta_1;\mathbf{p}_1;U_0) \ {}_{{\rm ev}_0}\times_{{\rm ev}_i} \mathcal{M}^{\rm main}_{k+1;\ell_2}(\beta;\mathbf{p}_2) \tag{28.6}$$

where $k_1+k_2=k$, $\ell_1+\ell_2=\ell$, $\mathbf{p}_1\cup\mathbf{p}_2=\mathbf{p}$ (disjoint union).

We recall that during the construction of the operator $\mathfrak{q}$, we constructed a T^n equivariant, disk component-wise Kuranishi structures on $\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p})$. (It is not component-wise however). We call it $\mathfrak{q}$ Kuranishi structure.

The next step is to construct a Kuranishi structure on $\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta;\mathbf{p};U_0)$ which is not necessarily T^n equivariant and is compatible to the fiber product description (28.5), (28.6) of the codimension one stratum. Here we use $\mathfrak{q}$ Kuranishi structure on the first factor of (28.5) and the second factor of (28.6). Moreover our Kuranishi structure is partially component-wise. We now define this notion precisely.

Let $((\Sigma, \vec z, \vec z^+), u, (x_1, \dots, x_\ell))$ be an element of $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta; \mathbf{p}; U_0)$. (Namely $((\Sigma, \vec z, \vec z^+), u) \in \mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta)$ and $x_i \in P_i$.) We consider a tree of sphere components Σ_a of Σ, which is rooted at a point w^a contained in a disc component. We denote the corresponding disc by $D_{i(a)}$. We define $z^a \in \partial D_{i(a)}$ as follows.

1. If the 0-th boundary marked point z_0 is on $D_{i(a)}$, then $z^a = z_0$.
2. Otherwise there exists a unique singular point of $D_{i(a)}$ such that every path joining $D_{i(a)}$ with z_0 contains it. We put this singular point to be z^a.

We next define points w_i, $i = 1, 2$ as follows:

1. If the i-interior marked point z_i^+ is on a disk component then we put $w_i = z_i^+$.
2. Otherwise there is a tree of sphere components containing z_i^+ and is rooted on a disk component. We set w_i to be the point on the disk component where this tree of sphere component is rooted.

Definition 28.2. Let ϵ be a sufficiently small positive number, which we determine later and Σ_a a maximal tree of sphere components.

We say Σ_a is of *Type I* if one of the following two conditions holds:

1. $w_1 \notin D_{i(a)}$.
2. Suppose $w_1 \in D_{i(a)}$. We require $w_1 \neq w^a$. Then $(D_{i(a)}; z^a; w_1, w^a)$ defines an element of $\mathcal{M}^{\text{main}}_{1;2} \cong D^2$. We then require
$$dist([D_{i(a)}; z_a; w_1, w^a], \partial \mathcal{M}^{\text{main}}_{1;2}) < \epsilon. \tag{28.7}$$

We say that Σ_a is of *Type II* if both of the following two conditions hold:

1. $w_1 \in D_{i(a)}$.
2. Either $w_1 = w^a$ or
$$dist([D_{i(a)}; z^a; w_1, w^a], [\Sigma_0]) < \epsilon. \tag{28.8}$$

If Σ_a is neither of Type I nor of Type II, we say it is of *Type III.*

We say a sphere component is of Type I (resp. Type II, Type III) if it is contained in a maximal tree of sphere components of Type I (resp. Type II, Type III).



Figure 28.1 (Type I)



Figure 28.2 (Type II)

Lemma 28.3. *Let a sufficiently small constant* $\epsilon > 0$ *be given. Then we may choose* U_0 *sufficiently small so that whenever* $((\Sigma, \vec{z}, \vec{z}^{\rm int}), w, (x_1, \dots, x_\ell))$ *lies in* $\mathcal{M}^{\rm main}_{k+1,\ell+1}(\beta; \mathbf{p}; U_0)$ *and* Σ_a *contains one of the first two interior marked points* z_1^+, z_2^+, *then* Σ_a *is of Type II.*

Proof. If Σ_a contains z_1^+, then $w_1 \in D_{i(a)}$ and $w^a = w_1$. The lemma follows in this case.

In case Σ_a contains z_2^+, then $w_2 \in D_{i(a)}$ also. Moreover by the assumption $((\Sigma, \vec{z}, \vec{z}^{\rm int}), w, (x_1, \dots, x_\ell)) \in \mathcal{M}^{\rm main}_{k+1,\ell+1}(\beta; \mathbf{p}; U_0)$, we have either $[D_{i(a)}; z^a; w_1, w^a] = [D_{i(a)}; z^a; w_1, w_2] \in U_0$ or $w_1 = w_2$. Hence we may take ϵ small so that Σ_a is of Type II. □

Definition 28.4. A system of Kuranishi structures on $\mathcal{M}^{\rm main}_{k+1;\ell+2}(\beta; \mathbf{p}; U_0)$ is said to be *partially component-wise* if the following holds:

We consider the stratification of $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta;\mathbf{p};U_0)$ such that each of its stratum is a fiber product of

$$\mathcal{M}^{\text{main}}_{k_i+1;\ell_i}(\beta_i;\mathbf{p}_i), \qquad i=1,\dots,m, \tag{28.9}$$

or

$$\mathcal{M}_{\ell'_j}(\alpha_j;\mathbf{p}_j), \qquad j=1,\dots,m' \tag{28.10}$$

or

$$\mathcal{M}^{\text{main}}_{k_i+1;\ell_0+2}(\beta_0;\mathbf{p}_0;U_0) \tag{28.11}$$

where we have $\sum k_i = k+m$, $\sum \beta_i + \sum \alpha_j = \beta$, $\sum \ell_i + \sum \ell'_j = \ell + m'$, and

$$\mathbf{p}_0 \cup \cdots \cup \mathbf{p}_m \cup \mathbf{p}'_1 \cup \cdots \cup \mathbf{p}'_{m'} = \mathbf{p},$$

a disjoint union. For the factor (28.10), we require the corresponding sphere components are of type II. (In other words, we include a tree of sphere components of Type I or III to the disk component on which it is rooted.)

Then the Kuranishi structure is compatible to this fiber product description of the stratum of this stratification. Here we use the $\mathfrak{q}$ Kuranishi structure for the factor (28.9), and the Kuranishi structure in this definition for the factor (28.11).

Lemma 28.5. *There exists a system of Kuranishi structures on $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta;\mathbf{p};U_0)$ with the following properties:*

1. *It is compatible with the forgetful map*
$$\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta;\mathbf{p};U_0) \to \mathcal{M}^{\text{main}}_{1;\ell+2}(\beta;\mathbf{p};U_0).$$
2. *It is invariant under the action of symmetric group of order $\ell!$ which exchanges the 3rd - $(\ell+2)$nd interior marked points and exchanges $P_1,\dots,P_\ell$ at the same time.*
3. *It is compatible with the fiber product description (28.5), (28.6) of the codimension one stratum of $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta;\mathbf{p};U_0)$. Here we use the $\mathfrak{q}$ Kuranishi structure for the first factor of (28.5) and the second factor of (28.6).*
4. *It is partially component-wise in the sense of Definition 28.4.*

Proof. We remark that if a sphere component of an element of a codimension one stratum is on the first factor of (28.5) or the second factor of (28.6) then this sphere component is of type I. So we do not require the component-wise-ness for this sphere component. Once this point is understood, we can use the induction on $\beta \cap \omega$ and k, ℓ, to prove the lemma in the same way as many other cases we have already discussed. □

We call the Kuranishi structure in Lemma 28.5 the *partially component-wise Kuranishi structure* or $\mathfrak{pcw}$ Kuranishi structure.

Lemma 28.6. *There exists a system of Kuranishi structure on $[0,1]\times\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta,\mathbf{p};U_0)$ with the following properties:*

1. *It is compatible with the forgetful map*
$$[0,1]\times\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta,\mathbf{p};U_0) \to [0,1]\times\mathcal{M}^{\text{main}}_{1;\ell+2}(\beta,\mathbf{p};U_0).$$
2. *It is invariant under the action of symmetric group of order $\ell!$ which exchanges the 3rd - $(\ell+2)$nd interior marked points and exchanges $P_1,\dots,P_\ell$ at the same time.*

(3) *It is compatible with the fiber product description* (*28.12*), (*28.13*) *of the codimension one stratum of* $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$.

$$\mathcal{M}^{\text{main}}_{k_1+1;\ell_1}(\beta_1, \mathbf{p}_1) \,{}_{\text{ev}_0}\times_{\text{ev}_i} ([0,1] \times \mathcal{M}^{\text{main}}_{k+1;\ell_2+2}(\beta, \mathbf{p}_2; U_0)) \tag{28.12}$$

where $k_1 + k_2 = k$, $\ell_1 + \ell_2 = \ell$, $\mathbf{p}_1 \cup \mathbf{p}_2 = \mathbf{p}$ *(disjoint union).*

$$([0,1] \times \mathcal{M}^{\text{main}}_{k_1+1;\ell_1+2}(\beta_1, \mathbf{p}_1; U_0)) \,{}_{\text{ev}_0}\times_{\text{ev}_i} \mathcal{M}^{\text{main}}_{k+1;\ell_2}(\beta, \mathbf{p}_2) \tag{28.13}$$

where $k_1 + k_2 = k$, $\ell_1 + \ell_2 = \ell$, $\mathbf{p}_1 \cup \mathbf{p}_2 = \mathbf{p}$ *(disjoint union).*

Here we use the $\mathfrak{q}$ *Kuranishi structure for the first factor of* (*28.12*) *and the second factor of* (*28.13*).

(4) *At* $\{1\} \times \mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$ *it coincides with* $\mathfrak{pow}$ *Kuranishi structure.*

(5) *At* $\{0\} \times \mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$ *it coincides with the* $\mathfrak{q}$ *Kuranishi structure, (that is a Kuranishi structure which is* T^n *equivariant, disk component-wise, but is not component-wise).*

Proof. We can prove the lemma by the induction argument which we used many times in this paper. □

Let $\chi : U_0 \to [0,1]$ be a smooth function such that

$$\chi(\mathbf{x}) = \begin{cases} 0 & \text{if } \mathbf{x} \text{ is in a small neighborhood of } \partial U_0, \\ 1 & \text{if } \mathbf{x} \text{ is in a small neighborhood of } [\Sigma_0]. \end{cases}$$

We then embed $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$ to $[0,1] \times \mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$ by

$$\mathbf{x} \mapsto (\mathbf{x}, \chi(\mathfrak{forget}(\mathbf{x})).$$

We pull back the Kuranishi structure of Lemma 28.6 by this embedding to obtain a Kuranishi structure on $\mathcal{M}^{\text{main}}_{k+1;\ell+2}(\beta, \mathbf{p}; U_0)$. (Here pull back of the Kuranishi structure means pull back of obstruction bundle.) We claim that we can use this Kuranishi structure to prove Lemma 9.7. In fact by Lemma 28.3, we find that the partially component-wise-ness of out Kuranishi structure in a neighborhood of $\mathfrak{forget}^{-1}([\Sigma_0])$ is enough to prove the compatibility with fiber product, which we need to prove Lemma 9.7 etc. On the other hand, Properties (1), (3) of Lemma 28.6 is enough to see that contribution of the codimension one stratum vanishes. □

Thus we describe the way to go around Remark 9.14 The way to handle Remark 20.13 is the same and is left for the reader.

29. APPENDIX 3: T^n EQUIVARIANT AND CYCLICALLY SYMMETRIC KURANISHI STRUCTURE

In [FOOO4] we gave the construction of a system of T^n invariant multisections of the moduli space of pseudo-holomorphic discs. In [FOOO5] we gave the construction of a continuous family of multisections of the moduli space of pseudo-holomorphic discs. In [Fu2] construction of a continuous family of multisections with cyclic symmetry is explained. In this paper we use continuous family of multisections that is T^n invariant and cyclically symmetric. We describe it in this and the next sections. In this section, we construct a system of T^n equivariant and cyclically symmetric Kuranishi structures. Namely we prove the following

Proposition 29.1. *There exists a system of Kuranishi structures on* $\mathcal{M}_{\ell,k+1}(\beta)$ *(*$k \ge 0, \ell \ge 0$*) with the following properties.*

1. *It is compatible with the map*
$$\mathfrak{forget} : \mathcal{M}_{\ell,k+1}(\beta) \to \mathcal{M}_{\ell,1}(\beta)$$
forgetting 1-st $\dots k$*-th marked points. (Namely only the* 0*-th boundary marked point remains.) (See* [Fu2] *Section 3 for the precise meaning of this compatibility.)*
2. *It is invariant under the cyclic permutation of the boundary marked points.*
3. *It is invariant under the permutation of the interior marked points.*
4. $$\mathrm{ev}_0 : \mathcal{M}_{\ell,k+1}(\beta) \to L$$
is weakly submersive.
5. *The Kuranishi structures are disk-component-wise.*
6. *The Kuranishi structures are* T^n *equivariant.*

Proof. The proof occupies the rest of this section.

Let $(\Sigma, \vec z, \vec z^{+})$ be a marked semi-stable curve and $u : (\Sigma, \partial\Sigma) \to (X, L)$ a pseudo-holomorphic map that is stable. (Here $\vec z, \vec z^{+}$ are boundary and interior marked points, respectively.) Hereafter we write $\vec z = (z_1, \dots, z_k)$, $\vec z^{+} = (z_1^{+}, \dots, z_m^{+})$ and put $\mathbf{x} = (\Sigma, \vec z, \vec z^{+}, u)$.

Definition 29.2. We define by $G(\mathbf{x})$ the group consisting of the pairs (g, φ) of $g \in T^n$ and a biholomorphic map $\varphi : \Sigma \to \Sigma$ such that:

1. $u \circ \varphi = gu$.
2. φ preserves $\vec z^{+}$ as a set.
3. φ fixes all the points of $\vec z$.

We define a group structure on $G(\mathbf{x})$ by
$$(g_1, \varphi_1) \cdot (g_2, \varphi_2) = (g_1 g_2, \varphi_1 \circ \varphi_2). \tag{29.1}$$

Here we allow φ to permute the interior marked points $\vec z^{+}$ because we want our Kuranishi structure to be invariant under the permutation of the interior marked points. By definition, we have the projections:

$$\begin{tikzcd} & G(\mathbf{x}) \arrow[dl, "\pi_{T^n}"'] \arrow[dr, "\pi_{\mathrm{Aut}}"] & \\ T^n & & \mathrm{Aut}(\Sigma) \end{tikzcd}$$

Lemma 29.3. *The natural projection* $G(\mathbf{x}) \to \mathrm{Aut}(\Sigma)$ *is injective.*

Proof. Let $(g, \varphi) \in G(\mathbf{x})$ with $\varphi = id$. By Definition 29.2 (1), we have $gu \equiv u$. Since u has its boundary lying on a Lagrangian torus orbit L on which T^n acts freely, it follows that $g = id$. ☐

We note that each element in $G(\mathbf{x})$ induces a permutation of the elements of $\vec z^{+}$, if we regard $\vec z^{+} = (z_i^{+})_{i=1,\dots,m}$ as an ordered set.

We use it to define a $G(\mathbf{x})$-action on the set of $\mathbf{x}' = (\Sigma', \vec z', \vec z'^{+}, u')$ by
$$(g, \varphi) \cdot (\Sigma', \vec z', \vec z'^{+}, u') = (\Sigma', \vec z', \rho\vec z'^{+}, gu') \tag{29.2}$$
where ρ is defined by $\varphi(z_a^{+}) = z_{\rho(a)}^{+}$ and $\rho\vec z'^{+}$ is defined by $(\rho\vec z'^{+})_a = z'^{+}_{\rho^{-1}(a)}$.

This is a left action. Namely

$$((g_1,\varphi_1)\cdot(g_2,\varphi_2))\cdot(\Sigma',\vec{z}',\vec{z}'^{+},u') = (g_1,\varphi_1)\cdot((g_2,\varphi_2)\cdot(\Sigma',\vec{z}',\vec{z}'^{+},u')).$$

We also remark that

$$(g,\varphi)\cdot\mathbf{x} = \mathbf{x}$$

by definition.

We also define an action of $T^n \times \mathfrak{S}_\ell$ on the set of $\mathbf{x}' = (\Sigma',\vec{z}',\vec{z}'^{+},u')$ by

$$(g,\rho)\mathbf{x}' = (\Sigma',\vec{z}',\rho\vec{z}'^{+},gu').$$

The isotropy group of $\mathbf{x}$ of this action can be identified with $G(\mathbf{x})$.

We begin with the following lemma. We fix a T^n invariant metric on X.

Lemma 29.4. *Let* $\mathbf{x} = (\Sigma,\vec{z},\vec{z}^{+},u)$ *and* $G(\mathbf{x})$ *as above. We consider a relatively compact* $G(\mathbf{x})$ *invariant open subset* $K \subset \Sigma \setminus \vec{z}^{+}$ *whose closure does not contain singular points and does not intersect with* $\partial\Sigma$*. Then there exists a subspace*

$$E \subset C^\infty(\Sigma; u^*TX \otimes \Lambda^{0,1})$$

such that

1. E *is* $G(\mathbf{x})$*-invariant.*
2. *Elements of* E *have compact support in* K*.*
3. E *is finite dimensional.*
4. *We have*

$$D_u\overline{\partial}\left(\{\xi \in W^{1,p}(\Sigma,u^*TX) \mid \xi = 0 \ \text{at}\ \vec{z}\cup\vec{z}^{+}\}\right) + E = L^p(\Sigma;u^*TX\otimes\Lambda^{0,1}).$$

Proof. The existence of E_0 that satisfies (2)-(4) can be proved by the unique continuation theorem in a standard way. We will modify construction of such E_0 so that it becomes to also satisfy (1). We define an action of $G(\mathbf{x})$ on $L^2(K;u^*TX\otimes\Lambda^{0,1})$ which extends the action

$$\eta \mapsto (g,\varphi)_*(\eta) = (dg\otimes(d\varphi^{-1})^*)(\eta).$$

More specifically as a one-form on Σ, $(g,\varphi)_*(\eta)$ is defined by

$$(g,\varphi)_*(\eta)(v_x) = dg(\eta(d\varphi^{-1}(v_x))), \quad v_x \in T_x\Sigma$$

for a smooth u^*TX-valued one-form $\eta \in C^\infty(u^*TX\otimes\Lambda^{0,1})$. We recall by definition of $G(\mathbf{x})$ we have the commutative diagram

$$\begin{tikzcd}
u^*TX\otimes\Lambda^{0,1} \arrow[r, "(g,\varphi)_*"] \arrow[d] & u^*TX\otimes\Lambda^{0,1} \arrow[d] \\
\Sigma \arrow[r, "\varphi"'] & \Sigma
\end{tikzcd}$$

(Here we recall $gu\circ\varphi^{-1} = u$ from the definition of $G(\mathbf{x})$.) Using the compactness of $G(\mathbf{x})$, we can equip a $G(\mathbf{x})$-invariant inner product with $L^2(K;u^*TX\otimes\Lambda^{0,1})$. Then we have a decomposition

$$L^2(K;u^*TX\otimes\Lambda^{0,1}) = \widehat{\bigoplus_{\lambda\in\mathfrak{L}}} V_\lambda$$

where V_λ is a finite dimensional representation of $G(\mathbf{x})$ and $\mathfrak{L}$ is a certain index set, which is countable. (Here we use compactness of $G(\mathbf{x})$.) We denote by $\widehat{\bigoplus}$ the L^2 completion of the direct sum. Moreover we may choose so that each element of V_λ

is smooth. In fact, we take $G(\mathbf{x})$ invariant metrics of Σ and u^*TX and use them to define a Laplace operator on $L^2(K; u^*TX \otimes \Lambda^{0,1})$. Then each of its eigenspaces is a finite dimenisonal vector space consisting of smooth sections. Moreover it is $G(\mathbf{x})$ invariant. Therefore we may take V_λ as the eigenspaces of this Laplace operator.

Let e_i, $i = 1, \dots, N$ be an orthonormal basis of E_0. We put

$$e_i = \sum_{\lambda=1}^{\infty} e_{i,\lambda} \tag{29.3}$$

where $e_{i,\lambda} \in V_\lambda$.

When we choose e_i's, we can choose them so that the series (29.3) converges uniformly and so does in L^p. (This is well known for the eigenspaces of Laplace operator.) We put

$$e_{i,k} = \sum_{\lambda=1}^{k} e_{i,\lambda}$$

and let E_k be the vector space generated by $e_{1,k}, \dots, e_{N,k}$. If k is large, E_k satisfies (3), (4). ((4) can be proved by, for example, open mapping theorem.) We put

$$E' = \bigoplus_{\lambda=1}^{k} V_\lambda,$$

which satisfies (1), (3), (4).

To replace E' by one satisfying (2) also we proceed as follows. Let $\chi_i : X \to [0,1]$ be a sequence of $G(\mathbf{x})$ invariant smooth functions with compact support in K such that χ_i converges to 1 pointwise everywhere in K. We put

$$E(i) = \{\chi_i s \mid s \in E'\}.$$

We put $E = E(i)$ for sufficiently large i. Then (4) holds. (1)(2)(3) is immediate. □

We will use Lemma 29.4 to construct a T^n-equivariant and cyclically symmetric Kuranishi structure on $\mathcal{M}_{k+1;\ell}(\beta)$. The main part of this construction is the proof of the next proposition.

Proposition 29.5. *There exists a system of Kuranishi structures on $\mathcal{M}_{0;\ell}(\beta)$ and $\mathcal{M}_{1;\ell}(\beta)$ with the following properties:*

1. *They are forgetful map compatible in the sense of* [Fu2] *Definition 3.1.*
2. *The evaluation map $\mathrm{ev}_0 : \mathcal{M}_{1;\ell}(\beta) \to L(u)$ at the (unique) boundary marked point is weakly submersive.*
3. *The codimension one stratum of $\mathcal{M}_{0;\ell}(\beta)$ is a union of*

   $$\mathcal{M}_{1;\ell_1}(\beta_1)_{\mathrm{ev}_0} \times_{\mathrm{ev}_0} \mathcal{M}_{1;\ell_2}(\beta_2)$$

   with $\ell_1 + \ell_2 = \ell$ and $\beta_1 + \beta_2 = \beta$, $(\ell_1, \beta_1) \neq (\ell_2, \beta_2)$, and of

   $$(\mathcal{M}_{1;\ell'}(\beta')_{\mathrm{ev}_0} \times_{\mathrm{ev}_0} \mathcal{M}_{1;\ell'}(\beta'))/\mathbb{Z}_2$$

   with $2\ell' = \ell$, $2\beta' = \beta$. (Here $\mathbb{Z}_2$ acts by exchanging the factors.)

   Our system of Kuranishi structures is compatible with this fiber product description.
4. *Our Kuranishi structures are T^n equivariant in the sense of* [FOOO5] *Definition B.4.*
5. *Our Kuranishi structures are invariant under the permutation of interior marked points.*

Proof. The strategy of the proof of Proposition 29.5 is similar to the proof of [Fu2] Proposition 3.1. We however need to be a bit careful for the local construction of the Kuranishi neighborhood since we need to perform the whole construction in a T^n equivariant way. We will discuss this point in detail below.

Let $\mathbf{x} = (\Sigma, \vec{z}, \vec{z}^+, u) \in \mathcal{M}^{\text{main}}_{k;\ell}(\beta)$. Let $G = G(\mathbf{x})$ be as in Lemma 29.4. The following is a T^n equivariant version of [Fu2] Lemma 3.1.

Lemma 29.6. *There exists $E = E(\mathbf{x})$ as in Lemma 29.4 with the following properties:*

1. (1)–(4) *of Lemma 29.4 hold.*
2. *We put*

$$\mathrm{Ker}(\mathbf{x}) = (D_u\overline{\partial})^{-1}(E(\mathbf{x})). \tag{29.4}$$

Then for **any** $z_0 \in \partial\Sigma$, *the map* $Ev_{z_0} : \mathrm{Ker}(\mathbf{x}) \to T_{u(z_0)}L$ *defined by*

$$Ev_{z_0}(v) = v(z_0) \tag{29.5}$$

is surjective.

Proof. We apply Lemma 29.4 to obtain $E(\mathbf{x})$ satisfying (1). Then (2) is an immediate consequence of the fact that $\mathrm{Ker}(\mathbf{x})$ contains the Lie algebra of T^n. □

Remark 29.7. Thus the proof of Lemma 29.6 is easier than that of [Fu2] Lemma 3.1. Namely in our case (2) is automatic.

For $g \in T^n$ we put

$$E(g\mathbf{x}) = g_*E(\mathbf{x}). \tag{29.6}$$

Here $g\mathbf{x} = (\Sigma, \vec{z}^+, gu)$ and

$$g_* : C^\infty(\Sigma, u^*TX \otimes \Lambda^{0,1}) \cong C^\infty(\Sigma, (gu)^*TX \otimes \Lambda^{0,1})$$

is an obvious map. We remark that $E(g\mathbf{x})$ depends only on $g\mathbf{x} \in \mathcal{M}_{k;\ell}(\beta)$. We consider the $T^n \times \mathfrak{S}_\ell$ orbit $(T^n \times \mathfrak{S}_\ell)\mathbf{x}$ of $\mathbf{x}$. For each element $(g,\rho)\mathbf{x}$ of it we put

$$E((g,\rho)\mathbf{x}) = g_*E(\mathbf{x}) \subset C^\infty(\Sigma, (gu)^*TX \otimes \Lambda^{0,1}).$$

Lemma 29.8. *If $(g,\rho)\mathbf{x} = \mathbf{x}$ in $\mathcal{M}^{\text{main}}_{k;\ell}(\beta)$, then*

$$E((g,\rho)\mathbf{x}) \cong E(\mathbf{x}). \tag{29.7}$$

(We will explain the meaning of $\cong$ during the proof.)

Proof. $(g,\rho)\mathbf{x} = \mathbf{x}$ means that there exists a biholomorphic map $\varphi : (\Sigma, \vec{z}) \to (\Sigma, \vec{z})$ such that $\varphi(z_i^+) = z^+_{\rho^{-1}(i)}$ and $gu = u \circ \varphi$. By Lemma 29.4 we have

$$E((g,\rho)\mathbf{x}) = (g_* \otimes \varphi_*^{-1})E(\mathbf{x}).$$

This is what we mean by (29.7). □

We will use this choice $E(g\mathbf{x})$ to define a T^n equivariant Kuranishi neighborhood of the T^n orbit of $\mathbf{x}$. (This is the main part of the proof of Proposition 29.5 and occupies the most of this section.)

29.1. **The case with one disc component.** We start with the case where the following condition is satisfied.

Condition 29.9. For each $\mathfrak{t} \in Lie(T^n)$ there exists a (disk or sphere) component Σ_a of Σ and $p \in \Sigma_a$ such that u is immersed at p but $\mathfrak{t}(u(p))$ is not tangent to Σ at p. Here we also denote by $\mathfrak{t}$ the Killing vector field generated by $\mathfrak{t}$.

We remark that Condition 29.9 implies that the group $G(\mathbf{x})$ is finite.

We emphasize that $(\Sigma, \vec{z}^{+})$ may not be stable. We choose finitely many points

$$\vec{w}^{+} = (w_a^{+})_{a \in A}$$

on Σ so that the following is satisfied. We denote $m = \# A$.

Condition 29.10.

1. w_a^{+} is disjoint from $\vec{z}^{+}$, the singular point set and the boundary of Σ.
2. $(\Sigma, \vec{w}^{+})$ is stable. Moreover it has no nontrivial automorphism.
3. u is an immersion at w_a^{+}.
4. For any $\mathfrak{t} \in Lie(T^n)$ there exists w_a^{+} such that $\mathfrak{t}(u(w_a^{+}))$ is not contained in $u_* T_{w_a^{+}}\Sigma$.
5. The subset $\vec{w}^{+} \subset \Sigma$ is $G(\mathbf{x})$ invariant, i.e., $\varphi(w_a^{+}) \in \vec{w}^{+}$ for all $a \in A$, $(g, \varphi) \in G(\mathbf{x})$.

Note we can find w_a^{+} satisfying item (3)(4) above by Condition 29.9.

We consider $\mathbf{v} = (\Sigma, \vec{z}^{+}, \vec{w}^{+}) \in \mathcal{M}_{0;\ell+m}$.

Let $\mathfrak{S}_\ell$, $\mathfrak{S}_m$ be permutation groups of order $\ell!$, $m!$, respectively. The group $\mathfrak{S}_\ell$ acts on $\mathcal{M}_{0;\ell+m}$ as permutation of 1-st,…,ℓ-th marked points. The group $\mathfrak{S}_m$ acts on $\mathcal{M}_{0;\ell+m}$ as permutation of $(\ell+1)$-st,…,$(\ell+m)$-th marked points.

Definition 29.11. We define $\mathrm{Aut}_+(\mathbf{v})$ as the set of all $\varphi : \Sigma \to \Sigma$ such that:

1. φ is biholomorphic.
2. There exists $\rho = (\rho_1, \rho_2) \in \mathfrak{S}_\ell \times \mathfrak{S}_m$ such that
$$\varphi(z_i^{+}) = z_{\rho_1(i)}^{+}, \quad \varphi(w_i^{+}) = w_{\rho_2(i)}^{+}.$$

By Condition 29.10 (2), $\varphi \mapsto \rho_2$ as above is an injective homomorphism :

$$\Psi^2 : \mathrm{Aut}_+(\mathbf{v}) \to \mathfrak{S}_m.$$

We also write the homomorphism $\varphi \mapsto \rho_1$ as

$$\Psi^1 : \mathrm{Aut}_+(\mathbf{v}) \to \mathfrak{S}_\ell.$$

We put $\Psi = (\Psi_1, \Psi_2)$.

Let $\rho = (\rho_1, \rho_2) \in \mathfrak{S}_\ell \times \mathfrak{S}_m$. Then $\rho\mathbf{v} = \mathbf{v}$ if and only if ρ is in the image of Ψ.

Condition 29.10 (2) implies that $\Psi : \mathrm{Aut}_+(\mathbf{v}) \to \mathfrak{S}_\ell \times \mathfrak{S}_m$ is injective. Using this fact we can define an action of $\mathrm{Aut}_+(\mathbf{v})$ on a neighborhood of $\mathbf{v}$ in $\mathcal{M}_{0;\ell+m}$ as follows: Let $\mathbf{v}' = (\Sigma, \vec{z}'^{+}, \vec{w}'^{+})$ be in a neighborhood of $\mathbf{v}$ in $\mathcal{M}_{0;\ell+m}$ and $\varphi \in \mathrm{Aut}_+(\mathbf{v})$. We put $\Psi(\varphi) = (\rho_1, \rho_2)$. We then define

$$\varphi \cdot \mathbf{v}' = (\Sigma, \rho_1 \vec{z}'^{+}, \rho_2 \vec{w}'^{+}), \tag{29.8}$$

where

$$(\rho_1 \vec{z}'^{+})_a = z'^{+}_{\rho_1^{-1}(a)}, \quad (\rho_2 \vec{w}'^{+})_a = w'^{+}_{\rho_2^{-1}(a)}.$$

In other words, the action changes the enumeration of the interior marked points in the same way as φ does on $\mathbf{v}$. This action induces an action of $G(\mathbf{x})$ on a neighborhood of $\mathbf{v}$ in $\mathcal{M}_{0;\ell+m}$. Namely

$$(g,\varphi)\cdot\mathbf{v}' = \varphi\cdot\mathbf{v}'.$$

(See (29.2).)

We take neighborhoods $\mathfrak{U}(\mathbf{v})$ of $\mathbf{v}$ in $\mathcal{M}_{0;\ell+m}$ with the following properties. There exists a neighborhood $\mathcal{U}(S(\Sigma))$ of the singular point set of Σ such that for any $\mathbf{v}' = (\Sigma', \vec{z}'^{+}, \vec{w}^{+}) \in \mathfrak{U}(\mathbf{v})$ we have a smooth embedding

$$i_{\mathbf{v}'} : \Sigma \setminus \mathcal{U}(S(\Sigma)) \to \Sigma' \tag{29.9}$$

such that

$$i_{\mathbf{v}'}(z_i^+) = z_i^+, \quad i_{\mathbf{v}'}(w_i^+) = w_i^+. \tag{29.10}$$

We may assume that $i_{\mathbf{v}'}$ is holomorphic on the support of $E(\mathbf{x})$. (See [FO] §12. The map in [FO] (12.10) is a version of our map $i_{\mathbf{v}'}$ for the case $\partial\Sigma = \emptyset$.)

Furthermore we assume the following condition (29.11).

Let $\mathbf{v}' = (\Sigma', \vec{z}'^{+}, \vec{w}'^{+}) \in \mathfrak{U}(\mathbf{v})$, and $\varphi \in \mathrm{Aut}_+(\mathbf{v})$. We put

$$\sigma = (\sigma_1, \sigma_2) = \Psi(\varphi)$$

and require

$$i_{\sigma\mathbf{v}'} = i_{\mathbf{v}'} \circ \varphi^{-1} : \Sigma \setminus \mathcal{U}(S(\Sigma)) \to \Sigma'. \tag{29.11}$$

Remark 29.12. We remark

$$(i_{\mathbf{v}'} \circ \varphi^{-1})(z_i^+) = i_{\mathbf{v}'}(z^+_{\sigma_1^{-1}(i)}) = i_{\sigma\mathbf{v}'}(z_i^+).$$

We also choose $i_{\mathbf{v}'}$ so that the identity $i_{\mathbf{v}'_{(2)};\mathbf{v}_{(2)}} = i_{\mathbf{v}';\mathbf{v}}$ holds when we change the enumeration of interior marked points of $\vec{z}^{+}, \vec{w}^{+}$ and of $\vec{z}'^{+}, \vec{w}'^{+}$ in the same way as we obtain $\mathbf{v}_{(2)}, \mathbf{v}'_{(2)}$. (Here we write $i_{\mathbf{v}';\mathbf{v}}$ etc. in place of $i_{\mathbf{v}'}$ to specify $\mathbf{v}$.)

We also require that $i_{\mathbf{v}'}$ depends continuously on $\mathbf{v}'$ in C^∞ sense.

We can prove the existence of such $i_{\mathbf{v}'}$ using the compactness of $G(\mathbf{x})$.

We fix $\mathbf{x} = (\Sigma, \vec{z}^{+}, u)$ and $\mathbf{v} = (\Sigma, \vec{z}^{+}, \vec{w}^{+})$. For each w_a^+ we choose a normal slice $N_{w_a^+}$ of u such that:

Condition 29.13. (1) $N_{w_a^+}$ is a smooth submanifold of codimension 2.

(2) $N_{w_a^+}$ intersects transversally with $u(\Sigma)$ at $u(w_a^+)$.

(3) If $(g,\varphi) \in G(\mathbf{x})$, then $gN_{w_a^+} = N_{\varphi(w_a^+)}$.

We remark that the choices of the above slices $N_{w_a^+}$'s depend only on $\mathbf{x} = (\Sigma, \vec{z}^{+}, u)$ and $\vec{w}^{+} = \{w_a^+\}_{a\in A}$ which will be *fixed once and for all* when the latter is given. They also carry the metrics induced from the one given in the ambient space X.

We next consider $\mathbf{x}' = (\Sigma', \vec{z}'^{+}, u')$ and the following condition on it. Let ϵ_1, ϵ_2 be positive real numbers.

Definition 29.14. Let $h \in T^n$. We say that $\mathbf{x}'$ is (ϵ_1,ϵ_2)-*close to* $h\mathbf{x}$ if there exists $\vec{w}'^{+}$ with the following properties.

(1) $\mathbf{v}' = (\Sigma', \vec{z}'^{+}, \vec{w}'^{+}) \in \mathfrak{U}(\mathbf{v})$ and $\mathrm{dist}(\mathbf{v}', \mathbf{v}) < \epsilon_1$. Here we use an appropriate metric on $\mathcal{M}_{0;\ell+m}$ to define the distance between $\mathbf{v}'$ and $\mathbf{v}$.

(2) The C^1 distance between the two maps $u' \circ i_{\mathbf{v}'}$ and hu is smaller than ϵ_1 on $\Sigma \setminus \mathcal{U}(S(\Sigma))$.

(3) $\mathrm{diam}(u'(\mathcal{S})) < \epsilon_1$ if $\mathcal{S}$ is any connected component of the complement $\Sigma' \setminus i_{\mathbf{v}'}(\Sigma \setminus \mathcal{U}(S(\Sigma)))$.

(4) $u'(w'^{+}_a) \in hN_{w_a^+}$ for each a and

$$\mathrm{dist}(u'(w'^{+}_a), hu(w_a^+)) < \epsilon_2.$$

Here $\vec{w}'^{+}_a = (w'^{+}_a)_{a\in A}$.

When $\vec{w}^{+}, \vec{w}'^{+}, h$ are specified in the above condition, we say that $\mathbf{x}' = (\Sigma', \vec{z}'^{+}, u')$ is (ϵ_1,ϵ_2)-*close to* $(h\mathbf{x}, \vec{w}^{+})$ with respect to $\vec{w}'^{+}$.

Note Definition 29.14 means that $(\Sigma', \vec{z}'^{+}, u')$ is sufficiently close to $h\mathbf{x}$.

Later on we will take ϵ_1, ϵ_2 so that ϵ_2 small and ϵ_1 is smaller than a number depending on ϵ_2 (and $\mathbf{x}$). (See the proof of Lemma 29.21.)

Lemma 29.15. *Consider u', $\mathbf{v}' = (\Sigma', \vec{z}'^{+}, \vec{w}'^{+})$, h, $\vec{w}^{+}$ given as above and put $\mathbf{x}' = (\Sigma', \vec{z}'^{+}, u')$. Let $\mathbf{g} = (g,\varphi) \in G(\mathbf{x})$ and $\sigma = (\sigma_1,\sigma_2) = \Psi(\varphi)$. Suppose $\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h\mathbf{x}, \vec{w}^{+})$ with respect to $\vec{w}'^{+}$. We consider $\sigma\mathbf{v}' = (\Sigma', \sigma\vec{z}'^{+}, \sigma\vec{w}'^{+})$, where $(\sigma z'^{+})_i = z'^{+}_{\sigma_1^{-1}(i)}$, $(\sigma w'^{+})_a = w'^{+}_{\sigma_2^{-1}(a)}$. We put $\sigma\mathbf{x}' = (\Sigma', \sigma\vec{z}'^{+}, u')$.*

Then $\sigma\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(g^{-1}h\mathbf{x}, \vec{w}^{+})$ with respect to $\sigma\vec{w}'^{+}$.

Proof. Definition 29.14 (1), (3) are easy to check. Let us check Definition 29.14 (2). We remark

$$u' \circ i_{\sigma\mathbf{v}'} = u' \circ i_{\mathbf{v}'} \circ \varphi^{-1}$$

by (29.11). On the other hand,

$$h \circ u \circ \varphi^{-1} = hg^{-1} \circ u,$$

since $(g,\varphi) \in G(\mathbf{x})$. By assumption $u' \circ i_{\mathbf{v}'}$ is C^1 close to $h \circ u$. We thus obtain (2). (Note $hg^{-1} = g^{-1}h$.)

Definition 29.14 (4) can be checked by

$$u'(w'^{+}_{\sigma_2^{-1}(a)}) \in hN_{w^{+}_{\sigma_2^{-1}(a)}} = hN_{\varphi^{-1}(w_a^+)} = hg^{-1}N_{w_a^+}.$$

$\square$

Lemma 29.16. *Let $\mu_1 \in \mathfrak{S}_\ell$ and let ϵ_1, ϵ_2 be sufficiently small. Suppose $\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h\mathbf{x}, \vec{w}^{+})$ with respect to $\vec{w}'^{+}_{(1)}$ and $\mu_1\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h\mathbf{x}, \vec{w}^{+})$ with respect to $\vec{w}'^{+}_{(2)}$.*

Then there exists $\psi \in \mathrm{Aut}_+(\mathbf{v})$ such that $\vec{w}'^{+}_{(2)} = \mu_2\vec{w}'^{+}_{(1)}$ with $\mu = (\mu_1,\mu_2) = \Psi(\psi)$.

Proof. We first prove that there exists $\mu_2 \in \mathfrak{S}_\ell$ such that

$$\vec{w}'^{+}_{(2)} = \mu_2\vec{w}'^{+}_{(1)}. \tag{29.12}$$

We put $\mathbf{v}'_{(k)} = (\Sigma', \vec{z}'^{+}, \vec{w}'^{+}_{(k)})$, $k = 1,2$. For each $a \in \{1,\dots,m\}$, we have ${}_{(2)}w_a^+ \in \Sigma$ such that

$$i_{\mathbf{v}'_{(1)}}({}_{(2)}w_a^+) = w'^{+}_{(2),a}.$$

(We remark $i_{\mathbf{v}'_{(2)}}(w_a^+)$ is close to $w'^{+}_{(2),a}$. But $i_{\mathbf{v}'_{(1)}}(w_a^+)$ may not be close to $w'^{+}_{(2),a}$. So ${}_{(2)}w_a^+$ may be far away from w_a^+.)

Since $hu({}_{(2)}w_a^+)$ is close to $u'(\vec{w}'^{+}_{(2),a})$, there exists ${}_{(3)}w_a^+ \in \Sigma$ that is close to ${}_{(2)}w_a^+ \in \Sigma$ such that u is an immersion at ${}_{(3)}w_a^+$ and

$$u(U_{{}_{(3)}w_a^+}) = u(U_{w_a^+}), \qquad u({}_{(3)}w_a^+) = u(w_a^+), \tag{29.13}$$

where $U_{{}_{(3)}w_a^+}$ and $U_{w_a^+}$ are neighborhoods of ${}_{(3)}w_a^+$ and w_a^+ in Σ, respectively.

We put ${}_{(3)}\vec{w}^+ = ({}_{(3)}w_a^+)_{a\in A}$. Then, $(\Sigma, \mu_1\vec{z}^+, {}_{(3)}\vec{w}^+)$ is close to $\mathbf{v}'_{(2)}$ in $\mathcal{M}_{0;k+\ell}$. On the other hand, by assumption, $(\Sigma, \vec{z}^+, \vec{w}^+)$ is close to $\mathbf{v}'_{(2)}$ in $\mathcal{M}_{0;k+\ell}$.

We remark that ${}_{(3)}w_a^+$ is in the finite set of points such that (29.13) holds. Therefore, we may take ϵ_1, ϵ_2 small enough so that the fact that $(\Sigma, \mu_1\vec{z}^+, {}_{(3)}\vec{w}^+, u)$ is close to $(\Sigma, \vec{z}^+, \vec{w}^+, u)$ implies that they are isomorphic each other. Namely there exists a biholomorphic map $\psi : \Sigma \to \Sigma$ such that $\psi(z_i^+) = z_{\mu_2(i)}^+$ and $\psi(w_i^+) = {}_{(3)}w_i^+$.

By Condition 29.10 (5) it implies that there exists μ_2 such that ${}_{(3)}\vec{w}^+ = \mu_2\vec{w}^+$. For this μ_2 we can prove (29.12) easily.

If we put $\mu = (\mu_1, \mu_2)$, $\Psi(\psi) = \mu$ follows. $\square$


$i_{\mathrm{v}'_{(2)}}\left({}_{(2)}w_a^+\right)$
$w'^+_{(2),a} = i_{\mathrm{v}'_{(1)}}\left({}_{(2)}w_a^+\right)$
Σ'
u'
$hN_{w_a^+}$
$i_{\mathrm{v}'_{(1)}}$
$w'^+_{(1),a}$
u'
$u'\left({}_{(2)}w'^+_a\right)$
$i_{\mathrm{v}'_{(2)}}$
$u'(\Sigma')$
$i_{\mathrm{v}'_{(1)}}$
${}_{(2)}w_a^+$
${}_{(3)}w_a^+$
$u'\left({}_{(1)}w'^+_a\right)$
$u'(\Sigma')$
hu
Σ
ψ
hu
$hu(\Sigma)$
hu
$hu\left({}_{(2)}w_a^+\right)$
$hu\left(w_a^+\right) = hu\left({}_{(3)}w_a^+\right)$
w_a^+

Figure 29.1

Suppose $\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h\mathbf{x},\vec{w}^+)$ with respect to $\vec{w}'^+$. We define

$$\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,h} : E(\mathbf{x}) \to C^\infty(\Sigma', u'^*TX \otimes \Lambda^{0,1}) \tag{29.14}$$

as follows.

For $x \in \Sigma$ in the support of $E(\mathbf{x})$, we consider the minimal geodesic joining $hu(x)$ to $u'(i_{\mathbf{v}'}(x))$ with respect to the given T^n-invariant (almost) Kähler metric. We also use the associated almost Hermitian connection and take the complex linear part of the parallel transport map

$$\mathrm{Pal}_{hu(x)\,u'(i_{\mathbf{v}'}(x)))} : T_{hu(x)}X \to T_{u'(i_{\mathbf{v}'}(x)))}X.$$

Using also the fact that $i_{\mathbf{v}'}$ is holomorphic on the support of $E(\mathbf{x})$ and h is holomorphic, we obtain (29.14) as follows: We take the successive compositions of the following target maps

$$(u^*TX)_x = T_{u(x)}X \xrightarrow{h_*} T_{hu(x)}X \overset{\mathrm{Pal}_{hu(x)\,u'(i_{\mathbf{v}'}(x)))}}{\longrightarrow} T_{u'(i_{\mathbf{v}'}(x)))}X, \tag{29.15}$$

and then take the tensor product with the map

$$\Lambda^{0,1}_x(\Sigma) \overset{(i_{\mathbf{v}'}^{-1})^*}{\longrightarrow} \Lambda^{0,1}_{i_{\mathbf{v}'}(x)}(\Sigma'). \tag{29.16}$$

Lemma 29.17. *In the situation of Lemma 29.16 we have*

$$\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(2)},h} = \mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(1)},h} \circ \psi_*.$$

This lemma is immediate from Lemma 29.16, Formula (29.11) and the definition.

Lemma 29.17 implies that the image of $\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,h}$ is independent of $\vec{w}'^+$.

It follows from Lemma 29.16 that if we perturb h a bit then $\vec{w}'^+$ is uniquely perturbed nearby so that Definition 29.14 remains to be satisfied. Therefore the map $\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,h}$ smoothly changes when we vary h.

We next modify $\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,h}$ and eliminate this h dependence. Suppose $\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h\mathbf{x},\vec{w}^+)$ with respect to $\vec{w}'^+$. We define

$$f(h;(\mathbf{x}',\vec{w}'^+)) = \sum_a \mathrm{dist}(u'(w'^+_a), hu(w^+_a))^2. \tag{29.17}$$

Here $\mathrm{dist} = \mathrm{dist}_{hN_{w^+_a}}$ is the distance function of the submanifold $hN_{w^+_a}$ induced from the given ambient Kähler metric. Recall the square of the distance function is strictly convex and smooth in a neighborhood of the diagonal.

Corollary 29.18. *Let* $(\mathbf{x}',h,\vec{w}'^+_{(1)})$ *and* $(\mu_1\mathbf{x}',h,\vec{w}'^+_{(2)})$ *be as in Lemma 29.16. Then*

$$f(h;(\mathbf{x}',\vec{w}'^+_{(1)})) = f(h;(\mu_1\mathbf{x}',\vec{w}'^+_{(2)})).$$

This is immediate from Lemma 29.16 and the definition of f.

Corollary 29.18, in the case of $\mu_1 = id$, implies that $f(h;(\mathbf{x}',\vec{w}'^+))$ depends only on $h,\mathbf{x}'$ and is independent of $\vec{w}'^+$. Hereafter we write $f(h;\mathbf{x}')$ instead of $f(h;(\mathbf{x}',\vec{w}'^+))$.

For given $\mathbf{x}',\vec{w}^+$, our function f is a function of h. More precisely speaking, we define the subset $\mathfrak{H}(\mathbf{x}') \subset T^n$ by

$$\mathfrak{H}(\mathbf{x}') = \{h \in T^n \mid \exists \mu_1 \in \mathfrak{S}_\ell,\ \ \mu_1\mathbf{x}' \text{ is } (\epsilon_1,\epsilon_2)\text{-close to } (h\mathbf{x},\vec{w}^+)\}.$$

This is an open subset of T^n for each given $\mathbf{x}'$. Then for each given $\mathbf{x}'$ and sufficiently small $\epsilon_1,\epsilon_2 > 0$, the assignment $h \mapsto f(h;\mu_1\mathbf{x}')$ defines a function on $\mathfrak{H}(\mathbf{x}')$. (Here μ_1 is as in the definition of $\mathfrak{H}(\mathbf{x}')$.) We write this function $f(h;[\mathbf{x}'])$

Since $\mathfrak{H}(\mathbf{x}')$ is an open subset of T^n, it carries a natural affine structure. We use this to define convexity used in the statement of the following lemma.

Lemma 29.19. *If ϵ_1, ϵ_2 are sufficiently small, then $f(\cdot;[\mathbf{x}']) : \mathfrak{H}(\mathbf{x}') \to \mathbb{R}_{\ge 0}$ is strictly convex.*

Proof. This is a consequence of strictly convexity of the square of the Riemannian distance function and Item (4) of Condition 29.10. □

Lemma 29.20. *Let $\mathbf{g} = (g, \varphi) \in G(\mathbf{x})$. Then we have*

$$f(h;[\mathbf{x}']) = f(h;[\mathbf{g}\mathbf{x}']).$$

Here $\mathbf{x}' = (\Sigma', \vec{z}^{+'}, u')$ and $\mathbf{g}\mathbf{x}' = (\Sigma', \Psi(\varphi)\vec{z}^{+'}, gu')$.

Proof. Lemma 29.15 imlies

$$f(h;[\mathbf{x}']) = f(g^{-1}h;[\Psi(\varphi)\mathbf{x}']).$$

On the other hand, it is easy to see that

$$f(h;[g\mathbf{x}']) = f(gh;[\mathbf{x}']).$$

The lemma follows easily. □

We put

$$|G(\mathbf{x})| = \{g \mid \exists \varphi, (g, \varphi) \in G(\mathbf{x})\}.$$

Let $\mathfrak{c} = \#|G(\mathbf{x})|$ be its order and write as $|G(\mathbf{x})| = \{g_1 = 1, \dots, g_{\mathfrak{c}}\}$. We put

$$G(\mathbf{x}) = \{\mathbf{g}_{i,j} = (g_i, \varphi_{i,j}) \mid i = 1, \dots, \mathfrak{c},\ j = 1, \dots, \mathfrak{d}\}.$$

Here $\mathfrak{d} = \#\{\varphi \mid (1, \varphi) \in G(\mathbf{x})\}$ and $\varphi_{1,1} = id$.

Lemma 29.21. *For each sufficiently small ϵ_2 we may choose ϵ_1 small enough so that the following holds: Whenever $\mathfrak{H}(\mathbf{x}')$ is nonempty, $f(\cdot;[\mathbf{x}']) : \mathfrak{H}(\mathbf{x}') \to \mathbb{R}_{\ge 0}$ assumes local minima at exactly $\mathfrak{c}$ points $h_1, \dots, h_{\mathfrak{c}}$.*

Furthermore the following also holds: For each i there exist $\mathfrak{d}$ choices $\vec{w}'^{+}_{(i,j)}$ and $\rho_{(i,j),1} \in \mathfrak{S}_\ell$ such that $\rho_{(i,j),1}\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h_i\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^{+}_{(i,j)}$.

If we put $\sigma_{(i,j)} = (\sigma_{(i,j),1}, \sigma_{(i,j),2}) = \Psi(\varphi_{i,j})$ for each $\mathbf{g}_{i,j} = (g_i, \varphi_{i,j})$, we have

$$h_i = h_1 g_i^{-1} \tag{29.18}$$

and

$$\rho_{(i,j),1} = \sigma_{(i,j),1}\rho_{(1,1),1}, \qquad w'^{+}_{(i,j)} = \sigma_{(i,j),2} w'^{+}_{(1,1)}. \tag{29.19}$$

Proof. We first consider the case $\mathbf{x}' = \mathbf{x}$. By Lemma 29.15 we have

$$f(h;\sigma_{(i,j),1}\mathbf{x}') = f(g_i^{-1}h;\mathbf{x}'). \tag{29.20}$$

Therefore, $(\Sigma, \rho_{(i,j),1}\vec{z}^+, u)$ is (ϵ_1, ϵ_2)-close to $(g_i^{-1}\mathbf{x}, \vec{w}^+)$ with respect to $\rho_{(i,j),2}\vec{w}^+$. Moreover, g_i^{-1} is a critical point of $f(\cdot;\sigma_{(i,j),1}\mathbf{x})$. Namely (29.18) and (29.19) are satisfied with $h_1 = 1$, $\rho(1,1), 1 = id$, $w'^{+}_{(1,1)} = \vec{w}^+$.

We can prove that they are all the choices in the same way as the proof of Lemma 29.16.

Let us consider the general $\mathbf{x}' = (\Sigma', \vec{z}'^{+}, u')$. Suppose $h \in \mathfrak{H}(\mathbf{x}')$. Replacing u' by $h^{-1}u'$ we may assume $h = 1$. Then we can find a short path $\mathbf{x}(t)$ joining $\mathbf{x}$ to $\mathbf{x}'$. By the strict convexity of f, the number of local minima is constant on this (short) path. We may choose ϵ_1 small enough so that the distance between $u'(w'^{+}_{(i,j),t;a})$

and $h_{i,t}u(w_a^+)$ does not go beyond ϵ_2 while t moves from 0 to 1. (Here $h_{i,t}$ and $w'^+_{a,t}$ are such that $\rho_{(i,j),1}\mathbf{x}(t)$ is (ϵ_1,ϵ_2)-close to $(h_{i,t}\mathbf{x},\vec w^+)$ with respect to $\vec w'^+_{(i,j),t}$.)

We observe that if (29.18), (29.19) hold and $\rho_{(1,1),1}\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h_1\mathbf{x},\vec w^+)$ with respect to $\vec w'^+_{(1,1)}$, then

$$u'(\vec w'_{(i,j),a}) = u'(\vec w'_{(1,1),\sigma^{-1}_{(i,j),2}(a)}) \in h_1 N_{w^+_{\sigma^{-1}_{(i,j),2}(a)}} = h_1 g_i^{-1} N_{w_a^+}.$$

Namely $\rho_{(i,j),1}\mathbf{x}'$ is (ϵ_1,ϵ_2)-close to $(h_i\mathbf{x},\vec w^+)$ with respect to $\vec w'^+_{(i,j)}$.

Using Lemma 29.20, we can show that $f(\cdot;[\mathbf{x}'])$ assumes local minima at h_i. The proof of (29.18) of Lemma 29.21 is complete.

We can prove (29.19) using the fact that it holds for $t=0$ and the continuity argument. □

Lemma 29.22. *In the situation of Lemma 29.21 we have*

$$\mathcal{P}_{\sigma_{(i,j),1}\mathbf{x}',\vec w^+,\vec w'^+_{(i,j)},h_i} = \mathcal{P}_{\sigma_{(1,1),1}\mathbf{x}',\vec w^+,\vec w'^+_{(1,1)},h_1} \circ (\varphi_{i,j})_*.$$

Proof. Let $\mathbf{v}'_{(i,j)} = (\Sigma',\rho_{(i,j),1}\vec z^+,\vec w'^+_{(i,j))}) = \rho_{(i,j)}\mathbf{v}'$.

By (29.15),(29.16), we have

$$\mathcal{P}_{\sigma_{(i,j),1}\mathbf{x}',\vec w^+_{(i,j)},\vec w'^+_{(i,j)},h_i} = \mathrm{Pal}_{h_iu(x),u'(i_{\mathbf{v}'_{(i,j)}}(x))} \otimes (i^{-1}_{\mathbf{v}'_{(i,j)}(x)})^*. \tag{29.21}$$

By (29.18), the right hand side of Formula (29.21) is

$$\mathrm{Pal}_{h_1g_iu(x),u'(i_{\rho_{(i,j)}\mathbf{v}'}(x))} \otimes (i^{-1}_{\rho_{(i,j)}\mathbf{v}'})^*.$$

By (29.11), we have

$$i_{\rho_{(i,j)}\mathbf{v}'} = i_{\sigma_{(i,j)}\rho_{(1,1)}\mathbf{v}'} = i_{\rho_{(1,1)}\mathbf{v}'}\circ\varphi_{i,j}^{-1} = i_{\mathbf{v}'_{(1,1)}}\circ\varphi_{i,j}^{-1}.$$

Moreover $h_1g_i^{-1}u = h_1u\varphi_{i,j}^{-1}$. The lemma follows. □

We now define $\mathcal{P}_{\mathbf{x}'} : E(\mathbf{x}) \to C^\infty(\Sigma', u'^*TX\otimes\Lambda^{0,1})$ by

$$\mathcal{P}_{\mathbf{x}'} = \mathcal{P}_{\mathbf{x}',\vec w^+,\vec w'^+_{(i,j)},h_i} \circ (\varphi_{i,j})_*^{-1}.$$

This is independent of $i=1,\dots,\mathfrak{c}$ by Lemma 29.22 and of $j=1,\dots,\mathfrak{d}$ by Lemma 29.16.

Lemma 29.23. *For any $h\in T^n$, we have*

$$\mathcal{P}_{h\mathbf{x}'} = h_*\mathcal{P}_{\mathbf{x}'}. \tag{29.22}$$

Proof. This is immediate from construction. □

We now put $E(\mathbf{x}') = \mathrm{Im}\mathcal{P}_{\mathbf{x}'}$.

So far we have fixed the choice of $\mathbf{x}$, $\vec w$ and the choice of $E(\mathbf{x})$. Actually, we took $T^n\times\mathfrak{S}_\ell$ orbit of $\mathbf{x}$ and $E(g\rho\mathbf{x})$ for each $g\in T^n$, $\rho\in\mathfrak{S}_\ell$ which is invariant in the sense of Lemma 29.8.

So we need to show that $E(\mathbf{x}')$ does not change if we replace $\mathbf{x}$ to $g\rho\mathbf{x}$. We will prove this fact below. During this proof we write $\mathcal{P}^{\mathbf{x},\vec w^+}_{\mathbf{x}'}$ in place of $\mathcal{P}_{\mathbf{x}'}$ to specify the choice of $\mathbf{x},\vec w^+$.

We first discuss the choice of $\vec w^+$. We consider the following two conditions:

Condition 29.24. (1) If $g\in T^n$, then we take the same choice $\vec w^+$ for $g\mathbf{x} = (\Sigma,\vec z^+,gu)$ as $\mathbf{x} = (\Sigma,\vec z^+,u)$.

(2) Let $\varphi : \Sigma \to \Sigma$ be an isomorphism that preserves $\vec{z}^+$ as a set and denote by $\varphi \cdot \vec{z}^+$ the induced ordered set of marked points defined by $(\varphi \cdot \vec{z}^+)_a = \varphi^{-1}(z_a^+)$. Then our choice of $\vec{w}^+$ for $\varphi \cdot \mathbf{x} = (\Sigma, \varphi \cdot \vec{z}^+, u)$ is $\varphi \cdot \vec{w}^+$ which is defined similarly as $\varphi \cdot \vec{z}^+$.

We remark that if we set $\mathbf{v} = (\Sigma, \vec{z}^+, \vec{w}^+)$ and $\Psi(\varphi) = (\rho_1, \rho_2)$, then the above choice made in (2) is consistent with the action given in (29.8).

However, there is *no* such choice of $\vec{w}^+$ that satisfies both of the above two rules. If $(g, \varphi) \in G(\mathbf{x})$, then

$$g\mathbf{x} \sim \varphi \cdot \mathbf{x} \tag{29.23}$$

in $\mathcal{M}_{0;\ell}(\beta)$. The left hand side gives the choice $\vec{w}^+$, the right hand side gives $\varphi\vec{w}^+$. This actually does not matter since the isomorphism (29.23) is realized by φ.

The point to take care of is the fact that this isomorphism may not be unique. Namely, if there exists $\psi : \Sigma \to \Sigma$ which fixes all the points of $\vec{z}^+$, then both of φ and $\varphi \circ \psi$ represent isomorphisms (29.23). But $\vec{w}^+$ and $\psi\vec{w}^+$ are different from each other unless $\psi = identity$. However they will give the same choice of the image of $\mathcal{P}_{\mathbf{x}'}$. Namely we find the following:

Lemma 29.25.

$$\mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\vec{w}^+} = \mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\psi\cdot\vec{w}^+},$$

if $\psi : \Sigma \to \Sigma$ is a biholomorphic map that fixes all the elements of $\vec{z}^+$.

Proof. We can easily prove the equality

$$\mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\vec{w}^+} \circ \psi_* = \mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\psi\cdot\vec{w}^+},$$

from which the lemma follows. □

We put

$$\mathrm{Aut}(\Sigma, \vec{z}^+) = \{\psi \mid \psi : \Sigma \to \Sigma \text{ is a biholomorphic map such that } \psi(z_i^+) = z_i^+\}.$$

Lemma 29.26. *We consider the $T^n \times \mathrm{Aut}_+(\Sigma, \vec{z})$ orbit of $\mathbf{x}$. Then for each $\varphi g\mathbf{x}$ of it we can associate an $\mathrm{Aut}(\Sigma, \varphi\vec{z}^+)$ orbit of the set of $\vec{w}^+$'s such that Condition 29.24 is satisfied.*

The proof is easy and is omitted. The proof of the next lemma is also easy.

Lemma 29.27. *Suppose that Condition 29.24 is satisfied. Then we have*

(1) $\mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{g\mathbf{x},\vec{w}^+} = g_*\mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\vec{w}^+}$.

(2) $\mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{\varphi\cdot\mathbf{x},\varphi\cdot\vec{w}^+} = \mathrm{Im}\mathcal{P}_{\mathbf{x}'}^{\mathbf{x},\vec{w}^+}$.

Consider the image $\Psi_1(\mathrm{Aut}_+(\Sigma, \vec{z})) \subseteq \mathfrak{S}_\ell$ and let $\rho_1, \dots, \rho_K$ be the representative of the left coset space $\mathfrak{S}_\ell/\Psi_1(\mathrm{Aut}_+(\Sigma, \vec{z}))$.

For each $(g, \rho) \in T^n \times \mathfrak{S}_\ell$ we can write $\rho = \rho_i\Psi_1(\varphi)$ for some $i = 1, \dots, K$ and $\varphi \in \mathrm{Aut}_+(\Sigma, \vec{z})$. We then choose $\vec{w}^+$ for $(g, \rho)\mathbf{x}$ as the same $\vec{w}^+$ associated to $g\rho\mathbf{x}$ by Lemma 29.26. It is well-defined up to the action of $\mathrm{Aut}((g, \rho)\mathbf{x})$.

We can then define $E(\mathbf{x}')$ for each $\mathbf{x}'$ sufficiently close to $(g, \rho)\mathbf{x}$ for some g, ρ. (We use Lemma 29.25 to prove that the $\mathrm{Aut}((g, \rho)\mathbf{x})$-ambiguity does not affect the choice of the subspace $E(\mathbf{x}')$.)

We use the next lemma and Lemma 29.27 to show that $E(\mathbf{x})$ is independent of $(g, \rho)\mathbf{x}$ that is close to $\mathbf{x}'$.

Lemma 29.28. *Let* $\mathbf{x}'$ *is* (ϵ_1, ϵ_2)*-close to* $\mathbf{x}$*. Then the following conditions for* $\rho \in \mathfrak{S}_\ell$ *are equivalent to each other:*

1. *There exists* $(g, \varphi) \in G(\mathbf{x})$ *such that* $\rho = \Psi_1(\varphi)$*.*
2. *There exists* $g \in T^n$ *such that* $\mathbf{x}'$ *is* (ϵ_1, ϵ_2)*-close to* $g\rho\mathbf{x}$*.*

Note we write "(ϵ_1, ϵ_2)-close to $g\rho\mathbf{x}$" instead of "(ϵ_1, ϵ_2)-close to $(g\rho\mathbf{x}, \vec{w}^+)$" since we already specified the choice of $\vec{w}^+$ for each $g\rho\mathbf{x}$.

Proof. In case $\mathbf{x}' = \mathbf{x}$ this is easy to see. The general case can be reduced to this case in the same way as the proof of Lemma 29.16. □

Thus we have associated $E(\mathbf{x}')$ to each $\mathbf{x}'$ that is close to the $T^n \times \mathfrak{S}_\ell$-orbit of $\mathbf{x}$. By construction $\mathbf{x}' \mapsto E(\mathbf{x}')$ is $T^n \times \mathfrak{S}_\ell$ equivariant.

We consider the set of $\mathbf{x}'$ satisfying

$$\overline{\partial} u' \in E(\mathbf{x}').$$

Together with E and

$$s(\mathbf{x}') = \overline{\partial} u',$$

this defines a Kuranishi neighborhood of the $T^n \times \mathfrak{S}_\ell$ orbit of $\mathbf{x}$. (29.22) implies that it is T^n invariant. By construction, it is invariant under the permutation of the interior marked points $\vec{z}^+$.

Remark 29.29. In [FO] Appendix, we took the minimum choice of extra marked points which breaks the symmetry with respect to the automorphism. The symmetry is recovered in [FO] Appendix later on. In this section we take a slightly different way and take more additional marked points so that they are preserved by the automorphisms. This approach may be more suitable than the one in [FO] in the situation where a continuous group of symmetry exists. (It occurs in the situation we handle next.)

We next remove Condition 29.9.

Lemma 29.30. *Suppose* $\mathbf{x} = (\Sigma, \vec{z}^+, u)$ *does not satisfy Condition 29.9. Then there exists a splitting* $T^n = S^1 \times T^{n-1}$ *with the following properties.*

1. *If* $\mathfrak{t} \in Lie(T^{n-1})$*, then there exists a (disk or sphere) component* Σ_a *of* Σ *and* $p \in \Sigma_a$ *such that* u *is an immersion at* p *and* $\mathfrak{t}(u(p))$ *is not contained in* $u_* T_p \Sigma$*.* p *is not a singular, boundary, or marked point. Here we regard* $\mathfrak{t}$ *as a vector field on* X*.*
2. *If* $\mathfrak{t} \in Lie(S^1)$ *and* $w \in \Sigma$ *is not a singular point and* u *is an immersion at* w*, then* $\mathfrak{t}(u(w)) \in u_*(T_w \Sigma)$*.*

Proof. Let Σ_a be a component the restriction of u to which is nontrivial. Since T^n orbits are totally real, there exists at most one dimensional subspace $\mathfrak{T} \subset Lie(T^n)$ such that the following holds: If $\mathfrak{t} \notin Lie(\mathfrak{T})$ and $p \in \Sigma_a$ is a generic point, then $\mathfrak{t}(u(p))$ is not contained in $u_* T_p \Sigma$. Moreover $\mathfrak{t} \in \mathfrak{T}$ satisfies (2) above.

Since Condition 29.9 is not satisfied, this subspace $\mathfrak{T}$ is nonempty. Moreover it is independent of the component Σ_a.

We note that the Lie group associated to $\mathfrak{T}$ must be compact since otherwise its closure in T^n would have dimension 2 or more and its Lie algebra generates the set of Killing vector fields which spans the tangent space of the image of u. This gives rise to a contradiction since u is J-holomorphic.

Since all S^1 in T^n are direct summands, we obtain the required splitting. □

Hereafter we write

$$T^n = S^1_0 \times T^{n-1}_0$$

for the splitting in Lemma 29.30. (The factor T^{n-1}_0 is not unique. However we take one and fix it throughout.)

Lemma 29.31. *Suppose* $\mathbf{x} = (\Sigma, \vec z^+, u)$ *does not satisfy Condition 29.9. Then one of the following two altermatives holds.*

1. *There exists a special (singular or marked) point* z_0 *on* $(\Sigma, \vec z^+)$ *such that* $u(z_0)$ *is not a fixed point of* S^1_0. *The group* $G(\mathbf{x})$ *is finite.*
2. *The connected component of* $G(\mathbf{x})$ *is* S^1. *The restriction of the map* $G(\mathbf{x}) \to T^n$, $(g, \varphi) \mapsto g$ *to the connected component induces a finite covering of* S^1_0.

Proof. Suppose there exists a special point z_0 such that $u(z_0)$ is not a fixed point of S^1_0. Then clearly the Lie algebra of S^1_0 does not vanish on $u(z_0)$. It follows easily that $G(\mathbf{x})$ is finite.

We next assume that, for any special point z_0 of $(\Sigma, \vec z^+)$, the point $u(z_0)$ is a fixed point of S^1_0. Then it is easy to see that the vector field generated by $\mathfrak{t} \in Lie(S^1_0)$ lifts to a holomorphic vector field on Σ that vanishes on the special points. So by integrating it we obtain a subgroup of $G(\mathbf{x})$ which is of positive dimension. Clearly the connected component of $G(\mathbf{x})$ maps to S^1_0 by the map $(g, \varphi) \mapsto g$. The lemma follows. (Note in case u is a multiple cover of multiplicity k on all the components where S^1_0 action is nontrivial, then the covering index of the connected coponent of $G(\mathbf{x})$ to S^1_0 is k.) ☐

We discuss the two cases in Lemma 29.31 separately. We first consider case (1). Let $(\Sigma, \vec z^+)$ be our marked bordered Riemann surface of genus zero and one boundary marked point and let $u : (\Sigma, \partial\Sigma) \to (X, L)$ be pseudo-holomorphic. We choose finitely many points $\vec w^+ = (w^+_a)_{a \in A}$ so that the following is satisfied.

Condition 29.32.

1. w^+_a is disjoint from $\vec z^+$, singular point set and the boundary of Σ.
2. $(\Sigma, \vec w^+)$ is stable. Moreover the group of its automorpisms is trivial.
3. u is an immersion at w^+_a.
4. For any $\mathfrak{t} \in Lie(T^{n-1}_0)$ there exists w^+_a such that $\mathfrak{t}(u(w^+_a))$ is not contained in $u_* T_{w^+_a}\Sigma$.
5. $\vec w^+$ is $G(\mathbf{x})$ invariant, i.e., $\varphi(w^+_a) \in \vec w^+$ for all $a \in A$, $(g, \varphi) \in G(\mathbf{x})$.

This is almost the same as Condition 29.10. The only difference is we take T^{n-1}_0 in place of T^n in (4). The existence of such $\vec w^+$ follows from Lemma 29.30.

Let $\vec w^+$ satisfy Condition 29.32. We put $\mathbf{v} = (\Sigma, \vec z^+ \cup \vec w^+)$. We consider its neighborhood $\mathfrak{U}(\mathbf{v})$ as before.

We next consider $\mathbf{x}' = (\Sigma', \vec z'^+, u')$ which satisfies the conditions in Definition 29.14. We define the set $\mathfrak{H}(\mathbf{x}')$ and a function $f(\cdot; \mathbf{x}')$ or $f(\cdot; [\mathbf{x}'])$ on it in the same way.

We remark however that f may not be strictly convex in the S^1_0 direction. (This is because we replace T^n by T^{n-1}_0 in Condition 29.32 (4).)

So to generalize Lemma 29.21 to our situation we need to appropriately modify f. We use the special points on $(\Sigma, \vec z^+)$ for this purpose as follows.

Let $\mathbf{x}' = (\Sigma', \vec z'^+, u')$. Let z_c be a special point of $(\Sigma, \vec z^+)$. Below, we associate a real number $f_c(h)$ to $(h; \mathbf{x}')$. If z_c is an (interior) marked point, then there exists a corresponding marked point z'_c on Σ'. We put

$$f_c(h;\mathbf{x}') = \mathrm{dist}(u'(z'_c), hu(z_c))^2. \tag{29.24}$$

Here dist is a Riemannian metric on X which is invariant under the T^n action.

We next consider the case when z_c is a singular point. If this point remains singular in $\mathbf{v}'$, there exists a corresponding singular point z'_c on Σ'. In that case we define $f_c(h;\mathbf{x}')$ by (29.24).

In case z_c is no longer singular in $\mathbf{v}'$, we proceed as follows. We take the unique Riemannian metric on the double of $\Sigma' \setminus (\vec{z}'^{+} \cup \vec{w}'^{+})$ which is complete of constant negative curvature -1, of finite volume and whose conformal structure is compatible with the complex structure of Σ'. By Margulis' lemma (see p.107 [BGS] for example), there exists a unique short closed geodesic which lies in the neck part that appears after resolving the singularity z_c. Let $\ell_c : [0,\epsilon_c] \to \Sigma'$ be this closed geodesic parametrized by arc length. We now put

$$f_c(h;\mathbf{x}') = \frac{1}{\epsilon_c}\int_{t\in[0,\epsilon_c]} \mathrm{dist}(u'(\ell_c(t)), hu(z_c))^2 dt. \tag{29.25}$$

We then define

$$f_+(h;\mathbf{x}') = f(h;\mathbf{x}') + \sum_c f_c(h;\mathbf{x}'). \tag{29.26}$$

Now by Lemma 29.31 (1), $f_+(\cdot;\mathbf{x}')$ is *strictly* convex. We can perform the rest of the construction in the same way using f_+ instead of f, and obtain $\mathcal{P}_{\mathbf{x}'}$ and $E(\mathbf{x}')$. We thus obtain a Kuranishi chart of a neighborhood of the $T^n \times \mathfrak{S}_\ell$ orbit of $\mathbf{x}$.

We next consider the case where Lemma 29.31 (2) holds. In our situation, the connected component of $G(\mathbf{x})$ is S^1 and is a finite cover of S^1_0.

Lemma 29.33. *There exists $\mathbb{S} = \cup_{b\in B} S^1_b$ with $S^1_b \cong S^1 \subset \Sigma$ satisfying the following properties:*

1. *Each of $u(S^1_b) \subset X$ is an S^1_0 orbit.*
2. *$\mathbb{S}$ is $G(\mathbf{x})$ invariant.*
3. *$\mathbb{S}$ does not intersect with singular or marked points. It is disjoint from boundary.*
4. *For each choice of $w^+_b \in S^1_b$ ($b \in B$), the pair $(\Sigma, \{w^+_b \mid b \in B\})$ is stable. Moreover its has no nontrivial automorphisms.*
5. *If $w \in \mathbb{S}$, then u is an immersion at w.*
6. *For each $\mathfrak{t} \in Lie(T^{n-1}_0)$ there exists b such that $\mathfrak{t}(w^+_b) \notin u_* T_{w^+_b}(\Sigma)$ for any $w^+_b \in S^1_b$.*

Proof. By S^1_0 equivalence if (5) holds for some w^+_b then it holds for all w^+_b. Once we remark this point the rest of the proof is easy and is omitted. □

Let

$$G_0(\mathbf{x}) = \{(g,\varphi) \in G(\mathbf{x}) \mid g \in T^{n-1}_0\}. \tag{29.27}$$

Then $G_0(\mathbf{x})$ is a finite group.

We consider $\vec{w}^+ = (w^+_a)_{a\in A}$ together a surjective map $A \to B$, $a \mapsto b(a)$ with the following properties. Hereafter we write $S_a = S_{b(a)}$ for $a \in A$.

Condition 29.34. (1) $w^+_a \in S^1_a$.

(2) If $(g,\varphi) \in G_0(\mathbf{x})$, then $\varphi(w^+_a) \in \vec{w}^+$.

(3) Let $m = \#A$ and we identify $A = \{1, \dots, m\}$. For each $(g, \varphi) \in G(\mathbf{x})$ there exists $\rho \in \mathfrak{S}_m$, $t_a \in S^1_0$ $(a \in A)$ such that

$$\varphi(w^+_a) = t_a w^+_{\rho(a)}$$

Remark 29.35. In general, $\#A \ge \#B$. We may need to take $\#A > \#B$ for item (2) of Condition 29.34 to be satisfied.

We define

$$\mathbf{S} = \{(w^+_a)_{a\in A} \mid \text{Condition 29.34 is satisfied.}\}. \tag{29.28}$$

We remark all the elements of $\mathbf{S}$ satisfy (3)-(6) of Lemma 29.33.

The connected components of $\mathbf{S}$ is identified with $T^{m'}$ for some $m' \le m$. The number of connected components are finite. So $\mathbf{S}$ has a canonical probability measure.

We put $\mathbf{v}(\vec{w}^+) = (\Sigma, \vec{z}^+ \cup \vec{w}^+) \in \mathcal{M}_{0;\ell+m}$. (Here $\mathbf{v} = (\Sigma, \vec{z}^+)$ and $\mathbf{x} = (\Sigma, \vec{z}^+, u)$.)

For each $\mathbf{g} = (g, \varphi) \in G(\mathbf{x})$ and $w \in S^1_a$ we choose N_w satisfying the following conditions.

Condition 29.36. (1) N_w is a smooth submanifold of codimension 2.
(2) N_w intersects transversally with $u(\Sigma)$ at $u(w)$.
(3) If $(g, \varphi) \in G(\mathbf{x})$, then $gN_w = N_{\varphi(w)}$.

We next consider $\mathbf{x}' = (\Sigma', \vec{z}', u')$, $\vec{w}^+ \in \mathbf{S}$ and the following condition on it. Let ϵ_1, ϵ_2 be positive real numbers.

Definition 29.37. Let $h \in T^n$. We say that $\mathbf{x}'$ is *(ϵ_1, ϵ_2)-close to $h\mathbf{x}$* if there exists $\vec{w}'^+ \subset \Sigma'$, and $h \in T^{n-1}_0$ with the following properties.

(1) $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+) \in \mathfrak{U}(\mathbf{v}(\vec{w}^+))$ and $\mathrm{dist}(\mathbf{v}', \mathbf{v}(\vec{w}^+)) < \epsilon_1$. Here we use an appropriate metric on $\mathcal{M}_{0;\ell+m}$ to define the distance between $\mathbf{v}'$ and $\mathbf{v}(\vec{w}^+)$.
(2) The C^1 distance between two maps $u' \circ i_{\mathbf{v}'}$ and hu is smaller than ϵ_1 on $\Sigma \setminus \mathcal{U}(S(\Sigma))$.
(3) $\mathrm{diam}(u'(\mathcal{S})) < \epsilon_1$ if $\mathcal{S}$ is any connected component of the complement $\Sigma' \setminus i_{\mathbf{v}'}(\Sigma \setminus \mathcal{U}(S(\Sigma)))$.
(4) $u'(w'^+_a) \in hN_{w^+_a}$ for each a and

$$\mathrm{dist}(u'(w'^+_a), hu(w^+_a)) < \epsilon_2.$$

Here $\vec{w}'^+_a = (w'^+_a)_{a\in A}$.

In case $\vec{w}^+$, $\vec{w}'^+$ are specified we say that $\mathbf{x}'$ *is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+)$* with respect to $\vec{w}'^+$.

This definition is almost the same as Definition 29.14. The only difference is we take T^{n-1}_0 in place of T^n in (2).

Lemma 29.38. *We consider u', $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+)$, h, $\vec{w}^+$ and put $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$. Let $\mathbf{g} = (g, \varphi) \in G(\mathbf{x})$. We put $\sigma = \Psi(\varphi)$. Suppose $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+$. We consider $\sigma\mathbf{v}' = (\Sigma', \sigma\vec{z}'^+, \sigma\vec{w}'^+)$, where $(\sigma z'^+)_i = z'^+_{\sigma_1^{-1}(i)}$, $(\sigma w'^+)_a = w'^+_{\sigma_2^{-1}(a)}$. Then $\sigma\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(g^{-1}h\mathbf{x}, \vec{w}^+)$ with respect to $\sigma_2\vec{w}'^+$.*

Proof. The same as the proof of Lemma 29.15. $\square$

Lemma 29.39. *Let $\mu_1 \in \mathfrak{S}_\ell$ and ϵ_1, ϵ_2 be sufficiently small. Suppose $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(1)}$ and $\mu_1\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(2)}$. Then there exists $\psi \in \mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$ such that $\vec{w}'^+_{(2)} = \mu_2\vec{w}'^+_{(1)}$ with $\mu = (\mu_1, \mu_2) = \Psi(\psi)$.*

Proof. The proof is the same as that of Lemma 29.16. □

Lemma 29.40. *For each given $\vec{\epsilon} = (\epsilon_1, \epsilon_2)$ there exists another $(\epsilon_1^0(\vec{\epsilon}), \epsilon_2^0(\vec{\epsilon}))$ such that the following holds for $\epsilon'_1 < \epsilon_1^0(\vec{\epsilon})$, $\epsilon'_2 < \epsilon_2^0(\vec{\epsilon})$:*

Suppose $\mathbf{x}'$ is $(\epsilon'_1, \epsilon'_2)$ close to $(h\mathbf{x}, \vec{w}^+_{(1)})$ with respect to $\vec{w}'^+_{(1)}$ and let $\vec{w}^+_{(2)} \in \mathbf{S}$. Then there exists $\vec{w}'^+_{(2)}$ such that $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+_{(2)})$ with respect to $\vec{w}'^+_{(2)}$.

Proof. Let $\vec{w}^+_{(k)} = (w^+_{(k),a})_{a\in A}$ $(k = 1, 2)$ and $\vec{w}'^+_{(1)} = (w'^+_{(1),a})_{a\in A}$. We can find a path $t \mapsto w^+_a(t) \in S^1_a$ (whose length can be bounded by a constant independent of ϵ_1, ϵ_2) such that $w^+_a(0) = w^+_{(1),a}$, $w^+_a(1) = w^+_{(2),a}$. By Definition 29.37 (2) and 29.36 (2), we can apply the implicit function theorem to find a unique C^∞ map $t \mapsto w'^+_a(t) \in \Sigma'$ such that

$$u'(w'^+_a(t)) \in hN_{w^+_{\rho(a)}(t)}, \quad u'(w'^+_a(0)) = w'^+_{(1),a}.$$

Then $w'^+_{(2),a} = u'(w'^+_a(1))$ has the required properties. (We use continuity of $i_{\mathbf{v}'}$ with respect to $\mathbf{v}'$ to obtain Condition (2) of Definition 29.37.) □


Σ'
$w'^{+}_{(1),a}$
$w'^{+}_{a}(t)$
$w'^{+}_{(2),a}$
u'
Σ
$w^{+}_{(1),a} = w^{+}_{a}(0)$
$w^{+}_{a}(t)$
$w^{+}_{(2),a} = w^{+}_{a}(1)$
hu
$u'\left(w'^{+}_{(1),a}\right)$
$hu\left(w^{+}_{(2),a}\right)$
$N_{w^{+}_{a}(t)}$
$N_{w^{+}_{(2),a}}$
$hu\left(w^{+}_{a}(t)\right)$
$u'(\Sigma')$
$hu(\Sigma)$
$hu\left(w^{+}_{(1),a}\right)$
$u'\left(w'^{+}_{a}(t)\right)$
$u'\left(w'^{+}_{(2),a}\right)$
$N_{w^{+}_{(1),a}}$

Figure 29.2

Remark 29.41. Note that $\vec{w}^+(t)$ may not be an element of $\mathbf{S}$ for $t \neq 0, 1$. This does not affect the above argument.

For given $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ and $\vec{w}^+ \in \mathbf{S}$, we define the subset $\mathfrak{H}(\mathbf{x}', \vec{w}^+)$ of T^{n-1} by

$$\mathfrak{H}(\mathbf{x}', \vec{w}^+) = \{h \in T^{n-1} \mid \exists \mu_1 \in \mathfrak{S}_\ell,\ \ \mu_1\mathbf{x}' \text{ is } (\epsilon_1, \epsilon_2)\text{-close to } (h\mathbf{x}, \vec{w}^+).\}$$

We define $f(\cdot; [\mathbf{x}', \vec{w}^+]) : \mathfrak{H}(\mathbf{x}', \vec{w}^+) \to \mathbb{R}$ by the same formula as (29.17). (It is independent of $\vec{w}'^+$ as before.)

Remark 29.42. In the previous situation, we chose $\vec{w}^+$ once and fixed it. Here $\vec{w}^+ \in \mathbf{S}$ is moving. So we include $\vec{w}^+$ in the notation of $\mathfrak{H}(\mathbf{x}', \vec{w}^+)$ and of f.

We define

$$|G_0(\mathbf{x})| = \{g \in T_0^{n-1} \mid \exists \varphi, (g, \varphi) \in G(\mathbf{x})\}$$

and $\mathfrak{c} = \#|G_0(\mathbf{x})|$ its order. Let $|G_0(\mathbf{x})| = \{g_1 = 1, \dots, g_{\mathfrak{c}}\}$. We put

$$G_0(\mathbf{x}) = \{\mathbf{g}_{i,j} = (g_i, \varphi_{i,j}) \mid i = 1, \dots, \mathfrak{c},\ j = 1, \dots, \mathfrak{d}\}.$$

Here $\mathfrak{d} = \#\{\varphi \mid (1, \varphi) \in G(\mathbf{x})\}$ and $\varphi_{1,1} = id$.

Lemma 29.43. *If (ϵ_1, ϵ_2) is sufficiently small, then the function*

$$f(\cdot; [\mathbf{x}', \vec{w}^+]) : \mathfrak{H}(\mathbf{x}', \vec{w}^+) \to \mathbb{R}_{\geq 0}$$

is strictly convex and attains local minima at exactly $\mathfrak{c}$ points $h_1, \dots, h_{\mathfrak{c}}$.

Moreover the following also holds: For each i we have $\mathfrak{d}$ choices of $\vec{w}'^+_{(i,j)}$ and $\rho_{(i,j),1} \in \mathfrak{S}_\ell$ ($j = 1, \dots, \mathfrak{d}$) such that $\rho_{(i,j),1}\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h_i\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(i,j)}$. We put $\Psi(\varphi_{i,j}) = \sigma_{(i,j)} = (\sigma_{(i,j),1}, \sigma_{(i,j),2})$.

Then

$$h_i = h_1 g_i^{-1}, \qquad \rho_{(i,j),1} = \sigma_{(i,j),1}\rho_{(1,1),1}, \qquad w'^+_{(i,j)} = \sigma_{(i,j),2} w'^+_{(1,1)}. \tag{29.29}$$

Proof. Let us first consider the case when $\mathbf{x}' = \mathbf{x}$. In this case it is easy to see that f attains local minimum 0 at $h_i = g_i^{-1}$ ($i = 1, \dots, \mathfrak{c}$) ($g_1 = h_1 = 1$). By taking ϵ_2 appropriately we may assume that there are no other local minima. We can take $\rho_{(1,1),1} = id$, $\rho_{(i,j),1} = \Psi_1(\varphi_{i,j})$. Moreover $\vec{w}'^+_{(1,1)} = \vec{w}^+$, $\vec{w}'^+_{(i,j)} = \varphi_{i,j} \cdot \vec{w}^+$.

By Condition 29.34 there exists $\sigma_{(i,j),2}$ such that $\varphi_{i,j} \cdot \vec{w}^+ = \sigma_{(i,j),2}\vec{w}^+$. We thus proved (29.29) in case $\mathbf{x}' = \mathbf{x}$.

The general case can be reduced to this case in the same way as Lemma 29.21 □

Lemma 29.44. *In the situation of Lemma 29.38, we have*

$$f(h; [\mathbf{x}', \vec{w}^+]) = f(h; [\mathbf{g}\mathbf{x}', \varphi\vec{w}^+]).$$

Here $\mathbf{x}' = (u', \Sigma', \vec{z}'^+)$, $\mathbf{g}\mathbf{x}' = (gu', \Sigma', \Psi_1(\varphi)\vec{z}'^+)$. ($\mathbf{g} = (g, \varphi)$.)

Proof. The same as the proof of Lemma 29.20. □

Remark 29.45. In the situation of Lemma 29.40, $f(h; \mathbf{x}', \vec{w}^+_{(1)})$ may be different from $f(h; \mathbf{x}', \vec{w}^+_{(2)})$.

Suppose $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+$. We define

$$\mathcal{P}_{\mathbf{x}', w^+, w'^+, h} : E(\mathbf{x}) \to C^\infty(\Sigma', u'^*TX \otimes \Lambda^{0,1}) \tag{29.30}$$

in the same way as (29.14).

Definition 29.46. Consider $\mathbf{x}'$ such that $\mathfrak{H}(\mathbf{x}', \vec{w}^+)$ is nonempty for some (hence for all by Lemma 29.40) $\vec{w}^+ \in \mathbf{S}$.

We denote by $h_i^{\vec{w}^+}$ $(i = 1, \dots, \mathfrak{c})$ the elements of $\mathfrak{H}'(\mathbf{x}, \vec{w}^+)$ at which local minima of $f(\cdot; [\mathbf{x}', \vec{w}^+])$ is attained.

Applying Lemma 29.43, we obtain $\rho^{\vec{w}^+}_{(i,j),1}$, $\vec{w}'^{+,\vec{w}^+}_{(i,j)}$ for $j = 1, \dots, \mathfrak{d}$. (We include $\vec{w}^+$ in the notation to show its dependence on $\vec{w}^+$.)

Lemma 29.47.

$$\mathcal{P}_{\rho^{\vec{w}^+}_{(i,j),1}\mathbf{x}', \vec{w}^+, \vec{w}'^{+,\vec{w}^+}_{(i,j)}, h_i^{\vec{w}^+}} = \mathcal{P}_{\rho^{\vec{w}^+}_{(1,1),1}\mathbf{x}', \vec{w}^+, \vec{w}'^{+,\vec{w}^+}_{(1,1)} h_1^{\vec{w}^+}} \circ (\varphi_{i,j})_*.$$

Proof. The proof is the same as the proof of Lemma 29.22. ☐

Lemma 29.48. *For each sufficiently small positive number ϵ_1, the number*

$$\sup_{\vec{w}_1^+, \vec{w}_2^+ \in \mathbf{S}} \sup_{\mathfrak{v} \ne 0} \frac{\Vert \mathcal{P}_{\rho^{\vec{w}_1^+}_{(i,j),1}\mathbf{x}', \vec{w}_1^+, \vec{w}'^{+,\vec{w}_1^+}_{(i,j)}, h_i^{\vec{w}_1^+}}(\mathfrak{v}) - \mathcal{P}_{\rho^{\vec{w}_2^+}_{(i,j),2}\mathbf{x}', \vec{w}_2^+, \vec{w}'^{+,\vec{w}_2^+}_{(i,j)}, h_i^{\vec{w}_2^+}}(\mathfrak{v}) \Vert}{\Vert \mathfrak{v} \Vert}$$

goes to 0 *as* $\epsilon_2 \to 0$.

Proof. From the proof of Lemma 29.40, we have

$$u' \circ i_{\mathbf{v}'_{(1)}} \sim hu \sim u' \circ i_{\mathbf{v}'_{(2)}}.$$

Namely $i_{\mathbf{v}'_{(1)}}$ is C^1 close to $i_{\mathbf{v}'_{(2)}}$.

We can also prove that $h_i^{\vec{w}_1^+}$ is close to $h_i^{\vec{w}_2^+}$ by the proof of Lemma 29.40.

Using these facts, the lemma follows from definition. ☐

We now define

$$\mathcal{P}_{\mathbf{x}'} : E(\mathbf{x}) \to C^\infty(\Sigma', u'^* TX \otimes \Lambda^{0,1})$$

by

$$\mathcal{P}_{\mathbf{x}'}(\mathfrak{v}) = \int_{\vec{w}^+ \in \mathbf{S}} \mathcal{P}_{\rho^{\vec{w}^+}_{(i,j),1}\mathbf{x}', \vec{w}^+, \vec{w}'^{+,\vec{w}^+}_{(i,j)}, h_i^{\vec{w}^+}}((\varphi^{\vec{w}^+}_{i,j})_*^{-1}\mathfrak{v}) d\vec{w}^+. \tag{29.31}$$

We use the Haar measure, that is, the canonical probability measure $d\vec{w}^+$ of $\mathbf{S}$ in the right hand side. Lemma 29.48 implies that $\mathcal{P}_{\mathbf{x}'}$ is injective if ϵ_2 is sufficiently small.

(29.31) is independent of $i = 1, \dots, \mathfrak{c}$, $j = 1, \dots, \mathfrak{d}$ by Lemmas 29.47, 29.16.

Let $\mathbf{g} = (g, \psi) \in G(\mathbf{x})$. We put $\psi(z_a^+) = z^+_{\sigma_1(a)}$ where $\sigma_1 \in \mathfrak{S}_\ell$. For $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ we define $\mathbf{g}\mathbf{x}' = (\Sigma', \sigma_1 \vec{z}'^+, gu')$ with $(\sigma_1 \vec{z}'^+)_a = z'^+_{\sigma_1^{-1}(a)}$. (See (29.2).)

Lemma 29.49. *We have*

$$\mathcal{P}_{\mathbf{g}\mathbf{x}'}(\psi_* \mathfrak{v}) = g_* \mathcal{P}_{\mathbf{x}'}(\mathfrak{v}).$$

Proof. We put $\sigma_2 = \Psi_2(\psi)$. We apply Lemma 29.43 to $\mathbf{x}'$, $\vec{w}^+$ and obtain h_i, $\vec{w}'^+_{(i,j)}$ etc. We apply Lemma 29.43 to $\mathbf{g}\mathbf{x}'$, $\sigma_2 \vec{w}^+$ and obtain $\overline{h}_i$, $\vec{\overline{w}}'^+_{(i,j)}$ etc.

In the same way as the proof of Lemmas 29.15 and 29.20 we can prove

$$\overline{h}_i = h_i g, \qquad \vec{\overline{w}}'^+_{(i,j)} = \sigma_2 \vec{w}'^+_{(i,j)}. \tag{29.32}$$

In fact, if (29.32) holds and

$$u'(w'^+_{(i,j),a}) \in h_i N_{u(w_a^+)},$$

then for $b = \sigma_2(a)$ we have

$$gu'(\overline{w}'^{+}_{(i,j),b}) = gu'(w'^{+}_{(i,j),a}) \in gh_i N_{w_a^+} = \overline{h}_i N_{(\sigma_2 w^+)_b}.$$

This is a part of Definition 29.14 (4). We can use this fact together with (the proof of) Lemma 29.44 to prove (29.32).

We put

$$\overline{\mathbf{v}}'_{(i,j)} = (\Sigma', \sigma_1(\vec{z}'^{+}), \vec{\overline{w}}'^{+}_{(i,j)}).$$

Then by the second formula in (29.32) and (29.11), we obtain

$$i_{\overline{\mathbf{v}}'_{(i,j)}} = i_{\mathbf{v}'_{(i,j)}} \circ \psi^{-1}.$$

Using the first formula in (29.32) and the definition also we have

$$\mathcal{P}_{\mathbf{g}\mathbf{x}',\sigma_1(\vec{w}^{+}),\vec{\overline{w}}'^{+}_{(i,j)},\overline{h}_i} = g_* \circ \mathcal{P}_{\mathbf{x}',\vec{w}^{+},\vec{w}'^{+}_{(i,j)},h_i} \circ \psi_*^{-1}.$$

(Here we assume $\rho_{(i,j),1} = id$ for simplicity.)

The lemma follows by taking the average over $\vec{w}^{+}$. □

Remark 29.50. Note we integrate over $\vec{w}^{+}$ to achieve the above $G(\mathbf{x})$ invariance. This is different from the situation of Lemma 29.47. Namely in that case elements of $G(\mathbf{x})$ only exchange the points of $\vec{w}^{+}$. In our situation $G(\mathbf{x})$ has positive dimension so we can not take (the finite set) $\vec{w}^{+}$ so that it is $G(\mathbf{x})$ invariant (as a set).

Lemma 29.51. *If $h \in T_0^{n-1}$, then*

$$\mathcal{P}_{h\mathbf{x}'} = h_* \circ \mathcal{P}_{\mathbf{x}'}.$$

Proof. This is immediate from the construction. □

We put $E(\mathbf{x}') = \mathrm{Im}\mathcal{P}_{\mathbf{x}'}$. By construction, it is independent of the permutation of interior marked points.

We next prove its S_0^1-invariance. We put $\mathbf{x}' = (\Sigma', \vec{z}'^{+}, u')$. Let $S \in S_0^1$. Note $S_* : u'^*TX \to (S \circ u')^*TX$ induces

$$S_* : C^\infty(\Sigma', u'^*TX \otimes \Lambda^{0,1}) \to C^\infty(\Sigma', (S \circ u')^*TX \otimes \Lambda^{0,1}).$$

Lemma 29.52. *We have:*

$$E(S\mathbf{x}') = S_* E(\mathbf{x}').$$

Proof. Let $\widehat{S} = (S, \widehat{\varphi}) \in G(\mathbf{x})$ which is a lift of S. We choose a path $\widehat{S}_t = (S_t, \widehat{\varphi}_t)$ in $G(\mathbf{x})$ such that $\widehat{S}_0 = (1,1)$ and $\widehat{S}_1 = \widehat{S}$.

We remark that we may take $\widehat{\varphi}_t$ so that $\widehat{\varphi}_t(z_a^+) = z_a^+$, for each a. (In general $\widehat{\varphi}(z_a) = z_b$, $b \neq a$ may occur.) In fact we take $\mathfrak{t} \in Lie(S_0^1)$ such that $\exp(\mathfrak{t}) = \rho$. $\mathfrak{t}$ induces a holomorphic vector field on Σ which is zero at $\vec{z}^{+}$ and at the singular points. By integrating it we obtain the desired $\widehat{\varphi}_t$.

We take $h_{i,t} \in T_0^{n-1}$, $\vec{w}'^{+}_{(i,j),t}$, $\varphi_{i,j,t}$ ($i = 1, \dots, \mathfrak{c}$) such that $\widehat{\varphi}_t\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(h_{i,t}\mathbf{x}, \widehat{\varphi}_t\vec{w}^{+})$ with respect to $\vec{w}'^{+}_{(i,j),t}$. We remark

$$(\widehat{\varphi}_t\vec{w}^{+})_a = \widehat{\varphi}_t(w_a). \tag{29.33}$$

We have

$$\begin{aligned}
f(h; S_t\mathbf{x}', \widehat{\varphi}_t\vec{w}^+) &= \sum_a \mathrm{dist}_{hN_{\widehat{\varphi}_t(w_a^+)}}(S_t u'(w'^+_{(i,j),t;a}), hu(\widehat{\varphi}_t w_a^+))^2 \\
&= \sum_a \mathrm{dist}_{S_t hN_{w_a^+}}(S_t u'(w'^+_{(i,j),t;a}), hS_t u(w_a^+))^2 \\
&= \sum_a \mathrm{dist}_{hN_{w_a^+}}(u'(w'^+_{(i,j),t;a}), hu(w_a^+))^2 \\
&= f(h; \mathbf{x}', \vec{w}^+).
\end{aligned}$$


Σ'
Σ
$w'_{(i,j),a}$
$w'_{(i,j),t,a}$
w_a^+
$\hat{\varphi}_t$
$\hat{\varphi}_t(w_a^+)$
$S_t u'$
u'
u'
hu
hu
$u'(\Sigma')$
$hu(\Sigma) = S_t h(\Sigma)$
$S_t u'(\Sigma')$
$hu(w_a^+)$
$hN_{w_a^+}$
$u'(w'_{(i,j),a})$
S_t
$S_t u'(w'_{(i,j),a})$
$S_t hN_{w_a^+} = hN_{\hat{\varphi}_t(w_a^+)}$
$hu(\hat{\varphi}_t(w_a^+)) = hS_t u(w_a$

Figure 29.3

Remark 29.53. Note that $S_t u'(w'^{+}_{(i,j),t;a}) = u'(\widehat{\varphi}_t(w'^{+}_{(i,j),t;a}))$ does *not* hold in general. (Actually the right hand side is not defined.) On the other hand, $S_t u(w_a^+) = u(\widehat{\varphi}_t(w_a^+))$ does hold. Moreover, $S_t u'(w'^{+}_a) = S_t u'(w'^{+}_{(i,j),t;a})$ holds, because both are the unique nearby intersection point of $S_t u'(\Sigma') \cap h S_t N_{w_a^+}$. (In other words, $w'^{+}_a = w'^{+}_{(i,j),t;a}$.)

Therefore we find that $h_{i,t}$ is independent of t and is equal to $h_i = h_{i,0}$. We can use this fact in the same way as the proof of Lemma 29.47 to show

$$\mathcal{P}_{S\mathbf{x}',\widehat{\varphi}\vec{w}^+,\Psi_2(\varphi)\vec{w}'^+,h_i} = S_* \circ \mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,h_i}.$$

The lemma follows by taking the average over $\vec{w}^+$. □

We thus proved that $E(\mathbf{x}')$ is invariant under the T^n and $G(\mathbf{x})$ actions.

We consider the set of $\mathbf{x}'$ satisfying

$$\overline{\partial} u' \in E(\mathbf{x}')$$

together with E and

$$s(\mathbf{x}') = \overline{\partial} u'.$$

This defines a Kuranishi neighborhood of the T^n orbit of $\mathbf{x}$.

In the same way as before we can extend it and define $E(\mathbf{x}')$ for $\mathbf{x}'$ in a neighborhood of the $T^n \times \mathfrak{S}_\ell$ orbit of $\mathbf{x}$. Then $E(\mathbf{x})$ is invariant under the T^n-action and the permutation of interior marked points.

We thus completed the construction of our Kuranishi neighborhood in the case when $\mathbf{x}$ has only one disc component.

29.2. **The case with more than one disk components.**

We finally consider the case when the domain may have 2 or more disk components.

Let $\mathbf{x} = (\Sigma, \vec{z}^+, u) \in \mathcal{M}_{0;\ell}(\beta)$. We decompose Σ into disk components (we include tree of sphere components to the disc component on which it is rooted). We write

$$\Sigma = \bigcup_{\alpha \in \mathfrak{A}} \Sigma(\alpha).$$

Each of the disk component together with the restriction of u and $\vec{z}^+$ induces an element

$$\mathbf{x}(\alpha) = (\Sigma(\alpha), \vec{z}^+_\alpha, u_\alpha) \in \mathcal{M}_{0;\ell_\alpha}(\beta(\alpha)).$$

Each of $\mathbf{x}(\alpha)$ satisfies one of the following three conditions.

1. $\mathbf{x}(\alpha)$ satisfies Condition 29.9.
2. $\mathbf{x}(\alpha)$ satisfies the conclusion of Lemma 29.31 (1).
3. $\mathbf{x}(\alpha)$ satisfies the conclusion of Lemma 29.31 (2).

We say that $\mathbf{x}(\alpha)$ is *of Type I, II, III* if it satisfies (1), (2), (3) above, respectively. Sometime we say α is of Type I, II, III also.

We take and fix a splitting of T^n as in Lemma 29.30. We write this splitting as $S^1_\alpha \times T^{n-1}_\alpha$. (This splitting depends on α (which is of Type III) so we include α in the notation.)

Hereafter we put

$$\mathbf{T}_\alpha = \begin{cases} T^n & \text{if } \alpha \text{ is of Type I or II,} \\ T^{n-1}_\alpha & \text{if } \alpha \text{ is of Type III.} \end{cases} \tag{29.34}$$

For each $\mathbf{x}(\alpha)$ of Type I or II, we take extra marked points w_α^+ so that Condition 29.10 or 29.32 is satisfied respectively.

For each $\mathbf{x}(\alpha)$ of Type III, we choose the union $\mathbb{S}_\alpha \subset \Sigma$ of circles as in Lemma 29.33. We then take $\mathbf{S}_\alpha$ as in (29.28). ($(\mathbf{S},\mathbf{x})$ in (29.28) corresponds to $(\mathbf{S}_\alpha,\mathbf{x}(\alpha))$ here.) We use it to define $\mathbf{S}$ as follows.

We defined the group $G(\mathbf{x})$ as in Definition 29.2. We remark that in our situation where the domain Σ has at least two disk components, $G(\mathbf{x})$ is necessarily a finite group. This is because the isotropy group of a boundary singular point is trivial. (See the proof of Lemma 29.31.)

Remark 29.54. An element $\mathbf{g} \in G(\mathbf{x})$ may exchange factors $\mathbf{x}(\alpha)$. We choose $\vec{w}_\alpha^+$ (the case of Type I and Type II component), $\mathbf{S}_\alpha$ (the case of Type III component) so that it is preserved under this action. In fact, we are proving Proposition 29.5 inductively. (Namely by induction on $E(\beta)$.) In other words, we are making the choices of $\vec{w}_\alpha^+$ or $\mathbf{S}_\alpha$ inductively on $E(\beta_\alpha)$. In our situation for which β is decomposed into $\beta = \sum \beta_\alpha$, those choices for each of β_α had already been made. Of course, it depends only on the isomorphism classes. So the action of $\mathbf{g}$ automatically preserves it.

We define

$$G_{00}(\mathbf{x}) = \{(g,\varphi) \in G(\mathbf{x}) \mid g \in \mathbf{T}_\alpha \text{ for all } \alpha\}.$$

This is also a finite group.

Let $\mathfrak{A}(\mathrm{III})$ be the set of all $\alpha \in \mathfrak{A}$ of Type III. We define an equivalence relation $\sim$ on $\mathfrak{A}(\mathrm{III})$ as follows: Let $\alpha,\alpha' \in \mathfrak{A}(\mathrm{III})$. Then $\alpha \sim \alpha'$ if there exists $(g,\varphi) \in G_{00}(\mathbf{x})$ such that $\varphi(\Sigma(\alpha)) = \Sigma(\alpha')$.

Let $(\vec{w}_\alpha^+)_{\alpha\in\mathfrak{A}(\mathrm{III})}$ and consider the following condition on it.

Condition 29.55.
1. Each of $\vec{w}_\alpha^+$ satisfies Condition 29.34. (Here we replace $\mathbf{x}$ in Condition 29.34 by $\mathbf{x}(\alpha)$.)
2. Let $\alpha \sim \alpha'$ and $\varphi(\Sigma(\alpha)) = \Sigma(\alpha')$ as above. Then there exists $\rho \in \mathfrak{S}_{m_\alpha}$ such that
$$\varphi(w_{\alpha,a}^+) = w_{\alpha',\rho(a)}^+.$$

We remark that if $\alpha \sim \alpha'$ then $\mathbf{T}_\alpha = \mathbf{T}_{\alpha'}$, $S_\alpha^1 = S_{\alpha'}^1$.

Definition 29.56. The set $\mathbf{S}$ is a subset of $\prod_{\alpha\in\mathfrak{A}(\mathrm{III})} \mathbf{S}_\alpha$ consisting of the elements $(\vec{w}_\alpha^+)_{\alpha\in\mathfrak{A}(\mathrm{III})}$ satisfying Condition 29.55.

Since we have already fixed $\vec{w}_\alpha^+$ if α is of Type I or II, we obtain $(\vec{w}_\alpha^+)_{\alpha\in\mathfrak{A}}$ for each element of $\mathbf{S}$. Hereafter we regard $(\vec{w}_\alpha^+)_{\alpha\in\mathfrak{A}} \in \mathbf{S}$.

Lemma 29.57. $\mathbf{S}$ *is invariant under the* $G(\mathbf{x})$ *action.*

Proof. This is a consequence of Condition 29.34 (3) and the fact that $G_{00}(\mathbf{x})$ is a normal subgroup of $G(\mathbf{x})$. (See also Remark 29.54.) ☐

For $(\vec{w}_\alpha^+)_{\alpha\in\mathfrak{A}} \in \mathbf{S}$, we consider

$$\mathbf{v}(\vec{w}^+) = \left(\Sigma, \vec{z}^+ \cup \bigcup_\alpha \vec{w}_\alpha^+\right) = \left(\Sigma, \vec{z}^+ \cup \vec{w}^+\right).$$

We put $\ell = \sum \ell_\alpha = \#\vec{z}^+$, $m = \sum m_\alpha = \#\vec{w}^+$.

We also define $\mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$ as in Definition 29.11. Let $\varphi \in \mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$. We then obtain $\rho = (\rho_1, \rho_2) = \Psi(\varphi) \in \mathfrak{S}_\ell \times \mathfrak{S}_m$. We remark that the image $\Psi(\mathrm{Aut}_+(\mathbf{v}(\vec{w}^+)))$ depends on $\vec{w}^+$ in general. However if $\mathbf{g} = (g, \varphi) \in G(\mathbf{x})$, then

$$\Psi(\mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))) = \rho\Psi(\mathrm{Aut}_+(\mathbf{g}\mathbf{v}(\vec{w}^+)))\rho^{-1}$$

where $\mathbf{g}\mathbf{v}(\vec{w}^+)) = (\Sigma, \varphi \cdot \vec{z}^+ \cup \varphi \cdot \vec{w}^+)$, $\rho = \Psi(\varphi)$.

We remark that if $(g, \varphi) \in G_{00}(\mathbf{x})$ then $\varphi \in \mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$ by Condition 29.55 (2).

We also choose N_w as in Condition 29.13 or 29.36.

We consider a neighborhood $\mathfrak{U}(\mathbf{v}(\vec{w}^+))$ of $\mathbf{v}(\vec{w}^+)$. Let $\mathbf{v}' \in \mathfrak{U}(\mathbf{v}(w^+))$ and $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+)$. Then we have $\mathcal{U}(S(\Sigma))$, a neighborhood of singular point sets of Σ, and

$$i_{\mathbf{v}'} : \Sigma \setminus \mathcal{U}(S(\Sigma)) \to \Sigma'$$

such that the following holds:

Condition 29.58. (1) $i_{\mathbf{v}'}(z_i^+) = z_i'^+$, $i_{\mathbf{v}'}(w_i^+) = w_i'^+$.

(2) $i_{\mathbf{v}'}$ is a diffeomorphism to its image.
(3) $i_{\mathbf{v}'}$ is biholomorphic on each of the support of $E(\mathbf{x}(\alpha))$.
(4) If $\sigma = (\sigma_1, \sigma_2) \in \Psi(\psi)$, $\psi \in \mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$ then

$$i_{\sigma\mathbf{v}'} = i_{\mathbf{v}'} \circ \psi^{-1} : \Sigma \setminus \mathcal{U}(S(\Sigma)) \to \Sigma'.$$

(5) When Σ has 2 or more disk components, $i_{\mathbf{v}'}$ is compatible with the maps inductively defined in each of the disk components.
(6) $i_{\mathbf{v}'}$ depends continuously on $\mathbf{v}'$ in C^∞ topology.

We will elaborate Condition (5) later during the proof of Lemma 29.71.

Let ϵ_1, ϵ_2 be small positive numbers. We take $(\Sigma', \vec{z}'^+)$ and $u' : (\Sigma', \partial\Sigma') \to (X, L)$.

Definition 29.59. For $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ and $\vec{w}^+$, we say that $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ is (ϵ_1, ϵ_2)-*close to* $(h\mathbf{x}, \vec{w}^+)$ with respect to $\vec{w}'^+$ if the following holds.

(1) $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+) \in \mathfrak{U}(\mathbf{v}(\vec{w}^+))$ and $\mathrm{dist}(\mathbf{v}', \mathbf{v}(\vec{w}^+)) < \epsilon_1$. Here we use an appropriate metric on $\mathcal{M}_{0;\ell+m}$ to define the distance between $\mathbf{v}'$ and $\mathbf{v}(\vec{w}^+)$.
(2) The C^1 distance between two maps $u' \circ i_{\mathbf{v}'}$ and hu is smaller than ϵ_1 on $\Sigma \setminus \mathcal{U}(S(\Sigma))$.
(3) $\mathrm{diam}(u'(\mathcal{S})) < \epsilon_1$ if $\mathcal{S}$ is any connected component of the complement $\Sigma' \setminus i_{\mathbf{v}'}(\Sigma \setminus \mathcal{U}(S(\Sigma)))$.
(4) $u'(w_a'^+) \in hN_{w_a^+}$ for each a and

$$\mathrm{dist}(u'(w_a'^+), hu(w_a^+)) < \epsilon_2.$$

Here $\vec{w}_a'^+ = (w_a'^+)_{a\in A}$.

We remark that this definition is exactly the same as Definition 29.14.

We can proceed in the same way as the case of Type I component using Condition 29.59 in place of Definition 29.14. However, then, the resulting Kuranishi chart will not be disk-component-wise. To obtain a disk-component-wise Kuranishi chart we need to consider the case where the element $h \in T^n$ above may vary depending on the disk component to which h is applied. For this purpose we state the next definition.

We denote

$$\vec{w}^+ = \bigcup_{\alpha\in\mathfrak{A}} \vec{w}^+_\alpha = \bigcup_{\alpha\in\mathfrak{A}} (w^+_{\alpha,a})_{a\in A_\alpha}.$$

We also write $\vec{w}^+ = (w^+_a)_{a\in A}$ sometimes.

In the next definition we consider $\rho = (\rho_1, \rho_2) \in \mathfrak{S}_\ell \times \mathfrak{S}_m$ which is induced by

$$\rho_0 \in \mathfrak{S}_{\#\mathfrak{A}}, \quad \rho_{0,1}, \rho_{0,2}$$

according to the formula

$$\rho_1(\alpha, a) = (\rho_0(\alpha), \rho_{0,1}(\alpha, a)), \quad \rho_2(\alpha, a) = (\rho_0(\alpha), \rho_{0,2}(\alpha, a)). \tag{29.35}$$

Here $\rho_{0,1}$ is a map from $\{(\alpha, a) \mid \alpha \in \mathfrak{A}, a = 1, \dots, \ell_\alpha\}$ to $\{1, \dots, \ell_{\rho_0(\alpha)}\}$ and $\rho_{0,2}$ is a map from $\{(\alpha, a) \mid \alpha \in \mathfrak{A}, a = 1, \dots, m_\alpha\}$ to $\{1, \dots, m_{\rho_0(\alpha)}\}$. (Actually $\ell_{\rho_0(\alpha)} = \ell_\alpha$ and $m_{\rho_0(\alpha)} = m_\alpha$.)

In such a case we write

$$\rho = (\rho_1, \rho_2) = [\rho_0, \rho_{0,1}, \rho_{0,2}].$$

In case $\rho = \Psi(\varphi)$ for $\varphi \in \mathrm{Aut}_+(\mathbf{v}(\vec{w}^+))$ the permutation ρ is of the form (29.35). In fact ρ_0 is determined by

$$\varphi(\Sigma(\alpha)) = \Sigma(\rho_0(\alpha)).$$

We also remark the following. Suppose $[\rho_0, \rho_{0,1}, \rho_{0,2}] = \Psi(\varphi)$ and $\rho_0(\alpha) = \alpha$. Then $(\rho_{0,1}(\alpha, \cdot), \rho_{0,2}(\alpha, \cdot))$ is an element of $\mathfrak{S}_{\ell_\alpha} \times \mathfrak{S}_{m_\alpha}$. Moreover it lies in the image of $\mathrm{Aut}_+(\mathbf{x}(\alpha))$.

Definition 29.60. Let $(h(\alpha))_{\alpha\in\mathfrak{A}}$ with $h(\alpha) \in \mathbf{T}_\alpha$. We say that $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ is *weakly* (ϵ_1, ϵ_2)*-close to* $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+$ if the following holds.

1. $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+) \in \mathfrak{U}(\mathbf{v}(\vec{w}^+))$ and $\mathrm{dist}(\mathbf{v}', \mathbf{v}(\vec{w}^+)) < \epsilon_1$. Here we use an appropriate metric on $\mathcal{M}_{0;\ell+m}$ to define the distance between $\mathbf{v}'$ and $\mathbf{v}(\vec{w}^+)$.
2. The C^1 distance between two maps $u' \circ i_{\mathbf{v}'}$ and $h(\alpha)u$ is smaller than ϵ_1 on $\Sigma(\alpha) \setminus \mathcal{U}(S(\Sigma))$.
3. $\mathrm{diam}(u'(\mathcal{S})) < \epsilon_1$ if $\mathcal{S}$ is any connected component of the complement $\Sigma' \setminus i_{\mathbf{v}'}(\Sigma \setminus \mathcal{U}(S(\Sigma)))$.
4. $u'(w'^+_{\alpha,a}) \in h(\alpha)N_{w^+_{\alpha,a}}$ for each α, a and
$$\mathrm{dist}(u'(w'^+_{\alpha,a}), h(\alpha)u(w^+_{\alpha,a})) < \epsilon_2.$$
5. $|h - h(\alpha)| < \epsilon_1$ if $\mathbf{x}(\alpha)$ is of Type I or II, and $|\pi(h) - h(\alpha)| < \epsilon_1$ if $\mathbf{x}(\alpha)$ is of Type III. Here $\pi : T^n \to T^{n-1}_\alpha$ is the projection.

Definitions 29.59 and 29.60 are related in the following way.

Lemma 29.61. *For each* $\vec{\epsilon} = (\epsilon_1, \epsilon_2)$ *there exists* $(\epsilon^0_1(\vec{\epsilon}), \epsilon^0_2(\vec{\epsilon}))$ *such that the following holds for* $\epsilon'_1 < \epsilon^0_1(\vec{\epsilon})$*,* $\epsilon'_2 < \epsilon^0_2(\vec{\epsilon})$*.*

1. *If* $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ *is* (ϵ_1, ϵ_2)*-close to* $(h\mathbf{x}, \vec{w}^+)$ *with respect to* $\vec{w}'^+$*, then* $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ *is weakly* (ϵ_1, ϵ_2)*-close to* $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$*. Here* $h(\alpha) = h$ *for all* α*.*
2. *If* $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ *is weakly* $(\epsilon'_1, \epsilon'_2)$*-close to* $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$*, then there exists* $\vec{w}''^+$ *such that* $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ *is* (ϵ_1, ϵ_2)*-close to* $(h\mathbf{x}, \vec{w}^+)$ *with respect to* $\vec{w}''^+$*.*

Proof. The proof is immediate from definitions. □

We will consider the situation of Definition 29.60.

We can generalize Lemma 29.39 in our present situation as follows.

Lemma 29.62. *Let $\mu_1 \in \mathfrak{S}_\ell$ and ϵ_1, ϵ_2 be sufficiently small.*

Suppose $\mathbf{x}'$ is weakly (ϵ_1, ϵ_2)-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(1)}$ and $\mu_1\mathbf{x}'$ is weakly (ϵ_1, ϵ_2)-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(2)}$.

Then there exists $(1, \psi) \in G_{00}(\mathbf{x}) \subset G(\mathbf{x})$ such that $\rho_{(2)} = \rho_{(1)}\mu$, $\mu = (\mu_1, \mu_2) = \Psi(\psi)$ and $\vec{w}'^+_{(2)} = \mu_2\vec{w}'^+_{(1)}$.

The proof is the same as that of Lemma 29.16.

Let α_0 be of Type I or II. We define the set $\mathfrak{H}(\mathbf{x}', \alpha_0, \vec{w}^+)$ of all $h(\alpha_0) \in T^n$ with the following properties: There exist $\vec{w}'^+$ and $h(\alpha) \in T^n$ for $\alpha \neq \alpha_0$ such that $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+$. We define a function

$$f_{\alpha_0}(\cdot; \mathbf{x}', \vec{w}^+) : \mathfrak{H}(\mathbf{x}', \alpha_0, \vec{w}^+) \to \mathbb{R}$$

by

$$f_{\alpha_0}(h(\alpha_0); \mathbf{x}', \vec{w}^+) = \sum_{a\in A_{\alpha_0}} \mathrm{dist}(u'(w'^+_{\alpha_0,a}), h(\alpha_0)u(w^+_{\alpha_0,a}))^2. \tag{29.36}$$

Lemma 29.63. *The right had side of (29.36) is independent of $\vec{w}^+_\alpha$ $(\alpha \neq \alpha_0)$ and $\vec{w}'^+$.*

Proof. When we fix $\vec{w}^+_\alpha$ (for $\alpha \neq \alpha_0$ also) but change $\vec{w}'^+$, the required invariance is a consequence of Lemma 29.62.

When we change $w^+_{\alpha,a}$ for $\alpha \neq \alpha_0$, the element $w'^+_{\alpha,a}$ may change but we may choose $w'^+_{\alpha_0,a}$ so that $u'(w'^+_{\alpha_0,a})$ does not change.

The lemma easily follows from those facts. □

If α_0 is of Type I, then f_{α_0} is strictly convex.

If α_0 is of Type II, we use special points on the component $\Sigma(\alpha_0)$ and define f_{α_0+} in the same way as (29.26) and can show that it is strictly convex.

We now consider the case of α of Type III. We first modify Lemma 29.40 for the present situation. Let us consider $\mathbf{x}' = (\Sigma', \vec{z}'^+, u')$ and $\vec{w}^+_{(k)} \in \mathbf{S}$ for $k = 1, 2$.

Lemma 29.64. *For each $\vec{\epsilon} = (\epsilon_1, \epsilon_2)$ there exists $(\epsilon^0_1(\vec{\epsilon}), \epsilon^0_2(\vec{\epsilon}))$ such that the following holds for $\epsilon'_1 < \epsilon^0_1(\vec{\epsilon})$, $\epsilon'_2 < \epsilon^0_2(\vec{\epsilon})$.*

Suppose $\mathbf{x}'$ is weakly $(\epsilon'_1, \epsilon'_2)$-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+_{(1)})$ with respect to $\vec{w}'^+_{(1)}$. Then there exists $\vec{w}'^+_{(2)}$ such that $\mathbf{x}'$ is weakly (ϵ_1, ϵ_2)-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{S}}, \vec{w}^+_{(2)})$ with respect to $\vec{w}'^+_{(2)}$.

Proof. The proof is the same as the proof of Lemma 29.40. □

For each choice of $\vec{w}^+$, α_0 of Type III, we define the set $\mathfrak{H}_0(\mathbf{x}', \alpha_0, \vec{w}^+)$ of all $h(\alpha_0) \in T^{n-1}_{\alpha_0}$ with the following properties: There exists $\vec{w}'^+$ and $h(\alpha) \in T^n$ for $\alpha \neq \alpha_0$ such that $\mathbf{x}'$ is (ϵ_1, ϵ_2)-close to $(\mathbf{x}, h, (h(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+$.

We then define

$$f_{\alpha_0} : \mathfrak{H}_0(\mathbf{x}', \alpha_0, \vec{w}^+) \to \mathbb{R}$$

by the same formula as (29.36).

Lemma 29.63 holds in the case when α is of Type II or III also.

We now put

$$\mathfrak{H}_0(\mathbf{x},\vec{w}^+) = \left\{ (h(\alpha))_{\alpha\in\mathfrak{A}} \;\middle|\; \begin{array}{l} h(\alpha) \in \mathfrak{H}(\mathbf{x},\alpha) \text{ if } \alpha \text{ is of Type I or II,} \\ h(\alpha) \in \mathfrak{H}(\mathbf{x},\alpha,\vec{w}^+) \text{ if } \alpha \text{ is of Type III,} \\ \exists h \ (h(\alpha))_{\alpha\in\mathfrak{A}}, h \text{ satisfies (5) of Definition 29.60.} \end{array} \right\}$$

Each of f_α, $f_{\alpha,+}$ may be regarded as a function on it. We also put

$$\mathfrak{H}(\mathbf{x},\vec{w}^+) = \bigcup_{(\rho_1,\rho_2)=[\rho_0,\rho_{0,1},\rho_{0,1}]\in\mathfrak{S}_\ell} \mathfrak{H}_0(\rho_1\mathbf{x},\vec{w}^+).$$

We next adapt Lemma 29.21 to the present situation. We put

$$|G_{00}(\mathbf{v})| = \{g \mid \exists\varphi \ (g,\varphi)\in G_{00}(\mathbf{v})\}.$$

Let $\mathfrak{c}$ be the order of $|G_{00}(\mathbf{v})|$.

Let $|G_{00}(\mathbf{x})| = \{g_1 = 1,\dots,g_{\mathfrak{c}}\}$. We put

$$G_{00}(\mathbf{x}) = \{\mathbf{g}_{i,j} = (g_i,\varphi_{i,j}) \mid i = 1,\dots,\mathfrak{c},\ j = 1,\dots,\mathfrak{d}\}.$$

Here $\mathfrak{d} = \#\{\varphi \mid (1,\varphi)\in G(\mathbf{x})\}$ and $\varphi_{1,1} = id$. We put

$$\Psi(\varphi_{i,j}) = [\sigma_{(i,j),0},\sigma_{(i,j),0,1},\sigma_{(i,j),0,2}] = (\sigma_{(i,j),1},\sigma_{(i,j),2}).$$

Lemma 29.65. *Assume $\mathfrak{H}(\mathbf{x},\vec{w}^+) \neq \emptyset$. Then, there exist exactly $\mathfrak{c}$ choices of $(h_i(\alpha))_{\alpha\in\mathfrak{A}}$ $i = 1,\dots,\mathfrak{c}$ such that all of f_α and $f_{\alpha,+}$ attain local minimum there.*

For each i, there exist $h_i \in T^n$ and $\mathfrak{d}$ choices of $\vec{w}'^+_{(i,j)}$ and

$$\rho_{(i,j),1} = [\rho_{(i,j),0},\rho_{(i,j),0,1}] \in \mathfrak{S}_\ell, \quad j = 1,\dots,\mathfrak{d}$$

such that $\rho_{(i,j),1}\mathbf{x}'$ is weakly (ϵ_1,ϵ_2)-close to $(\mathbf{x},h_i,(h_i(\alpha))_{\alpha\in\mathfrak{A}},\vec{w}^+)$ with respect to $\vec{w}'^+_{(i,j)}$. Let $\varphi_{i,j}(z^+_a) = z^+_{\sigma_{(i,j),1}(a)}$, $\varphi_{i,j}(w^+_a) = w^+_{\sigma_{(i,j),2}(a)}$.

Then we have

$$h_i(\alpha) = g_i^{-1}h_1(\alpha), \quad \rho_{(i,j),1} = \sigma_{(i,j),1}\rho_{(1,1),1}, \quad \vec{w}'^+_{(i,j)} = \sigma_{(i,j),2}\vec{w}'^+_{(1,1)}. \tag{29.37}$$

Proof. In the case when $\mathbf{x}' = \mathbf{x}$, we can prove that f_α (or $f_{\alpha,+}$) takes local minimum 0 at $h_i(\alpha) = g_i^{-1}$ and $\vec{w}'^+_{(i,j)} = \varphi_{i,j}(\vec{w}'^+_{(1,1)})$ by using Condition 29.55. We may choose ϵ_1,ϵ_2 so that there is no other local minimum. This implies the lemma for the case $\mathbf{x}' = \mathbf{x}$.

The general case can be reduced to this case by using the continuity argument similar to the proof of Lemma 29.21. ☐

Lemma 29.66. *For each given ϵ and $\vec{\epsilon} = (\epsilon_1,\epsilon_2)$, there exists $(\epsilon^0_1(\vec{\epsilon},\epsilon),\epsilon^0_2(\vec{\epsilon},\epsilon))$ such that the following holds for all $\epsilon'_1 < \epsilon^0_1(\vec{\epsilon},\epsilon)$, $\epsilon'_2 < \epsilon^0_2(\vec{\epsilon},\epsilon)$.*

If $\mathbf{x}' = (\Sigma',\vec{z}'^+,u')$ is weakly $(\epsilon'_1,\epsilon'_2)$-close to $(\mathbf{x},h',(h'(\alpha))_{\alpha\in\mathfrak{A}},\vec{w}^+)$ with respect to $\vec{w}'^+$, then there exists $(h(\alpha))_{\alpha\in\mathfrak{A}} \in \mathfrak{H}(\mathbf{x},\vec{w}^+)$ such that f_α (or $f_{\alpha,+}$) attains local minimum at $h(\alpha)$ for each α and $|h'(\alpha) - h(\alpha)| < \epsilon$.

Proof. If $\mathbf{x}' = \mathbf{x}$, the proof is immediate from Lemma 29.62. Then the general case follows from continuity argument (using strict convexity of f_α) in a way similar to the last part of the proof of Lemma 29.21. ☐

We now define the maps

$$\mathcal{P}_{\rho_{(i,j),1}\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(i,j)},\alpha} : E(\mathbf{x}(\alpha)) \to C^\infty_0(\Sigma',u'^*TX\otimes\Lambda^{0,1}). \tag{29.38}$$

Here the right hand side denotes the set of smooth sections whose supports are disjoint from the neighborhood $\mathcal{U}(S(\Sigma))$ of singular points of Σ.

We take and fix $\vec{w}^+$. We assume that f_α (or $f_{\alpha,+}$) assumes local minimum at $(h_i(\alpha))_{\alpha\in\mathfrak{A}} \in \mathfrak{H}(\mathbf{x},\vec{w}^+)$. We remark that $\rho_{(i,j),1}\mathbf{x}'$ be weakly (ϵ_1,ϵ_2)-close to $(\mathbf{x}, h_i, (h_i(\alpha))_{\alpha\in\mathfrak{A}}, \vec{w}^+)$ with respect to $\vec{w}'^+_{(i,j)}$.

We put $\mathbf{v}'_{(i,j)} = (\Sigma', \rho_{(i,j),1}\vec{z}'^+, \vec{w}'^+_{(i,j)})$. We decompose the image $i_{\mathbf{v}'}$ into $\cup_\alpha \Sigma'(\alpha)$ where

$$\Sigma'(\alpha) = \mathrm{Im}\ i_{\mathbf{v}'}(\Sigma_0(\alpha))$$

and $\Sigma_0(\alpha)$ is the intersection of the domain of $i_{\mathbf{v}'_{(i,j)}}$ and $\Sigma(\alpha)$.

Let $x \in \Sigma_0(\alpha)$. We have

$$(u^*TX)_x = T_{u(x)}X \overset{h_i(\alpha)_*}{\longrightarrow} T_{h_i(\alpha)u(x)}X \overset{\mathrm{Pal}_{h_i(\alpha)u(x),u'(i_{\mathbf{v}'_{(i,j)}}(x))}}{\longrightarrow} T_{u'(i_{\mathbf{v}'_{(i,j)}}(x))}X. \tag{29.39}$$

We also have:

$$\Lambda^{0,1}_x(\Sigma) \overset{(i^{-1}_{\mathbf{v}'_{(i,j)}})^*}{\longrightarrow} \Lambda^{0,1}_{u'(i_{\mathbf{v}'_{(i,j)}}(x))}(\Sigma'). \tag{29.40}$$

They induce (29.38).

We next prove that the image of the map $\mathcal{P}_{\rho_{(i,j),1}\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(i,j)},\alpha}$ is independent of $\vec{w}'^+_{(i,j)}$.

Lemma 29.67. *In the situation of Lemma 29.65, we have*

$$\mathcal{P}_{\rho_{(i,j),1}\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(i,j)},\alpha} = \mathcal{P}_{\rho_{(1,1),1}\mathbf{x}',\vec{w}^+,\vec{w}'^+_{(1,1)},\sigma_{(i,j),0}(\alpha)} \circ (\varphi_{i,j})_*.$$

Proof. Using Lemma 29.65, the proof is the same as that of Lemma 29.22 □

Remark 29.68. In general, $\mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+,\alpha}$ depends on $\vec{w}^+_{\alpha'}$ for $\alpha' \neq \alpha$. The element $h(\alpha)$ is independent of $\vec{w}'^+_{\alpha'}$ for $\alpha' \neq \alpha$. However the map $i_{\mathbf{v}'}$ may depend on $\vec{w}'^+$. In particular, it may depend on $\vec{w}'^+_{\alpha'}$ for $\alpha' \neq \alpha$. We will come back to this point in the course of the proof of Lemma 29.71.

Let us move $\vec{w}^+ \in \mathbf{S}$ in a one-parameter family $\vec{w}^+_s$. Here $s \in S^1$. We take $h_s(\alpha)$, $\vec{w}'^+_s$ for each $\vec{w}^+_s$ so that they continuously depend on $s \in (0,1)$. (Here we identify $S^1 \setminus \{\text{point}\} = (0,1)$.) Then we can use (the proof of) Lemma 29.40 to show that $\mathcal{P}_{\mathbf{x}',\vec{w}^+_s,\vec{w}'^+_s,\alpha}$ is close to the constant map. Therefore we use Lemma 29.65 to prove that $\vec{w}'^+_s$ can be extended to a family parametrized by $s \in S^1$. (We use the fact that the group $G_{00}(\mathbf{x})$ is a finite group here.)

We use this fact to show that there exists a $\vec{w}^+ \in \mathbf{S}$ parametrized *continuous* family of $\vec{w}'^+$. We denote it by $\vec{w}'^+_{\vec{w}^+}$.

We now take the average over $\vec{w}^+$ and define $\mathcal{P}_{\mathbf{x}',\alpha}$ by

$$\mathcal{P}_{\mathbf{x}',\alpha} = \int_{\vec{w}^+\in\mathbf{S}} \mathcal{P}_{\mathbf{x}',\vec{w}^+,\vec{w}'^+_{\vec{w}^+},\alpha}\, d\vec{w}^+. \tag{29.41}$$

We use the canonical probability measure of $\mathbf{S}$ in the right hand side.

By Lemma 29.67, the right hand side of (29.41) is independent of the choice of $\vec{w}'^+_{\vec{w}^+}$. We can generalize Lemma 29.48 and use it to show that $\mathcal{P}_{\mathbf{x}',\alpha}$ is injective.

We put

$$E(\mathbf{x}',\alpha) = \mathrm{Im}\mathcal{P}_{\mathbf{x}',\alpha}, \qquad E(\mathbf{x}') = \bigoplus_\alpha E(\mathbf{x}',\alpha).$$

Lemma 29.69. *If* $g \in T^n$, *then*

$$E(g\mathbf{x}', \alpha) = g_* E(\mathbf{x}', \alpha).$$

Proof. For each $\alpha \in \mathfrak{A}(\mathrm{III})$ we decompose g as $g = g_{\alpha,1} + g_{\alpha,2}$ where $g_{\alpha,1} \in S^1_\alpha$, $g_{\alpha,2} \in T^{n-1}_\alpha$.

Let us take $h_i^{\vec{w}^+}$, $h_{i,\alpha}^{\vec{w}^+}$, $\vec{w}'^{+\vec{w}^+}_{(i,j),\alpha}$, $\varphi_{i,j}^{\vec{w}^+}$ as in Lemma 29.65. (They depend on $\vec{w}^+$. So we include it in the notation.) We define $g\vec{w}^+ = (g_{1,\alpha}\vec{w}^+_\alpha)_{\alpha \in \mathfrak{A}}$. (We put $g_{\alpha,1} = 1$ if α is of Type I or II.) More precisely, we lift $g_{\alpha,1}$ to $\widehat{g}_{\alpha,1} = (g_{\alpha,1}, \psi_{\alpha,1}) \in G(\mathbf{x})_0$ (= the connected component of $G(\mathbf{x})$) and define

$$g\vec{w}^+ = (\psi_{\alpha,1}^{-1} \cdot \vec{w}^+_\alpha)_{\alpha \in \mathfrak{A}}.$$

We remark that $g_{\alpha,1} u = u \circ \psi_{\alpha,1}$. We also remark that

$$\psi_{\alpha,1}^{-1} \cdot \vec{w}^+_\alpha = \psi_{\alpha,1}(\vec{w}^+_\alpha).$$

See (29.33.)

Since $\alpha \sim \alpha'$ implies $S^1_\alpha = S^1_{\alpha'}$ and $T^{n-1}_\alpha = T^{n-1}_{\alpha'}$, it follows that $g\vec{w}^+ \in \mathbf{S}$.

We apply Lemma 29.65 to $\mathbf{gx}'$, $g\vec{w}^+$, $\mathbf{x}$ to obtain ${}^{\mathbf{g}}h_i^{g\vec{w}^+}$, ${}^{\mathbf{g}}h_{i,\alpha}^{g\vec{w}^+}$ ${}^{\mathbf{g}}\vec{w}'^{+,g\vec{w}^+}_{(i,j),\alpha}$, ${}^{\mathbf{g}}\varphi_{i,j}^{g\vec{w}^+}$. Then, by definition, we have

$${}^{\mathbf{g}}h_{i,\alpha}^{g\vec{w}^+} = g_{\alpha,2} h_{i,\alpha}^{\vec{w}^+}, \quad {}^{\mathbf{g}}\vec{w}'^{+,g\vec{w}^+}_{(i,j),\alpha} = \vec{w}'^{+\vec{w}^+}_{(i,j),\alpha}, \quad {}^{\mathbf{g}}\varphi_{i,j}^{g\vec{w}^+} = \varphi_{i,j}^{\vec{w}^+}.$$

(See also the proof of Lemma 29.52.)


$w^{+}_{\alpha,a}$
$\psi_{\alpha,1}\left(w^{+}_{\alpha,a}\right)$
$w'^{\vec{w}}_{\alpha,a}$
u
u'
$u'\left(w'^{\vec{w}}_{\alpha,a}\right)$
$u'(\Sigma'(\alpha))$
$h_{i,\alpha}u(\Sigma(\alpha))$
$h_{i,\alpha}N_{w^{+}_{\alpha,a}}$
$u(w^{+}_{\alpha,a})$
$h_{i,\alpha}$
g
$\hat{g}_{\alpha,1} = (g_{\alpha,1}, \psi_{\alpha,1}) \in G(\mathbf{x}_{\alpha})$
$g_{\alpha,1}$
$h_{i,\alpha}g_{\alpha,1}g_{\alpha,2} = h_{i,\alpha}g$
$gh_{i,\alpha}N_{w^{+}_{\alpha,a}}$
$gu'(w'^{\vec{w}}_{\alpha,a})$
$gu'(\Sigma'(\alpha))$
$u\left(\psi_{\alpha,1}\left(w^{+}_{\alpha,a}\right)\right)$
$h_{i,\alpha}g_{\alpha,2}$

Figure 29.4

Therefore

$$\mathcal{P}_{\mathbf{g}\mathbf{x}', g\vec{w}^+, \mathbf{g}\vec{w}'^{+g\vec{w}^+}_{(i,j)}, \alpha} = g_* \mathcal{P}_{\mathbf{x}', \vec{w}^+, \vec{w}'^{+\vec{w}^+}_{(i,j)}, \alpha}.$$

By taking average over $\vec{w}^+$ we obtain the lemma. $\square$

We next prove the $G(\mathbf{x})$ invariance of $E(\mathbf{x}'; \alpha)$. Let $\mathbf{g} = (g, \varphi) \in G(\mathbf{x})$ and $\Psi(\varphi)$ be of the form $[\rho_0, \rho_{0,1}, \rho_{0,2}]$.

We have a map

$$\mathbf{g}_* : C^\infty_0(\Sigma'; u'^* TX \otimes \Lambda^{0,1}) \to C^\infty_0(\Sigma'; (gu')^* TX \otimes \Lambda^{0,1})$$

as in the proof of Lemma 29.4.

Lemma 29.70. *We have*

$$E(\mathbf{g}\mathbf{x}; \rho_0(\alpha)) = \mathbf{g}_* E(\mathbf{x}; \alpha).$$

Proof. Let $\mathbf{g} = (g, \varphi)$ and $\vec{w}^+ \in \mathbf{S}$. By Lemma 29.57, we have $\varphi(\vec{w}^+) \in \mathbf{S}$. The rest of the proof is the same as the proof of Lemma 29.49. $\square$

We use it to define a Kuranishi chart of a neighborhood of the $T^n \times \mathfrak{S}_\ell$ orbit of $\mathbf{x}$ in the same way as before. It is invariant under the $G(\mathbf{x})$ action and of the permutation of the interior marked points.

We now prove

Lemma 29.71. *The Kuranishi chart we defined above is disk-component-wise.*

Proof. As we mentioned in Remark 29.68, $h(\alpha)$ depends only on $\vec{w}^+_\alpha$ and is independent of $\vec{w}^+_{\alpha'}$, $\alpha \ne \alpha'$. So we only need to study $i_{\mathbf{v}'}$ to prove the lemma.

Here we describe Condition 29.58 (5) more precisely. During this description we write $i_{\mathbf{v}';\mathbf{v}}$ in place of $i_{\mathbf{v}'}$. Namely we specify $\mathbf{v}$.

Let $\mathbf{v} = (\Sigma, \vec{z}^+, \vec{w}^+)$. We decompose

$$\Sigma = \bigcup_{\alpha \in \mathfrak{A}} \Sigma(\alpha),$$

where $\Sigma(\alpha)$ is a disk component (together with sphere components attached on it).

Definition 29.72. Let $\mathrm{Part}(\mathfrak{A})$ be the set of all $\mathbb{B} = \{\mathfrak{A}(\beta) \mid \beta \in \mathfrak{B}\}$ such that:

(1) $\mathfrak{A}(\beta) \subset \mathfrak{A}$.
(2) $\mathfrak{A} = \bigcup_{\beta \in \mathfrak{B}} \mathfrak{A}(\beta)$.
(3) If $\beta \ne \beta'$, then $\mathfrak{A}(\beta) \cap \mathfrak{A}(\beta') = \emptyset$.
(4) $\bigcup_{\alpha \in \mathfrak{A}(\beta)} \Sigma(\alpha)$ is connected for each $\beta \in \mathfrak{B}$.

Let $\mathbb{B} = \{\mathfrak{A}(\beta) \mid \beta \in \mathfrak{B}\}, \mathbb{B}' = \{\mathfrak{A}'(\beta') \mid \beta' \in \mathfrak{B}'\} \in \mathrm{Part}(\mathfrak{A})$. We say $\mathbb{B} < \mathbb{B}'$ if for each $\beta \in \mathfrak{B}$ there exists $\beta' \in \mathfrak{B}'$ such that $\mathfrak{A}(\beta) \subset \mathfrak{A}'(\beta')$.

For $\mathbb{B} = \{\mathfrak{A}(\beta) \mid \beta \in \mathfrak{B}\} \in \mathrm{Part}(\mathfrak{A})$ let $\mathfrak{U}(\mathbf{v}; \mathbb{B})$ be the set of elements $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+) \in \mathfrak{U}(\mathbf{v})$ such that the following holds:

If $\alpha, \alpha' \in \mathfrak{A}$ and $i_{\mathbf{v}';\mathbf{v}}(\Sigma(\alpha) \setminus \mathcal{U}(S(\Sigma))$ belongs to the same disk component as $i_{\mathbf{v}';\mathbf{v}}(\Sigma(\alpha') \setminus \mathcal{U}(S(\Sigma))$, then there exists $\beta \in \mathfrak{B}$ such that $\alpha, \alpha' \in \mathfrak{A}(\beta)$.

By definition, $\mathfrak{U}(\mathbf{v}; \mathbb{B})$ is a closed set. Moreover if $\mathbb{B} < \mathbb{B}'$, then $\mathfrak{U}(\mathbf{v}; \mathbb{B}) \subset \mathfrak{U}(\mathbf{v}; \mathbb{B}')$.

Let $\mathbf{v}' = (\Sigma', \vec{z}'^+, \vec{w}'^+) \in \mathfrak{U}(\mathbf{v})$ and $\beta \in \mathfrak{B}$. We put

$$\Sigma'(\beta) = \bigcup_{\alpha \in \mathfrak{A}(\beta)} \text{The component containing } i_{\mathbf{v}';\mathbf{v}}(\Sigma(\alpha) \setminus \mathcal{U}(S(\Sigma)).$$

By definition

$$\Sigma'(\beta) \cap \Sigma'(\beta') \subset S(\Sigma),$$

if $\beta \ne \beta'$. Namely

$$\Sigma' = \bigcup_{\beta \in \mathfrak{B}} \Sigma'(\beta) \tag{29.42}$$

is a 'disjoint' union. (More precisely, two different members intersect only at one boundary singular point, if any.)

We put

$$\overset{\circ}{\mathfrak{U}}(\mathbf{v};\mathbb{B}) = \mathfrak{U}(\mathbf{v};\mathbb{B}) \setminus \bigcup_{\mathbb{B}'<\mathbb{B}} \mathfrak{U}(\mathbf{v};\mathbb{B}').$$

Note that (29.42) is a decomposition to the disk components if and only if $\mathbf{v}'$ is an element of $\overset{\circ}{\mathfrak{U}}(\mathbf{v};\mathbb{B})$.

Let $\mathbf{v}' \in \mathfrak{U}(\mathbf{v};\mathbb{B})$. The component $\Sigma'(\beta)$ together with interior marked points on it defines an element $\mathbf{v}'(\beta)$. We have a map

$$i_{\mathbf{v}'(\beta);\mathbf{v}(\beta)} : \bigcup_{\alpha \in \mathfrak{A}_\beta} (\Sigma(\alpha) \setminus \mathcal{U}(S(\Sigma)) \to \Sigma'(\beta).$$

Suppose $\mathbf{v}' \in \overset{\circ}{\mathfrak{U}}(\mathbf{v};\mathbb{B})$ and $\mathbf{v}'' \in \mathfrak{U}(\mathbf{v}')$. Then we have a map

$$i_{\mathbf{v}'';\mathbf{v}'} : \bigcup_{\beta \in \mathfrak{B}} (\Sigma'(\beta) \setminus \mathcal{U}(S(\Sigma')) \to \Sigma''.$$

Now the following is the precise form of Condition 29.58 (5).

Condition 29.73. If $\mathbf{v}' \in \overset{\circ}{\mathfrak{U}}(\mathbf{v};\mathbb{B})$ and $\mathbf{v}'' \in \mathfrak{U}(\mathbf{v}') \cap \mathfrak{U}(\mathbf{v})$, then we have

$$i_{\mathbf{v}'';\mathbf{v}} = i_{\mathbf{v}'';\mathbf{v}'} \circ i_{\mathbf{v}'(\beta);\mathbf{v}(\beta)} \tag{29.43}$$

on $\bigcup_{\alpha \in \mathfrak{A}(\beta)} \Sigma(\alpha) \setminus \mathcal{U}(S(\Sigma))$.

Sublemma 29.74. *There exists $i_{\mathbf{v}';\mathbf{v}}$ which satisfies Condition 29.58 (1) - (4), (6) and Condition 29.73.*

The proof is by induction on the number of disk components.

Now using (29.43) and Remark 29.68 we can prove Lemma 29.71. □

The proof of Proposition 29.5 is now complete. □

Once Proposition 29.5 is established, we can prove Proposition 29.1 in the same way as [Fu2] Corollary 3.1. □

30. APPENDIX 4: T^n EQUIVARIANT CONTINUOUS FAMILY OF MULTISECTIONS

In this section we construct a continuous family of multisections with cyclic symmetry for the Kuranishi structure produced in the last section. Most of the construction is a straightforward combination of those employed in [FOOO4, FOOO5, Fu2]. We only need to prove the following Lemma 30.1.

We consider (U, E, Γ, s) and f, N with the following properties:

1. U is a smooth manifold with a T^n action.
2. E is a T^n equivariant vector bundle on U.
3. Γ is a finite group which has *right* action on U and E. Moreover E is a Γ equivariant vector bundle. The action of Γ on U is effective.

(4) The Γ action and T^n action commute and so defines a $T^n \times \Gamma$ action on E.
(5) N is a smooth manifold and $f : U \to N$ is a smooth submersion.
(6) s is a $T^n \times \Gamma$ equivariant section of E.

Lemma 30.1. *There exist a manifold W with a T^n action, and a multisection $\mathfrak{s}$ of $E/\Gamma \times E \to U_0/\Gamma \times W$ that satisfy the following properties:*

(1) *U_0 is a $T^n \times \Gamma$ invariant neighborhood of $s^{-1}(0)$.*
(2) *$\mathfrak{s}$ is smooth and is transversal to 0.*
(3) *The restriction of f to $\mathfrak{s}^{-1}(0)$ is a submersion to N.*
(4) *Each branch of $\mathfrak{s}$ can be chosen to be as close to s as we want in C^0 sense.*

Proof. Let $p \in U$. We put $I_p = \{g \in T^n \mid gp = p\}$, $\Gamma_p = \{\gamma \in \Gamma \mid p = p\gamma\}$, $G_{p,+} = \{(g,\gamma) \in T^n \times \Gamma \mid gp = p\gamma\}$, which contains $I_p \times \Gamma_p$.

We define a *left* action of Γ by $\gamma p = p\gamma^{-1}$. Hereafter we only use this left action. The group $G_{p,+}$ acts on X and E as a subgroup of $T^n \times \Gamma$. By this action, p is a fixed point of $G_{p,+}$. So $G_{p,+}$ acts on the fiber E_p.

We choose a submanifold X_p containing p which is a local transversal to the T^n orbit of p. We may choose X_p so that it is $G_{p,+}$ invariant. We take a $T^n \times \Gamma$ invariant metric of U and a T^n and Γ invariant connection of E.

We put $W'_p = E_p$ and define a section s_p of E on $X_p \times W'_p$ by

$$s_p(x,w) = s(x) + \Pi_p^x(w). \tag{30.1}$$

Here Π_p^x is the parallel transport along the minimal geodesic joining p and x. s_p is transversal to 0. Moreover (3) in the conclusion of Lemma 30.1 is satisfied. In fact the restriction of the projection $X_p \times W'_p \to X_p$ to $s_p^{-1}(0)$ is a submersion.

We take a closed subgroup $J_p \subset T^n$ such that the map $J_p \times I_p \to T^n$ defined by $(g,h) \mapsto gh$ is finite to one and surjective. We put $T_p = J_p \times G_{p,+}$. We define $\pi : T_p \to T^n$ by

$$\pi(g,(h,\gamma)) = gh.$$

π is a surjective homomorphism. We put $H_p = \mathrm{Ker}\pi \subset T_p$. H_p is a finite group. The restriction of the projection $J_p \times G_{p,+} \to G_{p,+}$ to H_p defines a homomorphism $H_p \to G_{p,+}$, that is injective.

We recall $I_p \subset G_{p,+}$ by the embedding $g \mapsto (g,e)$ where $e \in \Gamma$ is the unit. Therefore we can embed

$$J_p \cap I_p \to H_p$$

by $g \mapsto (g^{-1},(g,e))$.

We let J_p act trivially on W'_p. We then define the a diagonal action on $T_p = J_p \times G_{p,+}$. Its restriction to H_p defines an action of H_p on W'_p. The embeddings $H_p \to G_{p,+}$ and $I_p \cap J_p \to H_p$ preserves this action to W'_p. Here we regard $I_p \cap J_p \subset I_p \subset T^n$ to define $I_p \cap J_p$ action.

Let $U_p = T^n X_p$ be the T^n orbit of X_p, which is a neighborhood of $T^n p$, the T^n orbit of p. We remark that if $y \in U_p$ then there exist $g \in J_p$ and $x \in X_p$ such that $y = gx$. Moreover if $g'x' = gx$ as above, then $g^{-1}g' \in J_p \cap I_p$.

We have

$$s_p(\mathbf{g}x, \mathbf{g}w) = \mathbf{g}_* s_p(x,w)$$

where $\mathbf{g}_*$ in the right hand side is the map $\mathbf{g}_* : E_p \to E_{\mathbf{g}p}$ induced by the $G_{p,+}$ action on E.

We put

$$H_p = \{\mathbf{g}_i = (g_i,(h_i,\gamma_i)) \mid i = 1,\dots,\ell\}.$$

Now we define an ℓ-multisection s'_p on $U_p \times W'_p$ by

$$s'_p(y,w) = \{g s_p(x, \mathbf{g}_i w) \mid i = 1,\dots,\ell\},$$

where $y = gx$ ($g \in J_p$, $x \in X_p$). This is well-defined. In fact, if $p = g'x' = gx$ then $g' = gg_a$, $x = g_a x'$ for some $g_a \in J_p \cap I_p$. Therefore

$$g' s_p(x', \mathbf{g}_i w) = g s_p(x, \mathbf{g}_i g_a w).$$

This implies that s'_p is well-defined. (Note $I_p \cap J_p \subset H_p$.)

The multisection s'_p is T_p equivariant. In fact J_p equivariace is immediate from definition. If $\mathbf{g} = (h_1, (h_2, \gamma)) \in G_{p,+}$ then for $y = gx$ ($g \in J_p$, $x \in X_p$) we have

$$\begin{aligned} s'_p(\mathbf{g}y, \mathbf{g}w) &= \{g s_p(\mathbf{g}x, \mathbf{g}_i \mathbf{g} w) \mid i = 1,\dots,\ell\} \\ &= \{g\mathbf{g} s_p(x, \mathbf{g}_i w) \mid i = 1,\dots,\ell\} = \mathbf{g} s'_p(y,w). \end{aligned}$$

(Note H_p is a normal subgroup of T_p.)

It is easy to see that

$$s'_p(x, \mathbf{g}w) = s'_p(x,w)$$

for $\mathbf{g} \in H_p$.

We remark that the quotient space W'_p/H_p is an orbifold. (Note however that it may not be effective. Namely there may exist a nontrivial element of H_p whose action on W'_p is trivial.)

We take an open subset W_p of W'_p/H_p which is a manifold and is sufficiently close to the origin. The quotient group $T^n = G_{p,+}/H_p$ acts on W_p.

Then s'_p induces a W_p parametrized multisection s''_p on U_p.

We put

$$\Gamma_{p,+} = \{\gamma \in \Gamma \mid \gamma U_p \cap U_p \neq \emptyset\}.$$

We may choose X_p so small that $\gamma \in \Gamma_{p,+}$ holds only when $\gamma p \in T^n p$. Then we have a surjective homomorphism

$$G_{p,+} \to \Gamma_{p,+}, \qquad (g,\gamma) \mapsto \gamma.$$

For $(g_i, \gamma_i) \in G_{p,+}$, we write

$$g_i = h_i k_i, \qquad h_i \in J_p,\ k_i \in I_p. \tag{30.2}$$

For $x \in U_p$ and $w \in W_p$ we define ℓ^2-multisection $s^{(3)}_p$ by

$$s^{(3)}_p(x,w) = \{s''_p(x, k_i w) \mid i = 1,\dots,\ell\}. \tag{30.3}$$

(Note $\ell = \#G_{p,+}$.) We remark that $s''_p(x, fw) = s''_p(x,w)$ if $f \in J_p \cap I_p$. Therefore (30.3) is independent of the choice of the decomposition (30.2).

We now extend $s^{(3)}_p$ to $U^+_p = \Gamma U_p$ as follows. Let $(y,w) \in U^+_p \times W_p$. We have $y = \gamma x$ with $\gamma \in \Gamma$, $x \in U_p$. We put

$$s^{(4)}_p(y,w) = \gamma_* s^{(3)}_p(x,w). \tag{30.4}$$

We claim that (30.4) is well-defined. In fact let $y = \gamma x = \gamma' x'$, with $\gamma' \in \Gamma$, $x' \in U_p$. We have $\mu = \gamma^{-1}\gamma' \in \Gamma_{p,+}$, $x = \mu x'$. We take $g \in T^n$ such that $\mathbf{g} = (g,\mu) \in G_{p,+}$. We put $g = hk$ with $h \in J_p$, $k \in I_p$. Then

$$\begin{aligned} &\gamma' s^{(3)}_p(x',w) = \gamma\mu s^{(3)}_p(x',w) = \gamma g^{-1}\mathbf{g} s^{(3)}_p(x',w) = \gamma k^{-1}h^{-1} s^{(3)}_p(\mathbf{g}x', \mathbf{g}w) \\ &= \gamma k^{-1}h^{-1} s^{(3)}_p(\mathbf{g}x', w) = \gamma k^{-1} s^{(3)}_p(h^{-1}\mathbf{g}x', w) = \gamma s^{(3)}_p(k^{-1}h^{-1}\mathbf{g}x', k^{-1}w) \\ &= \gamma s^{(3)}_p(\mu x', k^{-1}w) = \gamma s^{(3)}_p(x, k^{-1}w) = \gamma s^{(3)}_p(x,w). \end{aligned}$$

It is easy to see that $s_p^{(4)}$ is $\Gamma \times T^n$ invariant. (30.1) implies that s_p satisfies (2) and (3) of Lemma 30.1. Therefore $s_p^{(4)}$ satisfies (2) and (3) of Lemma 30.1 also. Thus $s_p^{(4)}$ gives a required W_p parametrized family of multisections on U_p^+.

Then by gluing $s_{p_i}^{(4)}$ for various p_i by an appropriate partitions of unity, we obtain the required family of multisections. □

Using Lemma 30.1, the rest of the construction is the same as [FOOO4], [FOOO5] and [Fu2] and so is omitted.

31. Appendix 5: Hochschild and quantum cohomologies

In this paper we study the operators $\mathfrak{p}$, $\mathfrak{q}$ and the cyclically symmetric version $\mathfrak{q}^{\mathfrak{c}}$ of $\mathfrak{q}$, in the case of toric manifold and its Lagrangian submanifold that is an orbit of T^n action. We use it to prove our main result, Theorem 1.1. We use various properties of these operators for this purpose. We have proved them only in our toric case.

Actually those operators are defined in greater generality and most of the properties we use here indeed hold in greater generality. There are various literatures which conjecture/claim/prove/use them in various level of generality and of rigor.

The purpose of this section is to briefly summarize the properties of operators $\mathfrak{p}$, $\mathfrak{q}$ etc. which are expected to hold, in order to put the discussion of this paper in proper perspective. We also mention various earlier results. We might overlook some other existing literature because our knowledge are certainly incomplete.

We remark that the operators $\mathfrak{p}$, $\mathfrak{q}$ etc. are defined in [FOOO2] Chapter 3 and [FOOO3] Section 3.8 for any pair (X, L) where X is a symplectic manifold and L is its relatively spin closed Lagrangian submanifold. We first review its relation to Hochschild cohomology here.

Let C be a filtered $\Lambda_{0,\rm nov}$ module. We assume it is free. We consider the Hochschild complex:

$$CH(C[1], C[1]) = \prod_{k=0}^{\infty} Hom_{\Lambda_{0,\rm nov}}(B_k(C[1]), C[1]). \tag{31.1}$$

More precisely, for $\varphi_0 \in Hom_{\Lambda_{0,\rm nov}}(B_0(C[1]), C[1]) = C[1]$, we assume $\varphi_0(1) \equiv 0 \mod \Lambda_{0,\rm nov}^+$, in addition. Its boundary operator is defined by

$$\begin{aligned}(\delta\varphi)(x_1, \dots, x_k) = & \sum_{k_1+k_2=k+1} \sum_{j=1}^{k_1} (-1)^{*_1} \mathfrak{m}_{k_1}(x_1, \dots, \varphi_{k_2}(x_j, \dots), \dots, x_k) \\ & + \sum_{k_1+k_2=k+1} \sum_{j=1}^{k_1} (-1)^{*_2} \varphi_{k_1}(x_1, \dots, \mathfrak{m}_{k_2}(x_j, \dots), \dots, x_k).\end{aligned} \tag{31.2}$$

Here $\varphi = (\varphi_k)_{k=0}^{\infty}$ and

$$*_1 = \deg \varphi_{k_2}(\deg' x_1 + \ldots + \deg' x_{j-1}), \quad *_2 = \deg' x_1 + \ldots + \deg' x_{j-1}.$$

(See [FOOO3] Section 4.4.5.) Let us consider the case when $C = H(L; \Lambda_{0,nov})$ as in [FOOO3] Theorem A. The operators $\mathfrak{q}_{\ell;k}$ defines a map

$$\widehat{\mathfrak{q}} : E(H(X; \Lambda_{0,\rm nov})[2]) \to CH(H(L; \Lambda_{0,\rm nov})[1], H(L; \Lambda_{0,\rm nov})[1])$$

by

$$\widehat{\mathfrak{q}}(\mathbf{p}_1,\dots,\mathbf{p}_\ell)(x_1,\dots,x_k) = \mathfrak{q}_{\ell;k}(\mathbf{p}_1,\dots,\mathbf{p}_\ell;x_1,\dots,x_k). \tag{31.3}$$

It is easy to see that the image of this map consists of cycles. See [FOOO3] Section 7.4.

We next define product structure on $CH(C[1],C[1])$. Let $\varphi = (\varphi_k)_{k=0}^\infty$, $\psi = (\psi_k)_{k=0}^\infty$. We put

$$\begin{aligned}&\mathfrak{m}_2(\varphi,\psi)(x_1,\dots,x_k)\\ &= \sum_{k_1+k_2=k}(-1)^*\mathfrak{m}_2(\varphi_{k_1}(x_1,\dots,x_{k_1}),\psi_{k_2}(x_{k_1+1},\dots,x_k))\end{aligned} \tag{31.4}$$

where

$$* = \deg\psi_{k_2}(\deg' x_1 + \cdots + \deg' x_{k_1}).$$

Then

$$\varphi\cup\psi = (-1)^{\deg\varphi(\deg\psi+1)}\mathfrak{m}_2(\varphi,\psi)$$

defines an associative product.

Remark 31.1. (1) We can define $\mathfrak{m}_k(\varphi_1,\dots,\varphi_k)$ in a similar way. It defines a structure of A_∞ algebra on $CH(C[1],C[1])$.

(2) This A_∞ structure is a particular case of one defined in [Fu1] Theorem-Definition 7.55. Namely we consider the filtered A_∞ category with one object whose endomorphism algebra is the A_∞ algebra C. Then $CH(C[1],C[1])$ is identified with the set of all prenatural transformations from identity functor to identity functor. In [Fu1] Theorem-Definition 7.55, the filtered A_∞ structure of it is constructed. If we apply it to our particular case, it coincides with the above A_∞ structure on $CH(C[1],C[1])$.

Claim 31.2. *The map* $\widehat{\mathfrak{q}}$ *defines a ring homomorphism*

$$QH(X;\Lambda_{0,nov}) \to HH(H(L;\Lambda_{0,nov})[1],H(L;\Lambda_{0,nov})[1]). \tag{31.5}$$

Here we put quantum cohomology ring structure on the left hand side. The right hand side is δ*-cohomology of* $HH(H(L;\Lambda_{0,nov})[1],H(L;\Lambda_{0,nov})[1])$*, that is the Hochschild cohomology.*

Remark 31.3. (1) When we consider several Lagrangian submanifolds instead of one Lagrangian submanifold, it forms a filtered A_∞ category called Fukaya category. Then we can modify the statement so that the right hand side is a Hochschild cohomology of Fukaya category.

(2) We can include bulk deformation by an element $\mathfrak{b} \in H^{ev}(L;\Lambda_{0,\mathrm{nov}}^+)$ then Claim 31.2 still holds. Namely we put $QH(X;\Lambda_{0,\mathrm{nov}})$ the $\mathfrak{b}$-deformed quantum cup product. (See (2.33).) We use also use $\mathfrak{b}$-deformed A_∞ structure on Floer cohomology of the right hand side of (31.2). Then (31.2) becomes a ring homomorphism in this generality.

Conjecture 31.4. *We use quantum (higher) Massey product on* $QH(X;\Lambda_{0,\mathrm{nov}})$ *to obtain a filtered* A_∞ *structure on it. (See* [FOOO6] *Corollary 1.10.) Then* (*31.5*) *is extended to a filtered* A_∞ *homomorphism.*

Remark 31.3 (1)(2) apply to Conjecture 31.4.

Theorem 9.1 is closely related to Claim 31.2, but it looks slightly different. We will explain its relation now. (For the simplicity of notation we consider the case when $\mathfrak{b} = 0$, namely the case without bulk deformation. Including bulk deformations is fairly straightforward.)

We first consider the case when the potential function $\mathfrak{PO}$ is Morse. In that case there are finitely many $(L(u_i), b_i)$ $(i = 1, \ldots, \mathrm{rank} H(X))$, such that

$$HF((L(u_i), b_i), (L(u_i), b_i)) \cong H(T^n).$$

(See Lemma 24.5.) Moreover $HF((L(u_i), b_i), (L(u_i), b_i))$ is a (nondegenerate) Clifford algebra as a ring. So we have

$$HH(H(L(u_i)), H(L(u_i)); \Lambda) \cong \Lambda.$$

(Here we use the A_∞ structure $\mathfrak{m}_k^{b_i}$ to define Hochschild cohomology in the right hand side.) Thus Claim 31.2 implies that there exists a left homomorphism

$$QH(X; \Lambda) \to \prod_i \Lambda. \tag{31.6}$$

Using the identification (2.16) it is easy to identify $\prod_i \Lambda$ with $\mathrm{Jac}(\mathfrak{PO}) \otimes_{\Lambda_0} \Lambda$. Then the ring homomorphism (31.5) induces a ring homomorphism to $\mathrm{Jac}(\mathfrak{PO}) \otimes_{\Lambda_0} \Lambda$. It coincides with the Kodaira-Spencer map appearing in Theorem 9.1.

We next consider the case when $\mathfrak{PO}$ is not necessarily Morse. We consider the basis $\mathbf{e}$, $\mathbf{e}_i$ $(i = 1, \ldots, n)$, $\mathbf{e}_I$ $(I \subseteq \{1, \ldots, n\})$ of $H^*(T^n; \mathbb{Q})$. Here $\deg \mathbf{e} = 0$, $\deg \mathbf{e}_i = 1$, $\deg \mathbf{e}_I \geq 2$.

Let $\varphi = (\varphi_k)_{k=0}^{\infty}$ be an element of $CH(H(L; \Lambda_0)[1], H(L; \Lambda_0)[1]) \otimes_{\Lambda_0} \Lambda$. We put

$$f_{j_1, \ldots, j_k} = \varphi_k(\mathbf{e}_{j_1}, \ldots, \mathbf{e}_{j_k}) \in \Lambda.$$

We define

$$\mathfrak{k}(\varphi) = \sum_{k=0}^{\infty} \sum_{j_1, \ldots, j_k = 1}^{n} \frac{1}{k!} f_{j_1, \ldots, j_k} x_{j_1} \cdots x_{j_k} \in \Lambda_0[[x_1, \ldots, x_n]] \otimes_{\Lambda_0} \Lambda. \tag{31.7}$$

Lemma 31.5. *(31.7) induces a ring homomorphism*

$$HH(H(L; \Lambda_0)[1], H(L; \Lambda_0)[1]) \otimes_{\Lambda_0} \Lambda \to \frac{\Lambda_0[[x_1, \ldots, x_n]] \otimes_{\Lambda_0} \Lambda}{\left(\frac{\partial \mathfrak{PO}^{u_j}(x+b)}{\partial x_i}; i = 1, \ldots, n\right)}.$$

Here we use $\mathfrak{m}_k^b$ to define Hochschild boundary operator in the left hand side.

We will write the boundary operator δ in (31.2) as $\delta_{\mathfrak{b}}$ in case we use $\mathfrak{m}_*^{\mathfrak{b}}$.

Proof. It suffices to show that the image of the Hochschild boundary operator maps to the Jacobian ideal in the right hand side by the map (31.7). Let $\psi = (\psi_k)_{k=0}^{\infty}$ be the Hochschild chain. We define $h^i_{j_1, \ldots, j_k}$ for $i, j_1, \ldots, j_k = 1, \ldots, n$ by

$$\psi_k(\mathbf{e}_{j_1}, \ldots, \mathbf{e}_{j_k}) - \sum_{i=1}^{n} h^i_{j_1, \ldots, j_k} \mathbf{e}_i \in \Lambda[\mathbf{e}, \mathbf{e}_I; \deg \mathbf{e}_I \geq 2].$$

We then define

$$h_i = \sum_{k=0}^{\infty} \sum_{j_1, \ldots, j_k = 1}^{n} \frac{1}{k!} h^i_{j_1, \ldots, j_k} x_{j_1} \cdots x_{j_k} \in \Lambda_0[[x_1, \ldots, x_n]] \otimes_{\Lambda_0} \Lambda.$$

It is straightforward to check

$$\mathfrak{k}(\delta_{\mathfrak{b}}(\psi)) = \sum_{i=1}^{n} h_i \frac{\partial \mathfrak{PO}(x_1 + b_1, \ldots, x_n + b_n)}{\partial x_i}.$$

Hence the lemma. □

We can prove

$$\mathrm{Jac}(\mathfrak{PO}) \otimes_{\Lambda_0} \Lambda \cong \prod_j \frac{\Lambda_0[[x_1,\dots,x_n]] \otimes_{\Lambda_0} \Lambda}{\left(\frac{\partial \mathfrak{PO}^{u_j}(x+b)}{\partial x_i}; i = 1,\dots,n\right)}.$$

We remark that the left hand side is a quotient of the strictly convergent power series ring, while the right hand side is a quotient of the formal power series ring. We can however prove that they are isomorphic in the present situation in a way similar to Lemma 12.8.

Therefore, by Lemma 31.5, we obtain a ring homomorphism

$$HH(H(L;\Lambda_0)[1], H(L;\Lambda_0)[1]) \otimes_{\Lambda_0} \Lambda \to \mathrm{Jac}(\mathfrak{PO}) \otimes_{\Lambda_0} \Lambda. \tag{31.8}$$

Composition of (31.8) and (31.5) is the Kodaira-Spencer map appearing in Theorem 9.1.

We can prove Claim 31.2 in a way similar to Theorem 9.1. We do not try to prove this here since we do not use it in this paper.

The homomorphism $\widehat{q}$ in (31.5) is related to the homomorphism $i^{\#}_{\mathrm{qm},(b,u)}$ given in (17.2) as follows. Let $\varphi = (\varphi_k)_{k=0}^{\infty}$ be a Hochschild chain. We put

$$\overline{\varphi} = \varphi_0(1) \in C.$$

It is easy to see that $\overline{\varphi}$ is a $\mathfrak{m}_1$-cycle if φ is a Hochschild cycle. Thus we have

$$HH(H(L;\Lambda_0)[1], H(L;\Lambda_0)[1]) \otimes_{\Lambda_0} \Lambda \to HF((L(u),b),(L(u),b);\Lambda). \tag{31.9}$$

$i^{\#}_{\mathrm{qm},(b,u)}$ is a composition of (31.9) and $\widehat{q}$. Since (31.9) and $\widehat{q}$ are both ring homomorphisms

$$i^{\#}_{\mathrm{qm},(b,u)} : QH(X;\Lambda) \to HF((L(u),b),(L(u),b);\Lambda) \tag{31.10}$$

is also a ring homomorphism.

Remark 31.6. (1) We remark that $i^{\#}_{\mathrm{qm},(b,u)}$ is constructed in [FOOO2] and Theorem 3.8.62 [FOOO3]. However the fact that it is a *ring* homomorphism is neither stated nor proved there.

(2) For the case of monotone L, the fact that $i^{\#}_{\mathrm{qm},(b,u)}$ is a ring homomorphism is closely related to a statement proved in Theorem A [BC2].

Here are some more precise statements on their relationship.

(a) The algebra which is denoted by $QH_*(L;\mathcal{R})$ in [BC2] is isomorphic to the Floer homology. Namely

$$QH_*(L;\mathcal{R}) \otimes_{\mathcal{R}} \Lambda_{0,nov} \cong HF((L,0),(L,0);\Lambda_{0,nov}),$$

as rings. (Here it is assumed that L is monotone.) This fact had been proved in the year 2000 version [FOOO1] pages 287–291, *without the monotonicity assumption*, except the compatibility with ring structure. Note we may take $\mathcal{R} = \mathbb{Q}[T,T^{-1}]$ as the coefficient ring in the monotone case as explained in [FOOO3] Chapter 2.

(b) The map

$$QH(X;\mathcal{R}) \otimes QH(L;\mathcal{R}) \to QH(L;\mathcal{R}) \tag{31.11}$$

appears in [BC2] Theorem A (ii). Via the isomorphism mentioned above, this map coincides with the map

$$(\mathbf{p}, x) \mapsto \mathfrak{q}_{1;1}(\mathbf{p}, x). \tag{31.12}$$

It is proved in [BC2] that (31.11) determines a module structure on $QH(L;\mathcal{R})$ over the quantum cohomology ring $QH(X;\mathcal{R})$. In view of the fact that $\mathfrak{m}_0 = 0$ in monotone case, this statement is a special case of the Claim 31.2. (See also [Se2].)
We remark that (31.12) is slightly different from (31.10). The latter map corresponds to $\mathbf{p} \mapsto \mathfrak{q}_{1;0}(\mathbf{p})$.

(3) The homomorphism (31.5) had been known much earlier. It seems that the first reference where it is mentioned is [Ko2]. (In [Ko2], the right hand is replaced by Hochschild homology of Fukaya category which is a more general statement than the present context. There may also be several physics literature containing statements related to this homomorphism.) The category version of the map (31.5) is believed to be an isomorphism under suitable assumptions. This is stated explicitly as a conjecture by P. Seidel in Conjecture 1 [Se5] for the exact case. (In noncompact case such as the exact case, one needs to replace quantum cohomology by symplectic homology.) There are also earlier references such as [Se4] but [Se5] contains the statement in the most transparent way. Seidel also constructed the map (31.5) and proved that it is a ring homomorphism in the exact case in Section 6 (6B) [Se4]. ('The technicalities' mentioned in the 4-th line of Section 6 (6B) [Se4] can be easily taken care of by now, that is 10 years after [Se4] was written.)

Recently this ring homomorphism (31.5) is familiar to many experts and becomes a kind of 'folklore' in the field. However as far as the authors know, rigorous proof of the ring homomorphism property is written in detail only for the cases where L is either exact (Seidel), monotone (Biran-Cornea), or toric (in this paper),

We also remark that this kinds of application of 'open-closed Gromov-Witten invariant' in Floer homology first appeared in the literature in [Fl2].

There is a cyclic analogue of the map (31.5). Namely we can use the cyclically symmetric version $\mathfrak{q}^{\mathfrak{c}}$ of $\mathfrak{q}$ to define a map

$$\widehat{\mathfrak{q}}^{\mathfrak{c}} : QH(X;\Lambda_{0,nov}) \to HC(H(L;\Lambda_{0,nov})[1];\Lambda) \tag{31.13}$$

by

$$\widehat{\mathfrak{q}}^{\mathfrak{c}}(\mathbf{p}_1,\dots,\mathbf{p}_\ell)(x_1,\dots,x_k,x_0) = \left\langle \mathfrak{q}^{\mathfrak{c}}_{\ell;k}(\mathbf{p}_1,\dots,\mathbf{p}_\ell;x_1,\dots,x_k),x_0\right\rangle.$$

Here the right hand side of (31.13) is the cyclic cohomology. This homomorphism is expected to play an important role to pursue the line of ideas explained in Section 24.

The operator $\mathfrak{p}$ is constructed in [FOOO3] Section 3.8 for any pair (X,L). It induces a map

$$\widehat{\mathfrak{p}} : HC(H(L;\Lambda_{0,nov})) \to H(X;\Lambda_0) \tag{31.14}$$

by

$$[x_1 \otimes \ldots \otimes x_k] \mapsto \mathfrak{p}_k(x_1,\dots,x_k).$$

Here the left hand side of (31.14) is cyclic homology.

Remark 31.7. We can use $\mathfrak{p}_{\ell;k}$ of [FOOO3] Section 3.8.9 to extend (31.14) to a map

$$EH(X;\Lambda_{0,nov}) \otimes HC(H(L;\Lambda_{0,nov})) \to H(X;\Lambda_0)$$

in a similar way.

If b is a bounding cochain (namely if $\mathfrak{m}_0^b = 0$), then we have

$$HF((L,b);\Lambda_0) \to HC(H(L;\Lambda_{0,nov}))$$

by

$$[x] \mapsto [x] \in B_1^{\rm cyc} H(L;\Lambda_{0,nov})[1].$$

(We use A_∞ structure $\mathfrak{m}^b$ to define the above cyclic homology.) Combined with (31.14), we obtain

$$i_{L,b} : HF((L,b);\Lambda_0) \to H(X;\Lambda_0). \tag{31.15}$$

It coincides with the map $i_{\#,{\rm qm},(b,u)}$ in (17.1).

Theorem 20.1 is expected to hold in that generality. Namely we expect

$$\begin{aligned} &\langle i_{L,b}(\mathfrak{v}), i_{L,b}(\mathfrak{w})\rangle_{{\rm PD}_X} \\ &= \sum_{I,J} \pm g^{IJ} \langle \mathfrak{m}_2^{\mathfrak{c},b}(\mathbf{e}_I,\mathfrak{v}), \mathfrak{m}_2^{\mathfrak{c},b}(\mathbf{e}_J,\mathfrak{w})\rangle_{{\rm PD}_L}. \end{aligned} \tag{31.16}$$

Here $\mathbf{e}_I$ is a basis of the cohomology $H(X;\Lambda)$. See Proposition 26.17 for sign ±. Other notations are the same as in Section 20.

Many parts of the proof in Section 20 can be generalized to the non-toric case. However we take short cut in several places in Section 20. So several parts of the proof should be rewritten in the general case.

Remark 31.8.

(1) In the monotone case the map $i_{L,0}$ coincides with the map i_L appearing in [BC2] Theorem A (iii). [BC2] attributes those maps to the paper [Al]. Indeed, the map ξ_k and τ_k which appears in Theorem 1.5 [Al] (defined in the monotone case) coincide with (31.10) and (31.15), respectively.

    One of the main applications of Theorem 1.5 [Al] is to prove nontriviality of Lagrangian Floer cohomology under certain conditions. The fundamental chain of L is used for this purpose there. In the proof of [FOOO1] Theorem 6.13 (6.14.3) a similar argument had been used for the same purpose. The operators $\mathfrak{p}$ was introduced in [FOOO2] in order to make this argument more systematic.

(2) We used the moduli space of pseudo-holomorphic annuli to prove (31.16) in the toric case. The moduli space of pseudo-holomorphic annuli is also used in [BC1] and in [Ab].

References

[Ab] M. Abouzaid, *A geometric criterion for generating the Fukaya category*, arXiv:1001.4593.

[Al] P. Albers, *A Lagrangian Piunikhin-Salamon-Schwarz Morphism and Two comparison Homomorphisms in Floer homology*, Int. Math. Res. Not. IMRN 2008, no. 4, Art. ID rnm134, 56 pp.

[Ar] M. Artin, *Lectures on Deformation of Singularities,* Tata Institute of Fundamental Research lectures on Mathematics and Physics series 19, Anil D. Ved at Prabhat Printers, Bombay, 1976.

[Aur1] D. Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. 1 (2007), 51-91 (electronic), arXiv:0706.3207v2.

[Aur2] D. Auroux, *Special Lagrangian fibrations, wall-crossing, and mirror symmetry*, Surveys in differential geometry. Vol. XIII. Geometry, analysis, and algebraic geometry: 1–47, Surv. Differ. Geom., 13, Int. Press, Somerville, MA, 2009.

[BGS] W. Ballman, M. Gromov, V. Schroeder, Manifolds of Non-positive Curvature, Birkhauser, Boston, 1985

[Bar] S. Baranikov, *Semi-infinite Hodge structures and mirror symmetry for projective spaces*, preprint, arXiv:math/0010157.

[B1] V. Batyrev, *Quamtum cohomology rings of toric manifolds,* Journees de Geometrie Algebrique d'Orsay, 1992, Asterisque 218, 9-34.

[B2] V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties,* J. Alge. Geom. 3 (1994), no. 3, 493–535.

[Ber] B.G. Berkovich, Spectral Theory and Analytic Geometry over Non-Archimedean Fields. Mathematical Surveys and Monographs, 33. American Mathematical Society, Providence, RI, 1990.

[BC1] P. Biran and O. Cornea, *Quantum structures for Lagrangian submanifolds*, preprint, arXiv:0708.4221.

[BC2] P. Biran and O. Cornea, *Rigidity and uniruling for Lagrangian submanifolds*, Geometry and Topology 13 (2009), 2881–2991.

[BGR] S. Bosch, U. Güntzer and R. Remmert, Non-Archimedean Analysis. Asystematic Approach to Rigid Analytic Geometry, Grundlehren der Mathematischen Wissenschaften 261, Springer-Verlag, Berline, 1984.

[BT] R. Bott and L. W. Tu, Differential Forms in Algebraic Topology, Springer-Verlag, Berline,

[Cho1] C.-H. Cho, *Holomorphic discs, spin structures and the Floer cohomology of the Clifford torus*, Internat. Math. Res. Notices 35 (2004), 1803-1843.

[Cho2] C.-H. Cho, *Products of Floer cohomology of torus fibers in toric Fano manifolds*, Commun. Math. Phys. 260 (2005) 613–640, math.SG/0412414.

[Cho3] C.-H. Cho, *Non-displaceable Lagrangian submanifolds and Floer cohomology with non-unitary line bundle*, J. Geom. Phys. 58 (2008), 1465–1476, arXiv:0710.5454, arXiv:0710.5454.

[CL] C.-H. Cho and S. Lee, *Notes on Kontsevich - Soibelman's Theorem about Cyclic A_∞ algebras.* preprint, arXiv:1002.3653.

[CO] C.-H. Cho and Y.-G. Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, Asian J. Math. 10 (2006), 773-814.

[Con] A. Connes, Noncommutative Geometry. Academic Press, Inc., San Diego, CA, 1994. xiv+661 pp.

[Dub] B. Dubrovin, *Geometry of 2D topological field theories*, Lecture note in Math. 1620 (1996), 120 - 348.

[Ei] D. Eisenbud, Commutative Algebra with a View toward Algebraic Geometry, Graduate Text in Math. 150, Springer-Verlag, Berline, 1994.

[El] R. Elkik, *Algébrisation du module formel d'une singularité isolée.* Asterisque, 16 (1974) 133-144.

[Fl1] A. Floer, *Morse theory for Lagrangian intersections,* J. Differ. Geom. 28 (1988), 513-547.

[Fl2] A. Floer, *Cuplength estimates on Lagrangian intersections.* Comm. Pure Appl. Math. 42 (1989), no. 4, 335–356.

[Fu1] K. Fukaya, *Floer homology and mirror symmetry II*, Minimal Surfaces, Geometric Analysis and Symplectic Geometry (Baltimore, MD, 1999) Adv. Stud. Pure Math. 34, 31–127, 2002.

[Fu2] K. Fukaya, *Cyclic symmetry and adic convergence in Lagrangian Floer theory*, to appear in Kyoto J. Math., arXiv:0907.4219.

[Fu3] K. Fukaya, *Counting pseudo-holomorphic discs in Calabi-Yau 3 fold*, submitted, arXiv:0908.0148.

[FOOO1] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory-anomaly and obstruction,* Kyoto University preprint, 2000.

[FOOO2] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory-anomaly and obstruction,* expanded version of [FOOO1], 2006 & 2007.

[FOOO3] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory-anomaly and obstruction,* AMS/IP Studies in Advaneced Math. 46, International Press/ Amer. Math. Soc. (2009).

[FOOO4] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory on compact toric manifolds I,* Duke Math. J. 151 (2010), 23-174.

[FOOO5] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory on compact toric manifolds II: bulk deformation,* submitted, arXiv:0810.5774.

[FOOO6] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Anti-symplectic involution and Floer cohomology*, submitted, arXiv:0912.2646.

[FOOO7] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Toric degeneration and non-displaceable Lagrangian tori in* $S^2 \times S^2$, submitted, arXiv:1002.1660.

[FO] K. Fukaya and K. Ono, *Arnold conjecture and Gromov-Witten invariant*, Topology 38 (1999), no. 5, 933–1048.

[Ful] W. Fulton, Introduction to Toric Varieties, Annals of Math. Studies, 131, Princeton University Press, Princeton, 1993.

[Gi1] A. Giventa1, *Homological geometry and mirror symmetry*, in "Proceeding of the ICM-1994, Zürich, Basel, 1995, 472 - 480.

[Gi2] A. Givental, *A mirror theorem for toric complete intersections*, Topological field theory, primitive forms and related topics (Kyoto, 1996), 141–175, Progr. Math., 160, Birkhäuser Boston, MA, 1998.

[Goo] T. Goodwillie, *Cyclic homology, derivations, and the free loop space*, Topology 24 (1985), 187–215.

[GrPa] T. Graber and R. Pandharipande, *Localization of virtual classes*, Invent. Math. 135 (1999), 487-518

[Gro] M. Gross, *Mirror symmetry for* $\mathbb{P}^2$ *and tropical geometry*, preprint, arXiv:0903.1378.

[GPS] M. Gross, R. Pandharipande, and B. Siebert, *The tropical vertex*, preprint, arXiv:0902.0779.

[HL] A. Hamilton and A. Lazarev, *Cohomology theories for the homotopy algebras and noncommutative geometry*. preprint, arXiv:0707.3937.

[Ha] R. Hartshorne, Algebraic Geometry, Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.

[Ho] K. Hori, *Linear models in supersymmetric D-branes,* Proceedings of the KIAS conference - Mirror Symmetry and Symplectic Geometry- (Seoul, 2000), eds by K. Fukaya, Y.-G. Oh, K. Ono and G. Tian, World Sci., River Edge, New Jersey, 2001.

[HV] K. Hori and C. Vafa, *Mirror symmetry,* preprint, 2000, hep-th/0002222.

[MIRROR] K. Hori, S Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil and E. Zaslow, Mirror Symmetry, Clay Mathematics Monographs, Amer. Math. Soc., 2003

[Iri1] H. Iritani, *Convergence of quantum cohomology by quantum Lefschetz*, J. Reine Angew. Math. 610 (2007), 29–69.

[Iri2] H. Iritani, *Real and integral structures in quantum cohomology I: Toric orbifolds*, preprint, arXiv:0712.2204.

[Ka] C. Kassel *A Künneth Formula for the Cyclic cohomology of* $\mathbb{Z}_2$*-Graded Algebras*, Math. Ann. 275, 683–699 (1986).

[KKP] L. Katzarkov, M. Kontsevich and T. Pantev, *Hodge theoretic aspects of mirror symmetry*, in "From Hodge Theory to Integrabiliy and TQFT tt^* geometry, Proceeding Symposia in Pure Mathematics 78, (2008) 87-174.

[Ko1] M. Kontsevich, *Feynman diagram and low dimensional topology*, Proc. of the first European Congr. Math. Birkhäuser, Basel, (1994), 97–122.

[Ko2] M. Kontsevich, *Homological algebra of mirror symmetry,* ICM-1994 Proceedings, Zürich, Birkhäuser, 1995.

[KS1] M. Kontsevich and Y. Soibelman, *Homological mirror symmetry and torus fibration*, in: Symplectic Geometry and Mirror Symmetry, (Seoul, 2000), eds by K. Fukaya, Y.-G. Oh, K. Ono and G. Tian, pp 203-263, World Sci. River Edge, 2001.

[KS2] M. Kontsevich and Y. Soibelman, *Affine structures and non-archimedean analytic spaces*, in: The Unity of Mathematics (P. Etingof, V. Retakh, I. M. Singer, eds), pp 321-385, Progr. Math. 244, Birkhäuser 2006.

[Lo] J.L. Loday, Cyclic homology, Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1992.

[Ma] Y.I. Manin, Frobenius Manifolds, Quantum Cohomology, and Moduli Spaces, AMS Colloquium Publications, 47, American Mathematical Society, Providence, RI, 1999.

[Mat] H. Matsumura, Commutative Algebra, Benjamin, 1970.

[MT] D. McDuff and S. Tolman, *Topological properties of Hamiltonian circle actions*, Int. Math. Res. Pap. 2006, 72826, 1–77.

[Sab] C. Sabbah, *Universal unfolding of Laurent polynomials and tt^* structures*, in "From Hodge Theory to Integrabiliy and TQFT tt^* geometry", Proceeding Symposia in Pure Mathematics 78 (2008) 1 - 31.

[Sa] K. Saito, *Period mapping associated to a primitive form*, Publ. R.I.M.S. 19 (1983), 1231 - 1261.

[SaTa] K. Saito and A. Takahashi, *From primitive forms to Frobenius manifolds*, in "From Hodge Theory to Integrabiliy and TQFT tt^* geometry", Proceeding Symposia in Pure Mathematics 78 (2008) 31 - 48.

[MSa] M. Saito, *On the structure of Brieskorn lattices*, Ann. Institute Fourier (Grenoble) 39 (1989) 27 - 72.

[Se1] P. Seidel, *π_1 of symplectic automorphism groups and invertibles in quantum homology rings*, Geom. Funct. Anal. 7 (1997), no. 6, 1046–1095.

[Se2] P. Seidel *Graded Lagrangian submanifolds.* Bull. Soc. Math. France 128 (2000), 103–149.

[Se3] P. Seidel, *A biased view of symplectic cohomology*, preprint, arXiv:0704.2055.

[Se4] P. Seidel, *More on vanishing cycles and mutations*, in 'Symplectic Geometry and Mirror Symmetry' (Seoul, 2000), eds by K. Fukaya, Y.-G. Oh, K. Ono and G. Tian, pp 429–465, World Sci., River Edge, NJ, 2001.

[Se5] P. Seidel, *Symplectic homology as Hochschild homology.* Algebraic Geometry—Seattle 2005. Part 1, 415–434, Proc. Sympos. Pure Math., 80, Part 1, Amer. Math. Soc., Providence, RI, 2009.

[Ta] A. Takahashi, *Primitive forms, topological LG model coupled with gravity, and mirror symmetry*, preprint, arXiv:9802059.

Department of Mathematics, Kyoto University, Kyoto, Japan
E-mail address: `fukaya@math.kyoto-u.ac.jp`

Department of Mathematics, University of Wisconsin, Madison, WI, USA, & Department of Mathematics, POSTECH, Pohang, Korea
E-mail address: `oh@math.wisc.edu`

Graduate School of Mathematics, Nagoya University, Nagoya, Japan, & Korea Institute for Advanced Study, Seoul, Korea
E-mail address: `ohta@math.nagoya-u.ac.jp`

Department of Mathematics, Hokkaido University, Sapporo, Japan, & Korea Institute for Advanced Study, Seoul, Korea
E-mail address: `ono@math.sci.hokudai.ac.jp`